

Barcode : 2030020020207
Title - Indian Medical Plants
Author - Chopra R N
Language - English
Pages - 344
Publication Year - 1955
Barcode EAN.UCC-13



UNIVERSAL
LIBRARY

OU_174155


UNIVERSAL
LIBRARY

**A REVIEW OF WORK
ON
INDIAN MEDICINAL PLANTS**

Indian Council of Medical Research

SPECIAL REPORT SERIES

No. 30



सत्यमेव जयते

A REVIEW OF WORK
ON
INDIAN MEDICINAL PLANTS
(Including Indigenous Drugs & Poisonous Plants)

By

Bt.-Colonel R. N. CHOPRA

I. C. CHOPRA

NEW DELHI

1955

PRICE RS 3/-

FOREWORD

The use of indigenous drugs for the alleviation of human suffering is of considerable economic importance to our country. With this end in view the Indian Research Fund Association, now the Indian Council of Medical Research, has been sponsoring extensive researches in indigenous drugs for over two decades. The pioneering work in this field was done by Colonel Ram Nath Chopra at the School of Tropical Medicine, Calcutta and it may be said that we owe to him and to his students such knowledge as we today possess about the indigenous drugs of India. A review of the work done in this field was published by the Indian Research Fund Association in 1939. This review was greatly in demand. The Council therefore decided that it was essential to bring the subject up to date and requested Colonel Ram Nath Chopra to undertake the task. That he has done it very ably will be apparent from a perusal of the book. It gives in simple and non-technical language a brief account of the indigenous drugs so far investigated in India. The review should therefore be useful not only to practitioners of the indigenous systems of medicine in the country but also to scientists and to the lay public as well. It will indicate to the scientists which drugs require further investigation. The bibliography given at the end is a comprehensive one which, apart from its scientific value, will show the extent of interest which scientists both in India and abroad have taken in this subject.

Much yet remains to be done and many more drugs will have to be investigated scientifically by the application of modern methods of botanical, chemical, pharmacological and clinical research. I would, however, like to congratulate Colonel Ram Nath Chopra and his collaborators for their painstaking labour and for putting this material in such a concise and interesting form.

AMRIT KAUR,

Minister for Health, India.

September 16, 1954.

PREFACE

For many years past, the Indian Council of Medical Research, formerly Indian Research Fund Association, has given large grants to research workers all over India for the study of Indian Indigenous Drugs. In fact it was this body which was chiefly responsible for initiating work on this subject of great economic importance to India. The senior author was one of the first recipients of these grants when he was Professor of Pharmacology at the Calcutta School of Tropical Medicine and Medical College, Calcutta. As Professor at the former Institution, one of the duties laid down for him, was the study of Indian Indigenous Drugs. Financial resources from the School itself for this work were very limited then and it was the generous grants given by the Indian Council of Medical Research, extending over a period of more than three decades, both at the School and later at the Drugs Research Laboratory, Kashmir which enabled him and his co-workers to carry out screening studies of a large number of indigenous drugs. Considerable financial help was also received later from Indian Council of Agricultural Research and the Council of Scientific and Industrial Research.

In 1938 the senior author was asked to write a review of the work done on Indigenous Drugs under the Council upto that time, in simple non-technical language and a brochure was published in 1939. Large demand came from the public for this pamphlet and the edition was soon exhausted. In 1953 he was again requested to write a review of the work on Indian Indigenous Drugs, but this time it was not to be limited to work done under the Council but was also to include work done by other independent investigators. A thorough study of the whole literature on the subject was, therefore, undertaken by the present authors. It transpired that the work on Indigenous Drugs was being chiefly directed in the channel of investigation of Indian Medicinal Plants. Hence the change in the title of the present edition.

The general arrangement of the subject matter in the brochure of 1939 has been again adopted. Brief summaries of the work done on various drugs have been given in "telegraphic language". A comprehensive bibliography has been given, arranged subject wise and with full titles of papers. Lists of plants used in the treatment of various diseases in the indigenous systems of medicines have also been given after a careful analysis of the literature.

It is hoped that this review will be read by those interested in indigenous drugs and their practical use for alleviation of human ailments. As the descriptions are given in simple non-technical language and common vernacular names have been included, the present volume will be of interest not only to research workers but also to the public in general. The bibliography of nearly two thousand publications on the subject, collection of which has involved immense labour, will we trust, be useful to research workers and others.

In writing the present volume we wish to gratefully acknowledge the assistance we have received at every stage from Mr K L Handa who has helped in collecting references from literature and has checked up the chemical data. Messrs. L. D. Kapoor and B K Abrol have checked up botanical terms and synonymy and Dr Kartar Singh Pharmacological Assistant in Indigenous Drugs Enquiry has rendered valuable assistance

To the Governing Body of the Indian Council of Medical Research, its worthy President Rajkumari Amrit Kaur and its able Secretary Dr C G Pandit, the authors are grateful for the generous grants for work on Indigenous Drugs. These grants are being continued even now and are enabling the authors to carry on the work at the Drug Research Laboratory

Finances were also kindly provided by the Council for the preparation of the manuscript and the printing of this volume.

Drug Research Laboratory,
Jammu April, 1954

R. N. CHOPRA
I C. CHOPRA

C O N T E N T S

	PAGES
CHAPTER I	
INTRODUCTION . Climate and topography Features of Indian Vegetation Relationship to other floras ...	1
CHAPTER II	
MEDICINAL PLANTS OF INDIA Remedies used in Indigenous Medicine Collection of Vegetable drugs ...	3
CHAPTER III	
INDIAN INDIGENOUS DRUGS Main lines of basic research. Indigenous Drugs of India Historical review ...	6
CHAPTER IV	
BASIC REQUIREMENTS FOR STUDY OF INDIGENOUS DRUGS A Botanical Unit A Chemical Unit A Pharmacological Unit A Clinical Unit. ..	10
CHAPTER V	
SCOPE OF WORK ON INDIGENOUS DRUGS To make India self-supporting To discover remedies from the claims of Indian Medicine How to effect economy Indian Phar- macopoeia.	13
CHAPTER VI	
A RETROSPECT OF RESULTS ACHIEVED Value of negative results Summaries of work done on commonly used 183 drugs.	19
CHAPTER VII	
I TOXICOLOGICAL ASPECTS OF INDIAN PLANTS Crypto- gams (Flowerless Plants) Bacteria Algae Fungi— Parasitic on man and animals Those attacking food- stuffs Smuts Rusts Ergot Moulds. Mushrooms Poisonous and edible. Lichens Bryophyta. Pteridophyta. ...	52

CHAPTER VIII

II TOXICOLOGICAL ASPECTS OF PHANEROGAMS · (Flowering Plants) Plants poisonous to man and livestock Medicinal plants Food Poisons. List of 386 plants with a reference to their habitat, toxicological, insecticidal & piscicidal properties	59
---	----	-----	----	----

CHAPTER IX

MEDICINAL PLANTS USED IN INDIGENOUS MEDICINE Against Tuberculosis, Bowel Diseases (Dysentery & Cholera), Fevers Emenagogue & Abortifacient Irritants producing dermatitis Insecticidal and Piscicidal plants	...			96
--	-----	--	--	----

CHAPTER X

BIBLIOGRAPHY	122
SUPPLEMENT TO THE BIBLIOGRAPHY			...	254
APPENDIX OF SCIENTIFIC NAMES	I
APPENDIX OF VERNACULAR NAMES	39

A REVIEW OF WORK ON INDIAN MEDICINAL PLANTS

(including Indigenous Drugs and Poisonous Plants)

INTRODUCTION

The Indian Materia Medica, including the common household remedies at present includes about 2,000 different drugs. Many of these are in actual use in different parts of India. Of this number approximately two hundred are of mineral origin, about two hundred are of animal origin and the remainder are of vegetable origin. The mineral drugs are mostly compounds whose chemical composition is generally known and whose actions are in most cases not difficult to understand. The drugs of animal origin are a very varied assortment, few of which have been shown to have any rational usage. The study of indigenous drugs, therefore, chiefly means a study of Indian medicinal and allied plants in which there is a vast and very rich field for investigation.

Climate and topography The extreme variability that India presents in its meteorological and climatic conditions as also in its topographical features is perhaps unrivalled in the world. The tremendous range of variation of temperature from 126°F in summer in desert areas to minus 46°F in winter in Dras in Kashmir, and the prevalence of arctic conditions throughout the year in some of the Himalayan ranges are well known. The annual rainfall varies from 430 inches at Cherrapunji in the hills of Assam to less than 5 inches per annum in the desert areas of Rajputana. The air is saturated with moisture in the coastal districts and in the hills during the south-west monsoon and there are periods of practically zero relative humidity in the dry weather. These are some of the interesting contrasts presented by this vast country in which we live.

No less striking are its physical features. It has the most massive and the loftiest range of mountains in the world, the Himalayas. There are also lower hills and plateaus, extensive rich alluvial plains, sandy wastes and deserts, hills, streams, mighty rivers with their extensive deltaic and estuarine systems, numerous lakes, canals, ponds, tanks and extensive marshy tracts, sandy or rocky coasts spread over an area of more than a million and half square miles. The country is in fact an epitome of almost all climates, seasons, and soils of the world. Under these conditions it is not surprising that India possesses one of the richest and certainly the most varied flora of any area of similar size on the surface of the globe.

Features of Indian Vegetation Roughly speaking, one-fifth of the Indian sub-continent is occupied by forests of different kinds. There are ever-green forests occupying the west coast of the peninsula up to the ridges of the Western mountain chain, and the lower slopes of the

Note .— This work was initiated by the Indian Council of Medical Research formerly Indian Research Fund Association.

Eastern Himalayas, with their magnificent and majestic trees, the 'sky scrapers' many centuries old. A large part of the peninsula is covered by the deciduous type of the forests which are very valuable to the country. There are the dry forests of Rajputana and the Punjab which are also spread over a large area of Indian States. There is very little vegetation of any kind in the Great Indian Desert itself.

Many tidal creeks and backwaters along the coasts of India, chiefly the deltas of rivers, such as the Sunderbans are covered with another type of evergreen forest, the tidal or littoral, where many plants send up from their underground roots a number of aerial roots for respiration. The massive chain of the Himalayas with its tropical, temperate and alpine flora presents unique features. The eastern Himalayan flora differs remarkably from the western in several respects in addition to the disappearance of the tropical type as the extreme north-west is reached. It is in this chain that the highest limit of vegetable life on earth has been recorded. The tree limit in the Western Himalayas corresponds roughly with the snow-line, i.e. 12 000 ft. above sea level, above which a variety of beautifully coloured flowers flourish. Aquatic or semi-aquatic types of vegetation are found in the tanks, jhils, ponds, water courses and swamps all over India, particularly in Bengal. Lastly, there are areas in the plains, as well as in the hills, especially in the former, which have, through the influence of man in search of more and more agriculturally exploitable land, lost much of their original characters. Another feature of the Indian vegetation is that whereas in one season of the year, vast areas in the plains of India are the scene of plenty for man and animals, in another they become a dreary brown sunburnt waste where herbivores starve by the thousands and where it passes human understanding how any creature dependent on plant life can survive the dry season.

Relationship to other floras The Indian flora is closely related to that of some other countries, and it is interesting to remark that none of families of flowering plants is peculiar to this country. There is a very great preponderance of those genera and species which are also found in the adjacent countries. Malayan, Burmese and Chinese genera are specially well represented in Eastern India. Next in order come the European and Middle East flora which are particularly strongly represented in Western India. Of the typically European elements there are twice as many in the Western as in the Eastern Himalayas. Nearly 600 European genera are represented in India, many of them by a single species, and the Middle Eastern element is certainly, as is only to be expected, no less prominent. African, Australian and American elements follow these in decreasing proportions, the African flora being more noticeable in Western India. Recently there has been an influx of American species with their remarkable tendency to spread. Tibetan and Siberian floras only reach India in the alpine regions of the Himalayas, while Chinese and Japanese elements are strongly represented in its temperate belts. All these factors tend to make the Indian flora very rich and cosmopolitan and full of possibilities of economic importance.

CHAPTER II

MEDICINAL PLANTS OF INDIA

India abounds in all kinds of food plants, spices, perfumes, timber, fibres, gums, etc. which have been known all over the world from ancient times. There are more than about 700 important fodder plants including about 260 species of valuable fodder grasses. It is not surprising, therefore, that plants containing active and medicinal principles also grow abundantly within its bounds. More than 2,000 plants, out of a total of about 11,000 species found in India, are alleged to have medicinal properties of some description or other and have been enumerated in the literature of indigenous medicine. Nearly three-fourths of the drugs mentioned in the British and other Pharmacopoeias grow here in a state of nature and others can be easily grown. Indeed, this country is a veritable emporium of drugs. The families to which the larger numbers of medicinal plants belong are Leguminosæ, Compositæ, Labiataë, Euphorbiaceæ, Rubiaceæ, Rosaceæ, Gramineæ, Liliaceæ, Rutaceæ, Ranunculacæ, Umbelliferæ, Cucurbitaceæ, Solanaceæ, Apocynaceæ and Malvaceæ.

It stands to reason that all these 2,000 odd plants cannot possess the wonderful virtues attributed to them, but it is believed that there are some of these which might rightly deserve the reputation they have earned as cures. In order to determine what these were and what medicinal properties they possessed the study of these plants was begun in the early part of the last century.

The early studies were mainly confined to collection of available data from the literature of indigenous medicine as well as getting information about their popular use as household remedies by the people. A rough survey with regard to the occurrence of these plants in different parts of the country was undertaken by various workers and in some cases chemical analysis of a preliminary nature was also carried out. Modern medicine however, is intimately related to chemistry and detailed examinations of active principles of plants and other products form an essential part of it. Experimental work on the pharmacological side to determine the action of these active principles needs laboratories well equipped with modern apparatus and appliances which were not available in those days. Work on medicinal plants on modern scientific lines was thus started in the early part of this century with the main objective of making India self-sufficient and self-supporting by enabling her to utilize drugs produced in the country and by manufacturing them in a form suitable for administration. There are a number of drugs of established therapeutic value which are in use in the pharmacopoeias of different countries. Many of these grow wild and some in great profusion in different parts of India and a certain number are even cultivated. If these resources could be developed and utilized and the finished products manufactured, treatment of many diseases could be brought within the means of the Indian masses whose economic condition is unfortunately of a very low order.

A number of important drug plants extensively used by the medical profession, however, are neither found growing in a state of

nature nor have so far been cultivated in India. Cultivation of such plants is very important from an economic point of view, and scientific research in this direction, as is being carried out in other countries, would be very fruitful. It is a matter of regret that India is still importing quantities of crude drugs in spite of the fact that practically every conceivable pharmacopoeial drug can be grown within her bounds. The history of the cultivation of cinchona, eucalyptus, digitalis, etc. in India clearly shows that the cultivation of medicinal plants is pregnant with rich possibilities if taken up on proper lines. It is gratifying to note that in recent years a growing and increasing interest is now being taken in this direction.

Remedies used in Indigenous Medicine. The second objective has been to discover remedies from the claims of Ayurvedic, Tibbi and other indigenous resources suitable for employment by exponents of Western medicine. This is only possible if these are shown to have effective active principles in them. Since the period of decay and resuscitation of old systems of Indian medicine, knowledge of many of the effective remedies was lost while a number of uncertain ones crept in. Belief in their efficacy originates in some cases from the teachings of the ancient commentators and is based on clinical data. In others, however, there is no basis whatever. The object of research initiated by the Indian Council of Medical Research (formerly Indian Research Fund Association) was to determine which of these plants had the therapeutic efficacy claimed and which had not. Since this work was started about thirty years ago, several hundred medicinal plants have been partially screened and investigated by workers in this country. Some of these have been shown to be of great practical utility and have been actually brought into use. Many of the others were found to have effective medicinal properties and could be brought as cheaper substitutes for pharmacopoeial and Extra pharmacopoeial drugs. A large number of others were found to be entirely useless. Much, however, remains to be done and hundreds of plants with alleged medicinal properties, remain to be investigated.

Collection of drugs A few words may be said here with regard to the time of collection of drugs, for purposes of examination. This should be done at the proper time when the plant matures and the active principles are at their maximum. If this is not done even effective drugs may be found to have no activity. During the last twenty five years several drugs of Indian origin have assumed considerable importance from the point of view of foreign trade. Many firms of drug manufacturers in this country also use the locally produced raw materials for the manufacture of the finished products. It is a matter of very great concern, therefore, that the crude drugs collected locally are often not up to the required standard and this has resulted in considerable economic loss. Fortunately, this state of affairs is being gradually rectified. It must be stated however, that collection of drugs is not so simple a matter as most people seem to imagine. Collectors of medicinal drugs growing in a state of nature, and the present and prospective cultivators should bear in mind that there are certain factors which have to be considered in order to obtain standard medicinal products. There is a good deal of variation in the active principles in the different parts of a plant and in different seasons in the same part of the plant. Even the same part and at the same time of the year shows remarkable variations.

in the contents of its active principles For example, the young and the old leaves of a plant and unopened and opened flowers differ materially despite the fact that they are collected from the same plant and during the same season.

The soil also has an important influence on the active principles of plants It has often been observed that plants collected at the proper time, when the active principles have reached maturity and are at their maximum, give very effective results while the same plants when collected under other conditions are utterly useless Conditions for the collection of drugs in the case of plants under cultivation are more favourable and strict control over various factors can be exercised with greater ease than is the case with plants growing in a state of nature. It has, however, to be admitted that ideal conditions for the collection of even many of the common and important medicinal plants are not known with precision and research is urgently needed to determine the time when the active principles are at their maximum under the environments existing in this country, as was done by the Dutch in Java in the case of cinchona If this could be systematically carried out, India could supply itself and other countries with medicinal 'herbs'

CHAPTER III

INDIAN INDIGENOUS DRUGS

Main lines of basic research Having given a general account of the potentialities of this country with regard to medicinal plants, we will now briefly describe some of the important lines of basic research taken up by workers in this connection. The first of these is study of Indigenous Drugs of India.

Indigenous Drugs of India It is desirable to point out at the outset that the term 'Indigenous Drugs' has been used, for the purposes of research in its widest sense so as to include within its scope not merely those drugs which were originally the natives of India, but also those which have been introduced from outside and have become completely naturalised. Drugs which are cultivated in India, whether used in the indigenous system of medicine or in the pharmacopoeias of various western countries, have also been brought within the purview of this investigation. The term also includes locally manufactured preparations of mineral and animal origin which are used in medicine as practised in India.

Historical The study of Indian indigenous drugs was first begun in the early part of the last century and it was then confined chiefly to the collection of available information with regard to various medicinal plants growing in different parts of the country. The earliest contributions were from the writings of Sir William Jones whose memoir entitled "*Botanical Observations on Select Plants*", is well known. This was followed in 1810 by John Fleming's "*Catalogue of Medicinal Plants*", Ainslie's "*Materia Medica of Hindustan*" in 1813, and Roxburgh's "*Flora Indica*" in 1820. In 1844 O. Shaughnessy published his "*Bengal Pharmacopoeia*" which was the first book of its kind which dealt exclusively with the properties and uses of the medicinal plants used in Bengal. In 1868 a "*Pharmacopoeia of India*" was published under the able editorship of Waring. It signaled a new epoch in establishing and recording the value of indigenous medicinal products on modern lines. The more important drugs were officially recognised with a view to their eventual adoption in the British Pharmacopoeia. As a large number of the drugs, especially those in local use by practitioners and as household remedies, were not included in this work, Mohideen Sheriff published his "*Supplement to the Pharmacopoeia*" in the year 1869. He is also the author of "*Materia Medica of Madras*" which was edited and published after his death by Hooper. U. C. Dutt's translation of Sanskrit *Materia Medica* brought into prominence the drugs used in the old Hindu Medicine even now largely practised in India. Then Fluckiger's and Hanbury's "*Pharmacographia and Materia Medica of Western India*" 1883 was published. These were followed by the publication of that very comprehensive book on the Indian Medicinal plants the "*Pharmacographia Indica*" in 1885 under the joint editorship of Warden and Hooper. This treatise contains a mass of information regarding the uses of the indigenous *materia medica* in the Eastern and Western medicine. The most elaborate work of all is "*A Dictionary of the Economic Products of India*" published in 1895 by Sir George Watt, the Reporter on the Economic Products to the Government of India. This monumental work, compiled with the help of a large team of workers, refers

to all the previous work on medicinal plants and other plants of economic importance. Its pages are full of information of every description regarding the use of different barks, roots, flowers, leaves and woods for different medicinal and other purposes. Works published still later such as Kanai Lal Dey's "*Indigenous Drugs of India*" and Kirtikar and Basu's "*Indian Medicinal Plants*" are largely summaries and compilations from the above mentioned literature in a more systematised and elaborated form. In the last named book, plates illustrating various important medicinal herbs are given which greatly help the reader in differentiating them from plants with which they are apt to be confused.

The literature mentioned above is very valuable, as it contains not only information from Ayurvedic and Tibbi sources, but also gives the results of personal observations and experiences of some of the writers. There is no doubt that a considerable amount of botanical investigation into the scientific names of many medicinal plants was accomplished. More, however, remained to be done in the case of a large number of plants to clear up many points with respect to their exact botanical sources. Many drugs that had escaped the previous investigators require to be explored in all their details.

Admirable as all these attempts were, the pharmacology of most of the indigenous remedies remained an unexplored field till recent years. The reason for this is not far to seek. Investigations of this nature require a considerable outlay of money in the form of well-equipped chemical and pharmacological laboratories, while a liberal staff of competent chemists and pharmacologists is another essential prerequisite. Medicine we have observed is now intimately related to chemistry, and the ultimate solution of most problems, whether physiological or biological, rests on some physical or chemical basis. This is forcibly presented to us in the study of the action of drugs. The importance of the cooperation of chemists at every stage of research work can only be realised by the workers themselves. If satisfactory results have to be achieved and if the work is to be carried out on the same standard as in other civilised countries, the cooperation of competent chemists with experience is the first essential.

It must be pointed out that the time and labour required to work out the chemical composition of a single drug are enormous. This may be judged from the fact that it would take an experienced chemist several months, perhaps a year or more, to isolate in a pure state and roughly describe the nature of the different chemical constituents of a single crude drug. The determination of the chemical constitution of the active principles concerned would take a considerably longer time. The isolation of a sufficient quantity of the active principles and the testing of them pharmacologically would occupy many months. The magnitude of the task of working out all the drugs used in the indigenous systems of medicine in detail transcends all imagination. There is such an enormous scope for research in this field, and so little has been done, that it is impossible for any one individual or any one institution to cope with it adequately. The cooperation and intimate association of a large number of sincere and devoted workers of ability is needed to find the truth. This is now being gradually done. Chairs in Pharmacology have been founded by the various Universities and Medical Colleges and facilities for research work on modern scientific lines are now being made increasingly more available.

As the action of these drugs or their active principles can only be established by a careful chemical, pharmacological and clinical study, the investigation in all the three aspects should be carried on side by side. The experimental work on the pharmacological side can be done only in laboratories well equipped with all modern appliances. The first laboratory of its kind established in this country was at the School of Tropical Medicine, Calcutta in 1921. One of the main duties of the Professor of Pharmacology laid down was investigation of the indigenous drugs on scientific lines. The chemical department of this institution had a small team of chemists who worked out the chemical composition of drugs, isolated the active principles and handed them to the pharmacologist for determination of their action on the animal organism. The clinical testing of the drug was made possible by the Carmichael Hospital for Tropical Diseases a research hospital attached to this institution. In this way it was found possible to go through a number of drugs in all the varied phases of their investigation, i.e. from the isolation of their active principles to the testing of their action on animals and finally to the making of suitable preparations for trial on patients, and for recording the results of therapeutic trials.

The staff of the Departments of Pharmacology and Chemistry of this institution was only a nucleus to start this work. It was wholly inadequate to deal with such an immense task in addition to its ordinary duties of post-graduate teaching. Although a modest beginning was made in 1921 when the School started functioning, it was not really till 1926, when the Indian Research Fund Association, now known as the Indian Council of Medical Research, gave a grant to develop this work. Investigations were then taken up in right earnest and on proper systematic lines. During the years that followed, the importance of this work was appreciated with greater emphasis and the grants were considerably increased. It was realised by this body that this was the only effective reply to attempts at the wholesale revival of the indigenous system of medicine which were then in evidence. In 1935 the Imperial Council of Agricultural Research (now Indian Council of Agricultural Research), appreciating the importance of basic research of this type which was being done gave a grant for the investigation of the closely allied group of Poisonous Plants and Food Poisons of India which are such a menace to man and live-stock in this country. By this grant the research facilities for work on indigenous drugs were also further augmented and certain deficiencies which existed in the technical staff were made good. An efficient Botanical Unit was thus provided, the want of which had been very keenly felt in connection with the research work on medicinal plants.

With all these facilities made available it was possible to undertake the study of a large number of drugs in all the varied phases of their investigation i.e., from the isolation of their active principles to the testing of their action on animals and finally making suitable preparations for trials on patients and for recording results of therapeutic trials.

During the three decades that have followed, the research work on indigenous drugs has received considerable encouragement and has made satisfactory progress. The example of the Indian Council of Medical Research and Indian Council of Agricultural Research was followed by the Council of Scientific & Industrial Research. This Council

gave very generous grants to various Medical Institutions and other research bodies for this work. It also established in 1950, the Central Drug Research Institute at Lucknow as one of the eleven major National Laboratories of India. One whole division of this great Institution is devoted entirely to the study of Indian Indigenous Drugs. With the dawn of Independence, therefore, this research has been put on a sound and firm basis. Much has been accomplished, in the study of Indian Medicinal Plants on systematic and scientific lines, though much more remains to be done. A brief review of this work will be found in the following pages.

It is obvious from what has been stated above that the Indian Council of Medical Research with its predecessor the Indian Research Fund Association were the pioneers in encouraging the study of Indian Indigenous Drugs on scientific lines. It was the example of this body which stimulated interest on this important subject of vital importance to the country.

CHAPTER IV

BASIC REQUIREMENTS FOR STUDY OF INDIGENOUS DRUGS

We have said above that detailed study of Indigenous Drugs is not a one man job. It needs close collaboration and association of scientific workers in different allied subjects. The prerequisites for the work to be done properly on scientific lines are —

1. A Botanical Unit This Unit should consist of experienced botanists and technicians. They help not only in the collection and identification of genuine drugs, but also in carrying out a survey of the medicinal and poisonous plants growing in different parts of the country. The importance of this section must not be under-rated, as on the correct identification of the drugs depends the utility and authenticity of the whole work. In the early stages, when expert botanical help was not available, many mistakes were made in connection with the identity of medicinal plants, thereby causing much confusion and waste of effort. An extensive herbarium of these plants is being gradually built up by our botanists. This is a unique collection of its kind in India, and is rendering the correct identification of medicinal plants for any trained worker an easy matter.

2 A Chemical Unit This should consist of a band of expert research chemists with special experience in plant chemistry. They should be trained for work in different aspects of plant chemistry involving extraction and identification of active principles, such as alkaloids, glucosides, essential oils, neutral principles, antibiotics etc. The medicinal or poisonous plants sent by different forest officers or collected by the botanists, are first dealt with in this department and an attempt is made to determine the active constituents that may be contained in them. The preliminary chemical examination, the isolation of the different constituents which may have pharmacological action, their purification and elementary analysis all require considerable time, patience and skill. It may be stated here, that the isolation of the active principles of plant constituents is not an easy task of a routine nature, as may perhaps be imagined. Each plant has its own peculiar problems and difficulties which have to be solved. This can only be done by chemists with considerable practical experience and technical skill. Besides being engaged in the isolation and identification of active principles of plants, the chemical section is also required to assist in chemical and biochemical assays of different drugs, in the preparation of galenicals, extracts, etc. for purposes of clinical trial in the hospitals. The work involved, therefore, is of a varied and comprehensive character. Efficient teams of workers, have now been brought into existence after many years of training, and the analytical work is being considerably facilitated as more experience is being gained.

3 A Pharmacological Unit. This is composed, preferably of medical men especially trained in the technique of experimental pharmacology, who work under the guidance of a pharmacologist with large practical experience of this type of work. They test the different preparations or the purified active constituents isolated by the chemists by properly planned animal experimentation. They also carry out biological assays, test the toxicity of drugs and suggest suitable doses for therapeutic trials. Here also experience assists a great deal in assessing the

value of the reactions produced in animal tissues by drugs under investigation and their practical application to therapeutics in man

4 A Clinical Unit. This section carries out testing of drugs on patients in the out-patients department and in the hospital. This work is carried out under the direct guidance of experienced pharmacologists and clinicians. It has often happened that drugs have been tried in a haphazard fashion by medical practitioners and wrong conclusions have been drawn. The proper clinical evaluation of these drugs needs a well equipped hospital with a specially trained house staff of medical men and nurses, in which patients can be kept and watched for weeks and months if necessary. The clinician in charge of such investigation should also have experience of this type of work which needs considerable patience and sound judgment.


The existence of such teams all working in close cooperation, makes it possible to carry out this type of work on proper systematic lines. Research work of this nature started in institutions or university laboratories is often cut short at an early stage as the workers have to depend on institutions such as Hospitals for trials. Again some of the universities and colleges do not possess properly equipped laboratories for pharmacological studies. The procurement of all these facilities is expensive and make this work difficult. If such organisations as Indian Council of Medical Research, Indian Council of Agricultural Research and Council of Scientific & Industrial Research had not come forward with generous grants, it would not have been possible to make any headway.

It will thus be seen that from the stage of empirical knowledge available of a crude drug to bring it to its rational usage in modern medicine, it must pass through the following stages —


- (1) Botanical identification
- (2) Chemical examination and isolation of active principles
- (3) Determination of their pharmacological action and toxic effects
- (4) Chemotherapeutic studies in animals.
- (5) Clinical trials in man

In case of well-known drugs of established value in indigenous medicine, it may some time be convenient to start with clinical trials first and proceed to analysis and laboratory tests afterwards. A schematic idea of the work is given below —

PROCEDURE I



PROCEDURE 2



In whatever way the investigation is carried out on an indigenous drug, the work is time-consuming and can only be done by team-work of several groups of scientists each expert in his own field. The work should be co-ordinated by a medical scientist of experience who can draw logical conclusions from the data obtained. Haphazard approaches to this work have led to wrong conclusions and much confusion.

CHAPTER V

SCOPE OF WORK ON INDIGENOUS DRUGS

Work on indigenous drugs on modern scientific lines was initiated by the senior author when he was appointed Professor of Pharmacology at the School of Tropical Medicine, Calcutta with the following main objects in view :

1. To make India self-supporting by enabling her to utilise the drugs produced in the country, by manufacturing them in a form suitable for administration
- 2 To discover remedies from the claims of Ayurvedic, Tibbi and other indigenous sources suitable to be employed by the exponents of Western medicine
- 3 To discover the means of effecting economy, so that these remedies might fall within the means of the great masses in India whose economic condition is very low.
- 4 To prepare an Indian Pharmacopoeia including indigenous remedies of value

(I) TO MAKE INDIA SELF-SUPPORTING SO FAR AS DRUGS AND THEIR PREPARATIONS ARE CONCERNED

Pharmacopoeial and Allied Drugs The first proposition is likely to lead to great results, because a large number of drugs which grow in this country are known both to Eastern and Western medicine and the properties and actions in many cases are also known. The research here has been diverted into two main channels. Firstly, there are many drugs of established therapeutic value which are in use in the pharmacopoeias of different countries. The majority of these grow wild and in abundance in many parts of India and a certain number are even cultivated. Some of these are collected and exported, though only a small fraction of the quantity produced, to foreign countries and come back to us in the form of standardised pharmaceutical preparations and active principles in pure condition, probably at a price many times that of the original crude product. A host of others grow, mature and eventually die without being put to any practical use whatsoever. There are numerous examples which have been dealt with in detail elsewhere. A brief description of those investigated is given hereafter in this review with short notes with regard to the work done.

Besides these, there are a number of pharmacopoeial drugs which are widely used by the medical profession, but which do not naturally grow in this country. They thrive, however, when they are cultivated under proper conditions in suitable parts of the country. Examples of such drugs are numerous but a few of the important ones such as digitalis, ipecacuanha, eucalyptus, cinchona, jalap, etc., may be cited. They were introduced into India many years ago and are doing well. On account of the great demand for these drugs in medical practice, their production

in this country is of great economic importance. India, we have stated, possesses most wonderful variability so far as the temperature and general climatic conditions are concerned, and every conceivable drug ranging from those growing in the hottest tropical and damp climates to those growing in dry, temperate and very cold climates can be grown and acclimatised in some part or other. From the geological point of view also every grade of soil from alluvial deposits to hard rocky formation and sandy deserts are met with. Professor Greenish of the London School of Pharmacy rightly said, "India owing to the remarkable variations she possesses of climate, altitude and soil, is in a position to produce successfully every variety of medicinal herb required by Europe".

It should be remembered, however, that the soil, the season and the gathering time are some of the important variable factors with plants, and it can hardly be expected that the amount of active constituents would be constant under all conditions. In some cases the quality is good and constant, but in the majority of instances the percentage composition of active principles has to be determined by careful methods of chemical and biological assay, to show that these remedies, growing in a state of nature, are as good in quality as those required by the standards laid down in the Pharmacopoeias. If they do not come up to the required standard, the best method of bringing them into general use by improving the quality of the active principles by suitable cultivation, in parts of the country where this can be done economically, has to be determined.

The work done so far in connection with these two groups of plants, has been responsible for bringing into use a large number of crude drugs, which were formerly imported from foreign countries by the drug manufacturing industry, by showing that the active principles of many of the indigenous plants came up to the standards laid down by the pharmacopœia.

Secondly, a large number of plants grow in India which, though not exactly the same, have properties and actions similar to the imported and often expensive remedies and would form excellent substitutes. Not infrequently there are some closely allied species which are pharmacologically just as active. The properties of these plants are being worked out on scientific lines and they are being brought into use. Some examples may be cited here. *Colchicum luteum* growing on the slopes of the western temperate Himalayas has been shown to be an excellent substitute for the official *C. autumnale*. *Scilla indica* grows extensively on the sea-coast and on the drier hills of the lower Himalayas and the Salt Range and is being used as a substitute for *S. maritima*. *Ferula narthex*, growing in Kashmir, yields a gum resin resembling asafœtida. The properties of *Picrosma quassioides* and *Gentiana kurroo* resemble those of *Picrosma excelsa* and *Gentiana lutea* which are official in the British Pharmacopœia. Many other examples could be cited.

In both these groups there is a large field for research and development and considerable work has been done in connection with recognition of their active ingredients, determination of their percentage composition and establishing their pharmacological action. Pharmaceutical preparations are now manufactured from these indigenous products to the immense advantage of the country.

(2) TO DISCOVER REMEDIES FROM THE CLAIMS OF AYURVEDIC, TIBBI, AND OTHER INDIGENOUS SOURCES.

The Drugs used in the Indigenous Medicine. The second proposition of popularising and introducing new drugs to Western medicine is a more difficult one. It is believed that out of the very large number of drugs used in the indigenous systems of medicine for centuries past, and still in use, there must be some at least which might deserve the reputation they have earned as cures. Since the period of decay and recompilation of old Indian systems of medicine, many of the effective remedies have been lost and a number of uncertain ones have come in. The result is that in the indigenous systems, at the present time, almost every plant and shrub growing in the country has ascribed to it some medicinal virtue. These beliefs in some cases originate from the teachings of the ancient commentators and are based on clinical data, but in others have no foundation whatever. Their introduction was empirical and often a drug was used simply because a single case appeared to have derived some benefit from it. In this way remedies have multiplied without proof but by belief and as they hail from all parts of India, no one seems to have a correct notion about their uses and properties. The employment of a large number of them would thus appear to have been based on empirical evidence handed down from generation to generation. A thorough and complete research into all these drugs would constitute the life long work of innumerable chemists, pharmacologists and physicians. For practical purposes the method adopted by workers has been to make use of the experience of Vaidys, Hakims and others, and to take up for investigation those drugs which have a great local reputation before touching the less known remedies. Besides, many of these drugs have been clinically tried by medical men practising Western medicine, who have expressed their opinion regarding their efficacy, this has also been helpful in the selection of drugs to be investigated. In this way a large number of commonly used drugs have been scrutinised.

(3). HOW TO EFFECT ECONOMY AND BRING THE TREATMENT WITHIN THE MEANS OF THE MASSES.

The third proposition relates to the devising of expedients for effecting economy, so that these remedies may reach the masses. This is only possible if the price of the drugs can be considerably reduced, for in a poor country like India, there are millions of people who cannot afford any kind of treatment, whether cheap or expensive, and have consequently to depend upon charitable medical relief institutions. The cost of drugs is so heavy that most of these institutions, which have only a limited annual budget for drugs, are not able to cope with the demand for such common and essential drugs as quinine, castor oil, magnesia, etc., to say nothing of the expensive medicines which are sometimes necessary and even indispensable.

The only way in which drugs can be cheapened and brought within the means of the masses is to utilise the local resources and substitute the indigenous products for the more expensive imported preparations of Western medicine. This can be done by encouraging the production, collection and manufacture of the local materia medica by preparing pharmaceutical preparations in a systematic manner. By local production and substitution of equally potent drugs of Indian origin for the imported drugs, the cost of treatment can be considerably reduced.

We have already made a reference to some of these remedies and the possibilities of their development. Their active principles are now being isolated, and standardised preparations such as tinctures, extracts, powders, etc., are being prepared by a number of reliable manufacturers. If this aspect is further developed it will be possible to affect large scale economy.

It is a matter of satisfaction to note that a great deal has been done towards the development of a drug manufacturing industry in India, and towards encouraging the use of raw material produced in the country for the preparation of galenical and other preparations. The price of preparations has already come down considerably during recent years and by further development of this industry it is hoped that there will be further reduction in prices. The factors which were militating against the bringing down of the prices of medicinal preparations to the level of the low economic conditions of the masses were fully investigated by the Drugs Enquiry Committee (1930-31). These were firstly, the excise duty on alcohol used for bonafide medicinal purposes by the manufacturers, and secondly the high railway freight for purposes of transportation of drugs from one part of the country to another. Attention is being given to all these and drug industry in India is being put on a sound basis.

(4). INDIAN PHARMACOPOEIA.

The fourth proposition of preparing a Pharmacopoeia is by no means an easy one. A national Pharmacopoeia is primarily meant to meet the claims and satisfy the needs of a particular group of physicians at a particular time. There must exist, and there does exist, a great difference not only between the Pharmacopoeias of various countries, but also between various editions of the same Pharmacopoeia.

The object of a Pharmacopoeia is, in the words of the founders of the United States Pharmacopoeia, 1820, "to select from among substances which have medicinal power those the utility of which is most fully established and best understood, and to form from them preparations and compositions in which their powers may be exerted to the greatest advantage."

The modern Pharmacopoeia is above all a book of standards. Its fundamental object and scope is, "to provide standards for the drugs and medicines of therapeutic usefulness or pharmaceutical necessity sufficiently used in medical practice; to lay down tests for the identity, quality, and purity, to insure, as far as possible, uniformity in physical properties and active constituents." In other words, usage, rational usage and scientific usage are the basis of judgement.

The evidence placed before the Drug Enquiry Committee 1930-31 showed that India should have an official publication which would record what she recognizes as a trustworthy and approved *materia medica* upon which can be established modern food and drug acts, poison laws, systems of taxation, and the modernization of legal medicine. Her best scientific men should be enlisted to shoulder the responsibility of working out India's own standards and to develop the necessary analytical laboratories, biological institutes and pharmaceutical schools. It involves not only the development of machinery for the administration of laws based upon official standards, it means also finding an authoritative standard

which every doctor and pharmacist in India will hail as a real guide in every-day work

A consideration of what is said above will show that although the preparation of an Indian Pharmacopoeia is very desirable, a very large amount of work must necessarily be done before such a work can be prepared. The Drug Enquiry Committee 1930-31 carefully considered the question of the compilation of an Indian Pharmacopoeia and thought that the utility of such a work for the country cannot be overrated.

The pharmacopoeia which is in view ought to include the therapeutically active substances and, to find admission to it, a drug must be of known composition, of definite pharmacological action, and of well-established therapeutic use, and fully investigated for its toxicity and necessary standard for determining a safe maximum dose, with a chemical or biological standard. The large mass which do not satisfy this condition should be left out. Necessary tests have to be developed for the protection of doctor, pharmacist, and patient. India ought to set a standard of strength and purity for the material which is to appear on her markets.

This work should be on the lines of the British and the United States Pharmacopoeias including only drugs of known composition, of definite pharmacological action, of well-established therapeutic properties, with known toxicity and the necessary standards of chemical and biological assay for determining the safe maximum doses.

The work which is being done on the Indian Indigenous Drugs is exactly the type of work which will eventually lead to the preparation of an Indian Pharmacopoeia. But it is obvious that much more will have to be done before such an authoritative publication can be prepared. The Drug Control Laboratory at Calcutta and later the Central Drug Research Institute, Lucknow have helped greatly towards evolving standards suitable for this country. A large number of drugs commonly used have been fully worked out, and the conditions above stated are being satisfied with regard to them.

It is a matter of great satisfaction to note that work on these drugs has sufficiently advanced to lay the foundation of an *Indian Pharmacopoeia*. As a first step towards it an *Indian Pharmacopoeia List* was prepared by the Committee set up by the Government of India under the auspices of Drugs Technical Advisory Board in 1947. This was published in 1949 and is an authoritative document laying down standards for Indian climatic and other conditions for drugs sufficiently worked up to be included in the List. The Health Ministry then set up a Permanent Pharmacopoeia Committee. This Committee has been at work since and with the help of number of sub-committees composed of experts from all parts of India has now practically completed the labour of compiling the first edition of Indian Pharmacopoeia which will be published in the near future. The pioneering work done in connection with the preparation of Indian Pharmacopoeia by the Indian Council of Medical Research (formerly Indian Research Fund Association) cannot be overrated. As early as 1926 this body appreciated the importance of the study of Indigenous Drugs on scientific lines as a preliminary to compilation of an Indian Pharmacopoeia and started financing a number of investigations in this

connection with the object of working out of the indigenous drugs on scientific lines .

Recently an *Indian Pharmaceutical Codex* has been published under the auspices of Council of Scientific & Industrial Research by Dr B. Mukerji, Director, Central Drug Research Institute, Lucknow, which gives about 200 monographs and a similar number of formulary giving pharmacognostic, chemical, pharmacological and therapeutic data on indigenous drugs with their preparation and dosage for the medical and pharmaceutical professions.

CHAPTER VI

A RETROSPECT OF RESULTS ACHIEVED

During the period of three or four decades since this work was started, much has been accomplished though only the fringe of this vast problem has yet been touched. A number of important medicinal plants commonly used by the Kabirajes, Hakims and as popular household remedies by the people have been investigated on modern scientific lines. The chemical composition of many plants has been determined, the pharmacological action of the active principles of a large number has been worked out by animal experimentation, and finally suitable preparations made from a number of drugs have been tested on patients in properly equipped hospitals. It is only by such thorough investigations that the real merits of these drugs can be proved and a demand created for them not only in India but in other parts of the world. This laborious work has brought into prominence the merits and qualities of certain drugs and it has been shown that they may prove to be valuable additions to the armamentarium of the medical man to relieve the sufferings of humanity, if brought into general use. Such drugs unfortunately are not many. A few examples may be cited here. *Holarrhena antidysenterica* (Kurchi) has proved useful in the treatment of amoebic dysentery especially the chronic form, *Rauwolfia serpentina* has proved to be an effective sedative which brings down the blood pressure particularly the diastolic. This plant has attracted much attention of the Pharmacologists and clinicians in Europe and America. There are a number of others. *Plantago ovata*—Isabgul is an excellent sedative in irritative conditions of the guts, *Adhatoda vasica* is a good expectorant; *Psoralea corylifolia* is useful in leucoderma, *Cissampelos pariera* is likely to be a useful smooth muscle relaxant, *Deamia extensa* as smooth muscle stimulant.

A large number of those examined showed activity more or less the same as drugs already possessed by the pharmacopoeias. These can be brought into use as cheaper indigenous substitutes. Many others were effective enough to be used in the treatment of common minor ailments which form about 80% of total morbidity in this country. A fourth group of drugs which though used in indigenous medicine were found to have little or no activity as judged by our standards. Many drugs of questionable value and doubtful utility crept into the indigenous systems during the period of decay.

Full details of work done in this connection upto the end of 1953 will be found in the 2nd Edition of "*The Indigenous drugs of India*" which is a compendium of information with the regard to the whole subject of indigenous drugs and constitutes a leading work of reference. A Glossary of Indian Medicinal Plants is in the course of publication by the Council of Scientific & Industrial Research. In this book which has involved immense labour, brief reference has been made to scientific name of plants, their common vernacular names, the active principles contained,

Abbreviations used in description of Plants.

Afgh	..	Afghanistan
B	...	Bengali
Baluchi	...	Baluchistan
Baz	..	Bazar
Bhutia.	...	Bhutan and other Himalayan countries
Bo	...	Bombay
B P	...	British Pharmacopoeia
B P C	...	British Pharmaceutical Codex
Burm	.	Burmese
Dec	..	Deccan
H	..	Hindi
I P	...	Indian Pharmacopoeia
I P C.	...	Indian Pharmaceutical Codex
I P L.		Indian Pharmacopoeial List
K	..	Kanarese
Kash	...	Kashmiri
Kumaon	..	Kumaon Hills
M	...	Malayalam
Nep	...	Nepalese
P	..	Punjabi
Pers	..	Persian
S		Sanskrit
Singh	.	Singhalese
Santh.		Santhal
Vern		Vernacular

Abroma augusta Linn Devils cotton (Vern —H & B Ulatkambal
Bo —Olaktambol) I P C

Grows wild in Uttar Pradesh, Sikkim, Khasia Hills and Assam. Both root and bark are used as emmenagogue in menstrual disorders and as uterine tonic. Fresh viscid sap is used in dysmenorrhea in doses of 30 grains a day. An alkaloid 0.01%, a fixed oil, resins and water soluble basis have been found. Preliminary investigations of the alkaloid and water soluble bases revealed no remarkable pharmacological activity. No proper clinical trials have been carried out.

Abrus precatorius Linn. (Indian or wild Liquorice root) (Vern —S & Bo —Gunja, H —Gaungchi, B —Kunch, M —Gundumani)

Grows wild in plains of India and Ceylon and along the Himalayas up to altitudes of 3,000 ft. Seeds are used as tonic and aphrodisiac. Roots and leaves contain glycyrrhizin, an alkaloid abrine, a glycoside abralin and a fatty oil. Preliminary investigations of the alkaloid revealed its marked poisonous properties. No systematic study has been carried out with regard to its therapeutic efficacy.

Aconitum (Vern —Bachnag)

A paste prepared from the root is used as a remedy for neuralgia and other painful affections. The drug is used in the treatment of fever, rheumatism, cough, asthma and snake-bite.

Aconitum balfourii Stapf (Vern -Nep -Gobari, Darmiya gobriya)

It is found in sub-alpine Himalayas from Gharwal to Nepal at altitudes of 12,000 ft to 14,000 ft. Root contains total alkaloids 1.2% of which pseudo-acontine is 0.4%.

Aconitum chasmanthum Stapf. Indian Nepellus (Vern -Mohri, Kash -Banbalnag). I P L & I P C

Grows in Western Himalayas of Kashmir at altitudes of 7,000 ft to 12,000 ft Root contains 4.3 per cent of total alkaloids, aconitine being the principal

Aconitum deinorrhizum Stapf (Vern -Safed bikh, Mohra, Maurabikh)

It is found throughout the Central Himalayas from Kumaon to Nepal and Upper Bushahr in Himachal Pradesh Roots contain total alkaloids 0.9 per cent of which pseudo-aconitine is 0.4 per cent.

Aconitum ferox Wall (Vern -S -Visha, H & B -Bish, Bo -Vachnag, M -Vashanavi)

It is found in Northern Himalayas, Nepal and Kashmir. Root is used as a remedy for neuralgia and rheumatism and as a sedative, antipyretic and diaphoretic

Aconitum heterophyllum Wall (Vern -Atis, S -Ativisha, H -Atis, B -Ataicha, M -Ati vadayam)

It grows in sub-alpine and alpine zones of the Himalayas It is used as a bitter tonic and febrifuge in the treatment of fevers, diarrhoea, dyspepsia, cough and as an aphrodisiac Roots contain alkaloids atisine 0.4 per cent, heteratisine and hetisin

Aconitum laciniatum Stapf (Vern - Kalo bikhmo),

Grows in the sub alpine and alpine Himalayas of Sikkim and Tibet at altitudes of 10,000 to 14,000 ft.

Aconitum spicatum Stapf (Vern -Bikh, Kalo bikhoma donghi)

Grows in Nepal, Sikkim, Chumbi at altitudes of 10,000 to 12,000 ft It is used as a poison Roots contain 0.4 per cent of alkaloid named bikhaconitine

Aconitum violaceum Jacq (Vern -Tilia Kachnag)

Found in alpine zones of the Himalayas from Gilgit to Kumaon at altitudes of 10,000 to 15,000 ft The alkaloid aconitine is a cardiac irritant Locally it produces tingling followed by numbness Internally it slows the heart.

Acorus calamus Linn (The sweet flag) (Vern -H & B -Bach, Bo -Vaj, M.-Vashambu). I P C

Native of Europe and North America, but cultivated in India and Burma at altitudes of 3,000 to 6,000 ft Common in Manipur and Naga Hills. The rhizome is used as emetic, antispasmodic, carminative and expectorant. It is an old remedy against chronic, diarrhoeas Dried rhizome contains 1.5% of an essential oil, starch and tannin. No systematic pharmacological investigations have been carried out.

Actinodaphne hookeri Meissn (Vern.-Pichli, Bo -Pisa).

Grows in Sikkim and on the eastern and western Ghats. Infusion from leaves is used as a remedy for urinary disorders and oil from seeds is used as an external application in sprains of joints. Bark contains a crystalline alkaloid named as actinodaphnine. Leaves contain amorphous salts and seeds an essential oil (1.0%), fat and a fixed oil. No pharmacological or clinical works have been carried out.

Adhatoda vasica Nees (Malabar nut tree. Vern -Arusha, S.-Vasaka, H. & B.-Adulsa, B.-Bakas, M Adhatoda) B. P. C., I. P. C. & I. P. L.

Grows all over the plains of India and in the lower Himalayan ranges ascending to a height of 4,000 ft. Leaves, roots and flowers are used as a remedy against chronic bronchitis, asthma and phthisis. Juice of leaves is used in diarrhoea, dysentery and malarial fevers. It is also used locally as counter-irritant. It contains an alkaloid vasicine and an essential oil. The alkaloid vasicine has no very marked pharmacological action and produces only slight bronchodilation. The essential oil acts as an expectorant. Fluid extract from leaves relieves coughs. It is said to be effective in asthma and pulmonary tuberculosis. An antibiotic principle having a powerful action on tubercle bacillus has been discovered.

Aegle marmelos Correa (Bael fruit Vern -S.-Sripal, H, B. & Bo -Bael). B. P. C., I. P. C. & I. P. L.

Grows wild all over sub-Himalayan forests, in Bengal, South India and in Burma. Root, bark, leaves and fruit are used to cure gastrointestinal disorders. Roots, bark and leaves contain reducing sugars and tannin and fruit contains marmelosin, the most potent active principle. Marmelosin acts as laxative and diuretic in doses of 0.05 gm. Larger doses depress the heart. Extracts made from the fresh or dried unripe fruit are used effectively in treatment of chronic diarrhoeas and dysenteries.

Alangium salviifolium (Linn f) Wang Syn -*A. lamarckii* Thwaites (Vern -S -Ankota, H -Akola, B -Akar Kanta, Bo -Ankola, M.-Alangi).

Grows in forests throughout India and Burma. Root bark is used as an anthelmintic and purgative also as a remedy for leprosy and skin diseases. Bark contains an alkaloid alangin (0.82%) which produces a fall in blood pressure and depression of the heart. Respiration becomes irregular. Parasympathetic mechanism is stimulated. No clinical trials have been carried out.

Allium sativum Linn (Garlic Vern -S -Lasuna, H & Bo -Lasan, B -Rasun, M.-Vallarpundu). B. P. C. & I. P. C.

Is grown all over India and is used as a spice in cooking. It is considered to be hot and stimulant and useful in fevers, coughs and other debilitating conditions. Externally it acts as rubefacient in skin diseases. It contains a volatile oil (yield from 0.06 to 0.1%), allacin and a crystalline substance isolated from the alcohol-insoluble fraction of the residue. Garlic has an antiseptic action and slightly tonic effect on frogs' heart and a paralysing effect on isolated rabbit intestine. Garlic juice is used as an antiseptic for wounds. It is considered useful in digestive

disorders and respiratory diseases such as pneumonia, pulmonary tuberculosis and bronchitis. Contains a powerful antibiotic principle effective against tubercle bacillus.

Aloe vera ex Linn. (Vern.-Musabbar, S & B -Ghrita Kumari, H.-Ghi kanvar, M -Kattalai) Syn -*A. barbadensis* Mill B P , B P C., I P C & I P L

Native of North Africa but grows in East and West Indies, India and China. In India it is found in the dry valleys upto Cape Comorin. It is used as stomachic, purgative and emmenagogue, and in the treatment of piles, rectal fissures and as a poultice for application to inflamed parts. It contains a mixture of glucosides called aloin. It is valuable in the treatment of constipation.

Alpinia galanga Willd (The greater Galangal Vern -H -Bare kulinjan, B -Sugandha-vacha, M -Pera-rattai) I P C

Is a native of Sumatra and Java but commonly found in East Bengal and South India. Rhizome is used in bronchial catarrh and rheumatism, tubers and seeds are carminative and useful for impotence and nervous debility. Root contains compheride, galangin and alpinin. From the green rhizomes a pale yellow oil is obtained. The drug is depressant to cardio-vascular system and stimulates respiration in smaller doses but larger doses depress it. Bronchioles are dilated. It is used as an expectorant in respiratory troubles of children and in asthma, intestinal and biliary colics.

Alstonia scholaris R Br (Dita-bark Vern -S -Sapta parna, H -Chatium, B -Chhatim, M -Edakula-pala) I P C

Is cultivated throughout India and found in sub-Himalayan tract upto 3,000 ft. Also found in Bengal and in Southern India. Bark is used as tonic and alterative. It is considered useful in fevers and skin diseases. It contains a bitter principle ditain, alkaloids ditamine, echitamine, and acid and fatty resinous substances. The alkaloids are not protoplasmic poisons. They relax isolated strips of rat intestine. In the intact cat, the alkaloids produce a sharp fall of blood pressure. Larger doses produce irregularities of heart beat. Drug is used in the treatment of malarial fever, chronic diarrhoea and advanced stages of dysentery. Recently shown to have no specific curative action in malaria.

Amomum subulatum Roxb The greater Cardamom (Vern -S -Brihat-upa-kunchika, H & B -Bara-elachi, M -Periya-yelakkay) I P C

Grows in Nepal and Darjeeling Hills. Used for flavouring purposes. Seeds contain an oil rich in cineole.

Andrographis paniculata Nees (The Creat Vern -S -Bhunumba, H -Kiryat, B -Kalmegh, M -Nila vembu) B P C , I P C & I P L

Is found throughout the plains of India from Lucknow to Assam and is also cultivated in gardens in some parts of India. Leaves and juice are used to relieve griping, irregular stools and loss of appetite in children. Roots and leaves are used as febrifuge, tonic, alterative and anthelmintic. Leaves contain two bitter substances and traces of an essential oil. No systematic pharmacological investigations have been carried out. It has an intensely bitter taste.

Antiaris toxicaria Letsh. (The Upas tree Vern -Chanduka, Bo - Chandla, M -Nettavi, Sing -Riti, Burm -Hmyaseik)

Grows in Java, Malaya and Burma In India it is found in Concan, Canara, Travancore and Coorg Juice from leaves and bark is used as an arrow poison in Java, Malaya and Burma Bitter seeds are used as a remedy for dysentery and as a febrifuge in doses of half a seed three times a day. Milky juice of the plant contains antiarol, large amounts of potassium nitrate, antiaresin, a crystalline protein and three glucosides α antiarin, β -antiarin, γ -antiarin Preliminary toxicity tests on guinea pigs revealed that the drug is highly poisonous and the M L, D is 6 mgm per kilo body weight Death occurs due to heart failure It also produces marked tonic contraction of the isolated and intact intestines No therapeutic trials have been carried out

Arachis hypogaea Linn (Vern -S -Buchanaka, H -Mungphali, B -Chiner badam, Bo -Bhui-chane, M -Nila kadalai) Ground Nut B P., B P C, I P C & I P L

Originally a native of Brazil but now cultivated in all tropical or sub-tropical countries India is the largest producer The seeds are used as food, and oil as nutritive, laxative and emollient Seeds yield 40 to 50 per cent of a clear fixed oil which is used as a substitute for olive oil in India, as a basis for liniments and ointments Also used in the manufacture of soap and cosmetics, leather dressing and as substitute for tallow and diesel oil

Areca catechu Linn (The areca or betelnut palm) (Vern -S.- Gubak, H & B -Supari, Bo -Sopari, M -Kumugu) B P C, I P C & I P L

Grows in Sunda Islands but extensively cultivated in Southern India, Assam and Eastern Archipelago It is widely used in India and China as anthelmintic in man and animals, and as a masticatory to prevent the decay of teeth The seed contains alkaloids arecaine 0.1%, arecoline 0.07 to 0.1%, arecaidine, guvacoline, guvacine and choline, tannin 15% and fat 14% Arecoline is a highly toxic substance and its action resembles muscarine The seeds are used as anthelmintic in doses of 6 drams and for astringent action in relaxed condition of bowels.

Argemone mexicana Linn (Mexican Poppy) (Vern -Bharbhand, Kandiar, S -Srighala-kantaka, H & B -Siakanta, M -Birama-dandu)

Grows wild all over India Juice of plant is used to cure dropsy, jaundice and cutaneous affections The seeds and seed-oil is used as a remedy for dysentery and intestinal affections The plant contains alkaloids, berberine and protopine The seeds yield about 22% of argemone oil which is used as a purgative but has no special advantage over other purgatives It is used as an adulterant of mustard oil and is believed to produce epidemic dropsy.

Aristolochia indica Linn. (Vern.-S.-Rudrajata, H. & B.-Isharmul, M.-Ichchura-mula). I P.C. & I.P.L.

Grows all over tropical portions of India, Bengal, Konkan, Travancore and Coromandel Coast. Root is used in the treatment of bowel complaints in children and in fevers. It possesses emmenagogue and anti arthritic properties. Constituents of the root are an essential oil,

a bitter substance and an alkaloid aristolochin. The alkaloid produces cardiac and respiratory paralysis in frogs and mice. Skeletal muscles are stimulated in small doses. More detailed pharmacological investigations and clinical trials are indicated.

Artemisia Sp (Vern —Kirmala)

Grows abundantly in the Himalayas from Kumaon to Kashmir at altitudes of 4,000 to 12,000 ft, also in Baluchistan, Chitral and Afghanistan. Flowering tops are used as anthelmintic in 2 to 4 dram doses. The drug is also used as a remedy for dropsy and as cardiac and respiratory stimulant. It contains a volatile oil, santonin and an allied body—artemisin.

Artemisia absinthium Linn Wormwood (Vern -H. & Dec - Vilayatı afsantin).

Grows in Kashmir at altitudes of 5,000 to 7,000 ft. The oil has a tonic and stimulating effect on the digestive system. Contains 0.3 per cent. of an essential oil, a bitter glucoside absinthin and a crystalline compound.

Artemisia dracunculus Linn

Found in western Tibet at altitudes of 14,000 to 16,000 ft and in Lahul. Leaves are used for flavouring vinegar and as a spice. They contain about 0.3 per cent. of an essential oil.

Artemisia maritima Linn (Vern -S -Gadadhar, H -Kirmala, Bo. Karamani owa) I P C & I P L

Found in Western Himalayas from Kashmir to Kumaon at altitudes from 7,000 to 9,000 ft in Kashmir, Kurrum, Kagan, Bushher, Waziristan and Chamba.

Artemisia pallens Wall ex-DC (Vern -Davana)

Grows in certain parts of South India in Mysore State and is also cultivated near Poona. Leaves are used as floral decoratives and the oil from the herb is a popular article in perfumery.

Artemisia sacrorum Ledele (Vern P.-Tatwen)

Grows in Western Tibet, Kanawar and in Tibetan regions of Kumaon at altitudes of 10,000 to 12,000 ft. Contains 1 per cent. of an essential oil.

Artemisia vulgaris Linn (Vern -Tithwan, S -Nagadamani, H -Nagadouna, B -Nagdona).

Grows throughout the mountainous regions of India ascending to an altitude of 12,000 ft., in western Himalayas upto 5,000 to 8,000 ft, in Sikkim and Khasia Hills, also in Mount Abu and Western Ghats. Used as antiseptic expectorant and anthelmintic. Contains 0.2 per cent. of a volatile oil which is a good larvicide.

Asteracantha longifolia Nees Syn.-*Hygrophila spinosa* T Anders.
(Vern -S.-Kakilakshya, H -Talmakhana, B. Kuliakhara, Bo.-Talm khana,
M -Vallara1) I P C

Grows commonly in moist places throughout India from Himalayas to Ceylon. Roots are used as cooling, diuretic, stimulating and to remove dropsical fluids and kidney stones. They contain phytosterol, traces of an essential oil, a yellowish green wax, gum and maltose. Seeds contain an oil 23 per cent and an alkaloid. No pharmacological and clinical studies have been carried out.

Atropa acuminata Royle (Vern -Sag-angur) B P , B P C & I P C.

Grows in abundance in the Himalayan ranges extending from Simla to Kashmir, at altitudes of 6,000 to 12,000 ft. In Jammu & Kashmir State, it is found in the forest mountains round Kashmir valley. It is also found in Himachal Pradesh, Kulu, Parbatı valleys and Narkanda forests of Simla Hills. Belladonna and its alkaloid atropine are largely used as a sedative, antispasmodic and mydriatic in diseases of the eye, and as antidote in opium and muscarine poisoning. Alkaloids (as hyoscyamine) are present from 0.15 to 0.7 per cent in the leave and 0.3 to 0.66 per cent in the root. Traces of an inactive volatile alkaloid is also present. The plant is now cultivated and large amount of belladonna preparations in India come from this source.

Azadirachta indica A Juss (Vern -Nim) The Neem. I. P. C. & I P L

Grows throughout the greater part of India and Burma. Bark and leaves are used as a bitter tonic and astringent. It is considered useful in skin diseases. Fruit is used as purgative, emollient and for expelling intestinal worms. Considered beneficial in urinary diseases and in the treatment of piles. Bark contains an alkaloid margosine and seeds yield a fixed bitter oil. Margosates are toxic to protozoa and bacteria. Clinical trials revealed that the drug has a beneficial effect in skin diseases.

Bacopa monnieri (Linn) Pennell, Syn *Herpestis monniera* (Linn) H B & K (Vern -S -Brahmi, H -Braehmbhi, B -Brihmisak, M -Nirbrami) I P. C

Grows throughout India on the borders of streams and tanks. Roots, leaf and stalks are used as nerve tonic and as a remedy for insanity and epilepsy. It contains an alkaloid, herpestine, which is highly toxic. It produces a fall of blood pressure and stimulation of respiration. The smooth muscle of intestines and uterus is stimulated. In therapeutic doses it resembles strychnine. Powdered dried leaves gave satisfactory results in cases of asthenia and nervous break down. Further clinical trials are needed.

Bambusa bambos Druce, Syn.—*B. arundinacea* Willd. (Vern.-Kantabans, S.-Vansa, B. & H -Bans, Bo.-Mandgay).

Grows wild throughout the greater part of India particularly in the hilly forests of Western and Southern India upto an altitude of 3,000 ft. Leaves are considered useful in diseases of blood, leucoderma and

inflammatory conditions Tabashir found as concretions in it is used against blood diseases, tuberculosis, asthma and leprosy. Seeds contain water 11.0 percent, starch 73.7 percent, albuminoids 11.8 per cent, oil 0.6 per cent, fibre 1.7 per cent and ash 1.2 per cent. Young shoots contain a cyanogenetic glucoside. No systematic work has been done to determine its proper therapeutic efficacy.

Berberis aristata DC (The barberry) (Vern - Chitra, S - Daru haridra H - Dar-hald) I P L. & I P C

Grows in the temperate Himalayas at an altitude of 6,000 to 8,000 ft. A tincture made from root bark was used as a bitter tonic, stomachic, cholagogue, antiperiodic and alterative. Berberine is the chief alkaloid, also umbellatine is present. It is used as diaphoretic and antipyretic in malarial fever but has little specification.

Berberis asiatica Roxb. (Vern.-Sumlu, H - Kilmora, Nepal - Chitro). I P L.

Grows in dry valleys of the Himalayas at an altitude of 3,000 to 7,500 ft in Bhutan, Garhwal and Behar. Its uses are similar to those of *B. aristata* and its active principle is berberine.

Berberis coriacea—Brandis (Vern - Simla-Kashmal) I P L.

Grows in N W Himalayas at an altitude of 8,000 ft

Berberis floribunda Hort

Grows in Nepal. The roots contain eight alkaloids namely Oxycanthine, bebamine, berberine, epiberberine, palmatine, dihydrocarydeline, jatrorrhizine and columbamine. No detailed pharmacological studies have been carried out but the action of these alkaloids would appear to resemble berberine.

Berberis himalaica

Found in Bhutan. Bark contains berberine (0.03%), jatrorrhizine (0.01%) and an alkaloid limanthine. It has not been used in medicine although it belongs to berberis group of plants.

Berberis insignis Hook

Grows in humid forests of the Eastern Himalayas from Nepal and Sikkim to Bhutan at altitudes of 8,000-10,000 ft. Stem bark contains 1.52 per cent and root bark 2.5 per cent of total alkaloids consisting chiefly of umbellatine.

Berberis lycium Royle (Vern H - Kashmal, Bo - Darhald)

Grows in the Western Himalayas at altitudes of 3000-9000 ft from Garhwal to Hazara. It is used as febrifuge, carminative and aperient and is considered beneficial in piles, enlargement of liver and spleen and as external application to eyelids in conjunctivitis. It contains alkaloid umbellatine.

Berberis umbellata Wall.

Occurs on Himalayan ranges at altitudes from 9,000 to 12,000 ft. from Kashmir eastward to Bhutan. Bark contains Umbellatine which like berberine has a specific inhibitory action on *Leshmania tropica*. The cardiovascular system is depressed and para-sympathetic nervous system is stimulated. Respiration is depressed.

Berberis vulgaris Linn (The true barberry) Vern. Zırışk, P - Kashmal) I P L

Grows in the Himalayas from Nepal westwards at an altitude of over 8,000 ft. It is used as diuretic, astringent and refrigerant. In small doses it acts as a tonic and in large doses as a purgative. It contains berberine which has an inhibitory action on *leishmania*.

Blepharis edulis Pers (Vern —H -Utangan, Bo -Utangan).

Grows in the Punjab and Baluchistan. Leaves used as astringent in bowel complaints. It has aphrodisiac and alterative properties. It is useful in fevers, urinary discharges and leucoderma. Seeds contain a bitter glucoside belapharine (1.2 per cent) and dl-allantoin (2.1 per cent.) No detailed pharmacological studies have been carried out.

Boerhaavia repens Linn (Vern Thikri, S - Shothaghni, H -Sant, B -Punarnaba, Bo -Ghetuli, M -Mukuk-rattai) B P, I P C. & I P L.

Grows all over India. It is used as laxative and diaphoretic in oedema, anaemia, heart disease, cough and intestinal colic. It contains an alkaloid named punarnavine and an oily amorphous matter. Sulphates, chlorides, nitrates and chlorates occur in ash. The alkaloid is not toxic. It has a diuretic effect and raises blood pressure. The movements of intestines are depressed. The liquid extract in doses from one to four drams produces definite diuresis in patients with oedema and dropsy.

Bragantia wallichii R Br (Vern —Chakrani).

Grows in southern half of Bombay State near the Coast, Madras and Ceylon. Used to cure chronic sores and ulcers. Considered useful in cholera and diarrhoea. Roots contain a fatty oil and a yellow substance identical with isoaristolochic acid. No pharmacological studies or clinical trials have been carried out.

Butea monosperma Lam -Kutze (Vern. Palas) I P C

Grows throughout India and Burma extending in the N W. Himalayas as far as Jhelum. Gum (Kino) is used as astringent, diuretic and aphrodisiac. Seeds are used as an anthelmintic and contain 18 per cent. of a fixed oil, small quantities of resin and large amounts of water-soluble albuminoids. The oil does not possess any marked anthelmintic action. No systematic investigation has been carried out.

Butea superba Roxb. (Vern.-Palarvela, S Lata palas, H. & B.-Paas lata, M.-Kodi-murukkan).

Grows in the forests of Oudh and Bundhelkhand, Chotanagpur, Central and Southern India. It is used as a remedy against eruptions in children. Decoction of stem and leaves is emollient and is used as a

local application for piles. Root contains an esterogenic substance and a poisonous substance. Flowers contain butrin, butein and butin. An aqueous extract equivalent to 0.002 per cent of the dried powder causes estrus in spayed female mice. Further pharmacological studies are indicated.

Caesalpinia crista Linn Syn - *C. bonducella* Fleming. Fever nut (Vern -S -Kuberakshi, H -Kat-karanj, B -Nata-karanja, Bo -Sagara-goth, M -Gajega) I P.C.

Grows near the sea-coast, all over Bengal, Bombay and S India. Seeds, leaves and roots are used as anthelmintic, emmenagogue and febrifuge. Seeds contain starchy matter, a bitter principle called bonducin and an oil. The plant is not of much therapeutic value.

Calotropis gigantea Linn.- (Gigantic swallow root) (Vern Ak-Madar, S -Arka, H -Ak, B & Bo -Akanda, M -Erukku) I P C.

Grows in waste lands all over India upto altitudes of 3,000 ft. Milky juice is used for tanning and dyeing purposes. An intoxicating liquor is prepared from the juice. Latex contains water soluble matter 86 to 95.5 per cent. Stem bark contains calotropeolis, B-amyrin and giganteol. No marked therapeutic value.

Calycopteris floribunda Lam (Vern H -Kokoranj, Bo.-Ukshi, M -Marsada boli)

Grows in Central and Southern India in plains and upto an altitude of 2,500 ft. Leaves used as laxative, anthelmintic and the juice given in puerperal fevers. Also used in the treatment of dysentery and malarial fevers. The juice contains, chlorides, sulphates, nitrates, lime, ferric oxide, tannin, albuminoids and gummy matter. Leaves contain a yellow crystalline flavone, calycopterin. The drug is toxic to fish and earthworms, more toxic to earthworms than santonine or chenopodium oil. No detailed pharmacological and clinical investigations have been carried out.

Camellia sinensis Linn (O Kunfize) Vern -Cha Tea plant B P & B.P.C.

In India, tea and coffee grow luxuriantly. Coffee is grown principally in Madras, Coorg, Mysore, Travancore and Cochin. Tea is also found in Ceylon and is grown in Assam, Bengal, Madras and Travancore. Tea is used as a beverage in India and other parts of the world. Tea leaves contain from 2.5 to 3% of Caffeine. Coffee contains 15 per cent of caffeine. Caffeine is used in medicine for stimulation of central nervous system and circulation and as a diuretic.

Cannabis sativa Linn. (*Cannabis indica*) (Vern S.-Ganjika, H & Bo -Ganja, M -Bhang, Bhang) B P C, I.P C. & I.P L

Grows wild throughout the Himalayas from Kashmir to East of Assam and also in Southern India. Plant and its products are used for narcotic purposes, by smoking or taking internally. It contains a terpene 15 per cent, a sesquiterpene 17.5 per cent., small amount of paraffin carbon and a resin named as Cannabinol (33 per cent.). Hemp drugs are consumed all over the world for euphoric purposes. Their use alleviates the feeling of fatigue, encourages sleep and soothes restlessness.

Sensation of time and space is lost. Larger doses produce mental derangement and even insanity.

Carica papaya Linn. (The Papaya) (Vern Papita, H.-Papaya, B.-Papey, Bo -Papai, M -Pappayi). B.P , B.P.C , I P.C & I.P L.

Grows all over India and is used as anthelmintic, emmenagogue and digestive. The fruit pulp contains sucrose, invert sugar, resinous substance, papain, malic acid, salts of tartaric and citric acids and vitamins. Seeds contain, proteins, carbohydrates, fatty oil, volatile oil, an alkaloid carpaine, a glycoside carposide and an enzyme myrosin. Carpaine is a heart poison, it lowers blood pressure and depresses heart. No proper clinical trials have been carried out.

Carum carvi Linn The Caraway seed (Vern Zira, H.-Shiajira, B -Jira, M Shimai-shombu) B.P., B P C & I P C

Widely distributed in the temperate regions of both the hemispheres. Grows in North and Central Europe, extending to the Caucasus, Persia, Tibet and Siberia. In India it is cultivated on the hills of Baluchistan, Kashmir, Kumaon, Gharwal, Chamba at altitudes of 9,000 to 12,000 ft. Used as a cookery condiment. Oil is used for flavouring purposes and as a carminative. Seeds contain an essential oil with a marked carminative action.

Cassia absus Linn (Vern H —Chaksu, N —Karun Kanam).

Grows in lower parts of the Western Himalayas and Ceylon. Leaves are applied locally to tumours and internally in cough, asthma and bronchitis. Seeds are used as bitter, astringent and diuretic and contain two water soluble bases, chaksine and Isochaksine, as carbonates (15 per cent). Preliminary studies show, that chaksine sulphate is depressant to heart, respiration and nerve centres. More detailed studies are indicated. Considered useful in eye diseases. No clinical trials have been carried out.

Cassia angustifolia Vahl (Vern H -Hindisana, B.-Sonamukhi, M -Nilavakai) B.P , B P C & I P.C.

Cultivated in S India in Madras State (Madura). Leaves and pods are used as purgative. Leaves and pods contain cathartic acid, emodin and chrysophanic acid. Senna is useful in habitual constipation. It increases the peristaltic movements of the colon. The tendency to gripe is overcome by mixing it with aromatics.

Cedrela toona Roxb (Vern Mahalimbu, S. & B.-Nandibriksha, H -Tun, Bo.-Tuni, M -Tunumaram, P -Khushing).

Grows in Sub-Himalayan tracts from river Indus eastwards, Chittagong, Assam, Burma, Chotanagpur, Western Ghats of Bombay to the Nilgiris and other hills of the Deccan Peninsula. The plant is used as astringent in the treatment of ulcers, gleet and scabies. Flowers are used in menstrual disorders and contain a red colouring matter, nyctanthin, flavone and a glucoside. Bark contains, tannic acid, a bitter resin, citric acid and starch. No pharmacological or clinical investigations have been carried out.

Celastrus paniculatus Willd (Vern. S -Kanguni, H -Malkanguni, M -Valuluwai)

Grows in Sub-Himalayan tracts extending from Jhelum to Assam ascending to an altitude of 6,000 ft and in hilly parts of Bombay State, South Gujrat, Central India and Madras. Seeds and oil are used in the treatment of rheumatism, gout, paralysis and epilepsy. Oil is considered as nerve stimulant and brain tonic. Leaves contain an alkaloid, a glucoside and colouring matter. Seeds contain 52.2 per cent of a thick brownish oil. No pharmacological studies have been carried out.

Centella asiatica (Linn) Urban, Syn -*Hydrocotyle asiatica* Linn (Vern -Brahmi, S -Vanduka paini, H -Brahmo manduki, B -Tholkuri Bo, -Karivana, M -Vallara) I P C

Grows throughout India from Himalayas to Ceylon at an altitude of 2,000 ft. Plant is used as tonic and is considered beneficial in diseases of the skin, nervous system and blood. Also used in the treatment of leprosy and syphilitic affections. It contains an alkaloid, hydrocotyline and fresh leaves contain a glucoside asiaticoside. Hydrocotyle is shown to be a circulatory stimulant. Further work is indicated.

Centrathrum anthelminticum Kuntze, Syn -*Vernonia anthelmintica* Willd (Vern -S, H & B -Sōmaraj, Bo -Kaliyiri, M -Kattu-Shiragam). I P C & I. P. L.

Is found in waste lands near villages throughout India. The seeds are used as a remedy for leucoderma and other skin diseases, and as anthelmintic. They contain a bitter principle 1 per cent, a fixed oil 18 per cent and a small amount of an essential oil. No systematic pharmacological investigations have been carried out. Clinical trials revealed that it is effective in thread-worm infections. Further work is needed.

Cephaelis ipecacuanha A Rich Syn -*Psychotria ipecacuanha* Stokes B P, B P C & I P C.

Grows wild in Brazil. In India it has been cultivated in South India and Bengal—in Darjeeling hills. Contains emetine and other alkaloids used in the treatment of amoebic dysentery.

Cerbera odollam Gaertn Syn -*C. manghas* Linn. (Vern B -Dhakur, M -Katarali).

Grows all over India particularly in salt swamps in the Malabar Coast, Ceylon and Burma. Plant is intensely poisonous. The leaves and sap have emetic and purgative properties. Seeds contain three glycosides named as cerberin, odollin and cerbeside. Cerberin is non-irritant to the skin and mucous membranes and produces vomiting, diarrhoea and even syncope when given subcutaneously. Stimulates cardiovascular system and smooth muscles. Further clinical trials and experimental work is indicated.

Chenopodium ambrosioides Linn B P, B.P C, I.P.C. & I.P.L.

Grows in Central America, East Indies. In India found in Bengal, Sylhet, the Deccan (Coimbatore). Infusions from leaves and seeds used as a remedy against intestinal worms. Seeds contain a volatile oil which

contains ascaridole from 45 to 70%. Small portions of glycol-anhydride, mixture of hydrocarbons and traces of fatty acids also occur. Clinical trials have shown its efficacy against hookworms and round worms.

Chenopodium botrys Linn

Grows in the Himalayas from Kashmir to Sikkim. Used as a substitute for *C. ambrosioides*. Contains a volatile oil 0.03 to 0.04 per cent.

Cichorium intybus Linn (Vern - Kasni)

Grows in North Western India and is cultivated in Nadiad, Broach and Amalsad in Bombay State. Used as fodder. Cultivated sweet variety is considered beneficial in acne, ophthalmia and throat inflammations. Roots used as diuretic and to enrich and purify blood. Wild variety used as emmenagogue and to cure asthma. Plant contains a glucoside cichorin. Seeds contain, a semi-drying oil, water, gum, glucose, bitter substances, inulin and fibre. No systematic studies of its therapeutic action carried out.

Cinchona Cinchona bark B.P., B.P.C., I.P.C. & I.P.L.

Cinchona grows on the Eastern slopes of Central Western chain of the Andes mountains in South America, at altitudes of 2,500 ft to 9,000 ft. It is cultivated in India in the Nilgiris Hills, Mungpoo in Ranchi Valley and the Karen Hills in Burma. The bark contains quinine, cinchonidine, quinidine and cinchonine. It is used in the treatment of malaria.

Cinnamomum Blume.

Grows in Asia and Australia. About 20 species occur in India. Used as an aromatic and spice.

Cinnamomum camphora Nees B.P. & B.P.C.

Cultivated in India at Dehradun, Saharanpur, Calcutta, Nilgiris and Mysore.

Cinnamomum zeylanicum Breyn (Vern — Dalchini, S -Gudatreaka) B.P., B.P.C. & I.P.C.

Grows wild on Western Ghats from Konkan southwards and in forests of Tenasserim in Burma. Found abundantly in Ceylon. Used as carminative, astringent, stomachic, flavouring agent and as a spice. Bark contains an essential oil 0.5 to 1%. Leaves yield a dark coloured oil 1%.

Cissampelos pareira Linn (Vern — Akanadi, S -Patha, H & B -Nirbisi) I.P.C.

Grows throughout tropical and sub-tropical parts of Asia, East Africa and America. Root is used to relieve pain and is considered useful in febrile conditions, dysentery, heart troubles and urinary disorders. Contains an amorphous alkaloid named pelosine, saponin and ammonium salts, also hayatin and hayatinin. Hayatin methiodide, methochloride have curariform activity equal to tubocurarine chloride. They produce fall of blood pressure which is prevented by antihistaminic drugs. Further studies are in progress. A plant likely to be of great therapeutic value.

Citrullus colocynthis Schrad. (Vern.—Indrayan, S.-Indra-varuni, H. & Bo-Indrayan, B.-Makhal, M.-Peyt-tumatti). B.P., B.P.C. & I.P.C

Grows in arid tracts of N West, Central and South India and is met with in the Punjab, Sind and on the Coromandel coast. Fruit is used as cathartic to relieve biliousness and constipation, used in treatment of fever and intestinal parasites. Plant contains a bitter principle colocynthin and traces of an alkaloid. Used in medicine as a drastic purgative.

Citrus aurantii-folia (Christm) Swingle Syn - *C medica*, var *acida*.
The Lime Tree.

Citrus limon (Linn) Burm f Syn, - *Citrus medica*, var. *limonis*.
The Lemon Tree B.P., B.P.C., I.P.C & I.P.L

Lime and lemon are found growing wild in the warm valleys of the North West Himalayas. Also largely cultivated in the plains and up to altitudes of 4,000 ft. Used as flavouring agent in medicine and also as a carminative and stomachic. Peel contains an essential oil and juice citric acid upto 5.9 per cent.

Cleome icosandra Linn. (Vern — Aurhur).

Grows throughout India but commonly found in Bengal and S. India. Seeds are used as anthelmintic, rubeficient and vesicant. They contain a fixed oil (36.59 per cent) and a new flavone called 'Viscosin'. No work on its pharmacological action and therapeutic properties has been carried out.

Clerodendron infortunatum Linn (Vern.—H & B.-Bhant, Bhat)

Grows in waste lands throughout India and in damp forests of Ceylon, upto an altitude of 5,000 ft. Used as a vermifuge, anthelmintic and a substitute for Chireta. Leaves contain ash 8.04 per cent, protein 21.1 per cent., crude fibre 18.84 per cent, total sugars 17.05 per cent and a bitter substance clerodin. The last named has no haemolytic and bactericidal action. It is toxic to earthworms, small fish and mosquito larvae. No clinical trials carried out.

Coccinia cordifolia Cogn Syn - *Cephalandra indica* Naud, (Vern.-S - Bimba, H - Kanduri-ki-Bel, B - Telakucha Bo - Bhimb)

Grows wild in Bengal and other parts of India. It is said to have antidiabetic properties. It contains an enzyme, a hormone and traces of an alkaloid. Did not show any action on blood sugar in animals nor by clinical trials in diabetic patients.

Colchicum luteum Baker (Vern — Surinjan) I.P.C. & I.P.L.

Grows extensively in the Western temperate Himalayas, extending from Murree hills to Kashmir and Chamba. Used as an alterative and aperient in gout, rheumatism and diseases of liver and spleen. Bitter variety contains alkaloid colchicine, starch and oily resinous matter. Galenical preparations of the crude drug and the alkaloid colchicine are used in the treatment of gout. Recently colchicine has been widely used in plant breeding to induce polyploidy.

Commiphora mukul Engl. (Gum'gugul) (Vern -S'-Guggulu, H ,B. & Bo.-Guggul, M.-Gukhulu).

Grows in Rajputana, Khandesh, Sind, East Bengal and Assam. The gum resin from bark is used as demulcent, aperient, carminative and alterative and is useful in leprosy, rheumatism, syphilitic disorders, nervous and skin diseases and urinary disorders. Gum resin contains 32 per cent. gum, 19.5 per cent. mineral matter and 1.45 per cent of an essential oil. The pharmacological action of this oleoresin resembles the action of copaiba and cubebs. Taken internally it acts as bitter, stomachic and carminative. It also acts as diaphoretic, expectorant and diuretic. It is said to be uterine tonic and regulates menstrual functions. No proper chemical trials have been carried out.

Coptis teeta Wall. (Vern H -Mamira, B -Tita, Bo-Mahmira) I P.L.

Grows in upper Assam. It is used as an eye salve in indigenous medicine. Active principles berberine 85 per cent, coptine (0.08 per cent), palmatine (traces), coptisine 0.02 per cent and jatrorrhine (0.01 per cent). It is used as a bitter tonic in the same way as Calumba.

Coscinium fenestratum Colebr. (Vern —Halgach, S.-Daru hari-draka, B -Haldi-gach, M -Mara-manjal) I.P C.

Grows in the forests of Western India and is used as a bitter tonic. It contains berberine.

Crocus sativus Linn (Vern.—Kesar, Safran, S -Kumkuma, H & Bo.-Jafran, M -Kungumapu) B P C , I P C & I P L

Is commonly found in Kashmir and Quetta and also cultivated in Spain. It is considered to be stomachic, stimulant, antispasmodic and aphrodisiac. It is used as flavouring and colouring agent in cookery. It contains three crystalline colouring matters, α -crocetin, β -crocetin, γ -crocetin, a fatty oil, a volatile oil and a bitter substance. The essential oil has the characteristic features of other essential oils. No proper clinical trials have been carried out.

Curcuma longa Linn Syn -*C domestica* Valetton (Vern — S -Haridra, H.-Haldi, B -Halood, Bo -Halada, M -Munjal). B. P C. & I P C.

Grows throughout Southern parts of Asia, in India, China and East Indies. Also cultivated in almost all the States of India, particularly in Madras, Bengal and Bombay. Used as a spice. Hakims use it as a stomachic, tonic and blood purifier. The juice of rhizome has anti-parasitic action in affections of skin. Rhizome contains an essential oil 5.8 per cent., protein, mineral matters, carbohydrates, and vitamin A. The oil has slight anti-bacterial properties. Taken internally it has antacid action and relaxes the intestines. Used as a stomachic and carminative.

Curcuma zedoaria Rosc. (Vern.—S.-Sati, H, & Bo.-Kachura, B -Shori, M.-Pulan-ki-zhanga).

Grows in Eastern Himalayas and Kanara, Ceylon and China. It is also cultivated in many Indian States. Rhizomes are used as aromatic, stomachic, stimulant and carminative. Externally applied on bruises

and sprains. Contains an essential oil 1.5 per cent., starch 82.6 percent., ash 1.01 per cent and moisture 13.1 per cent. The essential oil is carminative and produces relaxation of the intestines

Cuscuta reflexa Roxb. (Vern —S -Amaravela, H -Akasbel, B.-Algusi, Bo -Nirmuli, M.-Sitama-purgonalu).

Grows throughout the plains of India upto an altitude of 5,000 ft. Used as carminative, purgative, diuretic and purifier of blood. Seeds contain a bitter substance, a glucosidal resin and quercetin. Plant also contains, cuscutin, cuscotalin, brown wax and reducing sugars. No pharmacological studies have been carried out.

Daemia extensa R Br (Vern —S.-Phala kantak, H.-Utran, B.-Chagulbanti, Bo & M -Utarni) I.P C

Grows throughout the hotter parts of India upto an altitude of 3,000 ft. Used as cooling, anthelmintic, laxative and antiperiodic. Considered useful in eye troubles, uterine disorders and urinary troubles. The plant contains 2.4 per cent of inorganic salts, a bitter resin, and three bitter substances. Bitter principle C is the most active pharmacologically. Has a stimulant action on uterus, intestines, due to direct action on the involuntary muscle. Further pharmacological studies are in progress.

Datura stramonium Linn (Vern —B -Sada dhutura, P -Tattu dattura, M -Umatai) B P, B P C & I P C

Grows throughout the temperate Himalayas from Kashmir to Sikkim. Used as intoxicant, emetic, digestive and as antispasmodic in asthma and whooping cough. Seeds and leaves contain hyoscyamine, atropine and hyoscine. Datura possess properties similar to belladonna.

Dichora febrifuga Lour. (Vern.—H -Basak, Nep -Aseru).

Grows in the temperate Himalayas from Bhutan to Khasia hills, at altitude of 4,000 to 8,000 ft, also in Upper Burma, Malayan Peninsula and China. Roots are used to cure malaria in China and as a febrifuge in India. Roots and leaves contain two alkaloids, febrifugine and Isofebrifugine. Clinical trials have revealed that the drug has antipyretic action like quinine (1/5 times) but no anti-parasitic action. Further work is indicated.

Didymocarpus pedicellata R Br

Grows in subtropical regions from Chamba to Kumaon at altitude between 2,500 and 5,000 ft. Leaves are used as a remedy for stones in kidney and bladder. The plant contains pedicin 1 per cent., iso-pedicin 0.4 per cent, pedicinin 0.3 per cent. and pedicellin 1 per cent which are crystalline. No detailed pharmacological and clinical trials have been carried out.

Digitalis lanata Ehrh. Grecian Foxglove B P, B.P.C. & I.P.C

Grows in Kashmir at altitudes about 7,000 ft. Fresh leaves contain 3 natural glucosides lanatosides A, B & C which on hydrolysis give digitoxin, gitoxin and digoxin. Digoxin produces the same cardiac effects as digitalis. It is 300 times more potent than prepared digitalis.

and is of particular value for rapid digitalization. Digitalis increases the force of systolic contractions and efficiency of the decompensated heart. Slows the heart rate and reduces cardiac oedema with diuresis.

Digitalis purpurea Linn Foxglove. B P., B P.C. & I P. C.

Cultivated or found as an escape in hilly regions at altitudes of 5,000–6,000 ft. Originally a native of Western Europe but now extensively grown in many parts of the world. In India, it is found in Kashmir, Himachal Pradesh, Darjeeling area and other parts. It is used as a myocardial stimulant in congestive heart failure. Used as a heart tonic and diuretic. Contains several glucosides but the leaves contain digitoxin, gitoxin and gitalin.

Dregea volubilis Benth. ex Hook (Vern—S—Madhu malati, H.—Nakchikni, B—Titakunga, Bo—Dodhi, M—Kodicpalay)

Grows wild in Assam, Bengal, Deccan, Peninsula from Konkan southward to Ceylon. The plant is used to cure piles, leucoderma, asthma and urinary discharges. It contains a substance of glucosidic nature with low toxicity and traces of an alkaloid. It stimulates all organs having cholinergic nerve supply and causes a prolonged fall of blood pressure. Further work is indicated.

Elettaria cardamomum Maton (Vern—S—Ela, H. & B—Chotrelachi, Bo—Elachi, M—Ellakai) B P., B P C., I P C & I P L.

Grows in Western and Southern India in the forests of Kanara, Mysore, Coorg, Wynad, Travancore, Cochin. Also found wild in Burma, Ceylon, China and Malaya Archipelago. Used as a spice and masticatory and in medicine as carminative and flavouring agent. Seeds contain an oil to the extent of 2 to 8 per cent.

Entada phaseoloides (Linn) Merr Syn.—*E. pursaetha* DC., *E. scandens* Benth

Grows in Central and Eastern Himalayas at altitudes of 4,000 ft and is also found in damp forests of Eastern Bengal, Bihar, Orissa, Eastern and Western Ghats. Seeds are used as soap to wash hair. Locally it is applied to swellings, it is a fish poison. Seeds contain two saponins, entada-saponin A & B and 8 per cent of a fixed oil. Both saponins are toxic to rabbits and guinea pigs. They produce hæmolysis of red blood cells and a depressant action on circulatory and respiratory systems. Movements of smooth muscles are inhibited. No clinical trials have been carried out.

Ephedra gerardiana Wall Syn.—*E. vulgaris*. and allied sp. (Vern Amsania) B P C., I P C & I P L.

All species grow in Northern India, Bashahr Division, Chakrata, Kangra, Kulu, Baluchistan, Kashmir, Hazara, Kagan and Waziristan. Contains ephedrine and pseudo-ephedrine. Ephedrine is a useful remedy for asthma. Tincture of Ephedra is an excellent cardiac stimulant in toxic conditions.

Erythroxylum coca Lam. B P. & B P.C.

Originally a native of South America but is grown in the West Indies, India, Ceylon and Java. Leaves used for euphoric purposes and

contain cocaine to the extent of 0.15 to 0.8 per cent. and other alkaloidal substances. Cocaine is used for local anaesthetic purposes and also as a euphoric

Eucalyptus L. W. Erit (Vern.—Karpura Maram) Blue gum tree, B.P., B.P.C., I.P.C. & I.P.L.

Grows naturally in Australia but cultivated in other parts of the world. The oil from leaves is used in soap industry and as an antiseptic and disinfectant. Leaves and terminal branches yield an essential oil

Eugenia aromatica Kuntze Syn -*Caryophyllus aromaticus* (Vern. Long) B.P. & I.P.C.

Native of Malacca Islands and is largely cultivated in Zanzibar, Pemba, the Amboyna Islands, Penang, Madagascar, Seychelles, Reunion, Mauritius and Ceylon and Southern India. Dried buds contain an essential oil and are used as aromatic stimulant and carminative

Euonymus tingens Wall (Vern —Barphali, H -Kungku) (Dogwood) I.P.C.

Grows in tropical regions of Asia, Malaya, Europe and America. Used to stimulate torpid liver producing flatulence and indigestion.

Euphorbia hirta Linn Syn -*E pilulifera* Linn (Vern —H -Dudhi, B -Bara keru, Bo -Nayeti, M.-Amumpatchaiarissi). B.P.C. & I.P.C.

Grows all over the hotter parts of India. It is used as a remedy for respiratory troubles, dysentery, colic and worms in children. It contains gallic acid, quercetin, traces of an alkaloid and an essential oil. The extract possesses depressant action on heart and respiration and produces relaxation of bronchioles. It is used to relieve asthma and other spasmodic conditions of the respiratory tract

Ferula foetida Regel. (Vern —S -Hingu, H & B.-Hing, Bo -Kayam) Asafoetida B.P.C., I.P.C. & I.P.L.

Found in Turkestan. Gum resin used as a carminative, antispasmodic and as flavouring agent in spices. Contains a volatile oil, resin, gum and impurities. Used in medicine as carminative and antispasmodic in nervous disorders of children and women

Foeniculum vulgare Mill (Vern —S -Madhurika, H —Bari-saunf, B -Pan-mauri, Bo -Bari sophia, M -Shombu). B.P., B.P.C., I.P.C. & I.P.L.

Cultivated throughout India and also in cold places upto an altitude of 6,000 ft. Used as aromatic, stimulant and carminative. Fruit contains a volatile oil with pleasant aromatic odour, chief constituent of which is anethol but small quantities of substances like fenchone are also present.

Gaultheria fragrantissima Wall. (Vern.—Jar Gandapuro). Indian winter green. I.P.C. & I.P.L.

Grows freely in the Nilgiris, Travancore and Assam. Oil of winter green used as an external application for rheumatic affections, sciatica,

neuralgia and as a flavouring agent in tooth pastes, etc. Leaves contain an oil which mainly consists of methyl-salicylate.

Glycyrrhiza glabra Linn. (Vern —S -Yashti-madhu, H -Jethimadh, B. & Bo -Jashti madhu, M -Atimaduram) B P., B P.C. & I.P C.

Found in sub-Himalayan tracts from the Chenab westwards, Burma and the Andaman islands Used as a tonic, demulcent in catarrh of respiratory tract and genito-urinary passages and as laxative Roots contain a tribasic acid glycyrrhizic acid and potassium and calcium salts, glucose, sucrose and fat 0.8 per cent. Used in medicine to cover the acrid taste of many nauseous drugs.

Gymnema sylvestre R Br (Vern —Gur mar, S -Meshasringi, H & B —Merasingi, Bo -Kavali, M -Shiru kurunja)

Grows in Central and Southern India and in Tropical Africa It is used as stomachic, diuretic, antiperiodic, antidiabetic and in urinary disorders It contains resins, a glycoside named as gymnemic acid, a bitter principle and some tartaric acid and calcium oxalate Experiments on animals showed no reduction of blood sugar The leaves stimulate heart and circulatory system and increase urine secretion Clinical trials showed no effect on blood or urine sugar of diabetic patients

Helicteres isora Linn (Vern —Kapasī) The East Indian screw tree

Is found throughout Central and Western India It is chiefly used in intestinal disorders such as colic, flatulence and diarrhoea It contains a demulcent substance and tannins The pods are used even today in some parts of India in the treatment of chronic diarrhoeas but clinical trials showed no marked action

Hemidesmus indicus R Br (Vern —S -Ananta, H -Magrabu, B. Anantamul, M.-Nannari) Indian Sarsaparilla I.P C.

Grows in Northern India, common in Bengal and in the Deccan, extending to Travancore and Ceylon and Bombay State. Used in the treatment of nutritional disorders, syphilis, chronic rheumatism skin affections and as a blood purifier Roots contain a steroptene, an essential oil, sterols, resin, tannin and small amount of glycosides Clinical trials show that its medicinal value is in no way inferior to imported Sarsaparilla.

Holarrhena antidysenterica Wall (Vern —Dhudi, S -Kutaja, H.-Karchi, B -Kurchi, Bo.-Pandhrakura, M.-Kashappuvetpalarishi) B P., B.P C, I P.C. & I P L

Grows throughout India upto an altitude of 3,500 ft and even as far south as Travancore Bark and seeds are used as a powerful antidysenteric, astringent, febrifuge, anthelmintic, carminative and aphrodisiac Bark and seeds contain kurchine, conessine, kurchicine and holarrhine Conessine is toxic to protozoa and is locally irritant The circulatory system is depressed and intestinal movements are stimulated, it has narcotic action on frogs The total alkaloids have similar action The extract from bark (containing one grain of total alkaloids in 4 c.c) given in doses of 2 drams 3 times a day has beneficial effect in amoebic dysentery especially the chronic form. It has proved effective against trichomona infestations.

Holarrhena febrifuga Klotizseh

Grows in Kenya, Tanganyika, Northern Rhodesia and Nyasaland. Used in the treatment of influenza, bilharzia, syphilis and also as substitute for quinine. Stem bark and leaves contain conessine 6.0 per cent and a subsidiary norbase isoconessiomine. Further studies indicated.

Hydnocarpus anthelmintica Pierre.

It grows in Siam, Northern Cochin and Gamboja

Hydnocarpus wightiana Blume (Vern.—Janglibadam, Kowti, Bo-Kava, M.-Yetti). B P., I P.C & I P L

Grows in western parts of the Indian Peninsula from South Konkan along the coastal range. The oil has been used in the treatment of leprosy for centuries. Seeds contain Chaulmoogra oil and Palmitic acid. Chaulmoogra oil has a bacteriostatic action on acid fast bacilli. It is extremely irritant to the tissues and for this reason ethyl esters such as moogrol (British) and Antileprol (German) are used by injection in the treatment of leprosy

Hyoscyamus niger Linn. (Vern —S -Parasikaya, H -Khurasani-ajvayan), B -Khorasani ajowan, Bo -Khorasani-owa, M -Khorasani-yomam) B P , B P C & I P C

Grows wild in temperate Himalayas at altitudes of 6,000 to 12,000 ft and also cultivated. Leaves contain alkaloids hyoscine and hyoscyamine

Ipomoea hederaceae Jacq & **I turpethum** Br (Vern —H , B & Bo -Kaladana, M -Jirkivirai) B.P C , I P C & I.P.L

Found throughout India ascending to altitudes of 3,000 ft. The resinous substance from the root bark is an excellent substitute for Jalap which is a drastic purgative. It is official in the I P L

Juniperus communis Linn. (Vern. H -Aaraar, P -Petthri) I.P C

Found in the Western Himalayas, Kumaon and Kurram valleys at an altitude of 11,000 ft above the sea level. It is used as digestive and diuretic. Berries contain a volatile oil 0.2 per cent, invert sugar about 33 per cent, resin 10 per cent, a bitter principle, organic acids and wax. Beverage gin is alcohol flavoured with juniper berries or the essential oil

Luffa acutangula Roxb. (Vern. S -Koshatak, H & Bo -Tora, B.-Jhinge, M -Pikumka)

Grows in N W India, Sikkim, Assam and East Bengal and also cultivated. Leaves used locally in splenitis, haemorrhoids, leprosy and granular conjunctivitis. Fruit contains a bitter substance luffein and seeds contain a fatty oil and a saponin. The latter is toxic to frogs and has a haemolytic action. It shows digitalis-like action and produces vomiting and diarrhoea. Further work is indicated

Luvunga scandens Ham. (Vern.—Jivanti, S -Lavangalata, B -Labangaphal).

Grows in Eastern Bengal, Assam, Khasia hills, Chittagong & Burma. Roots and berries are used to cure tuberculosis of lungs, biliousness

and fevers. Berries contain four crystalline neutral substances. Further chemical and pharmacological investigations are indicated

Madhuca indica J. F. Gmel. Syn - *M. latifolia* Mcbrd. The Mahu tree. (Vern -Mohua). I.P.C.

Grows in forests of Madhya Pradesh and is cultivated all over India. Also found in Uttar Pradesh, Bihar, Gujrat and Bombay. Its flowers, seeds and timber are of economic value. The seed-oil is edible and is also used for soap manufacture. The seeds contain 50 to 55% of a fatty oil and a saponin. Leaves contain a glucosidic saponin and traces of an alkaloid. Flowers contain sugar and enzymes. The plant is used as an astringent. It is largely employed as a lotion for chronic ulcers and bleeding spongy gums. Also considered beneficial in chronic bronchitis and chronic rheumatism. No proper clinical trials have been carried out.

Madhuca longifolia Linn. Macbd (Vern.—Mohua) Syn -*Bassia longifolia*. I.P.C

Is found in forests of Western India, Kanara, Malabar and Mysore in the same way as of *Madhuca latifolia*. Seeds contain 40 per cent. of fatty oil and sapo-glucoside called 'Mowrin'. Fruit contains saccharose 4.6 to 16.2 per cent. and maltose 2.39 per cent, tannins and enzymes. Acts in the same way as *M. latifolia*. The sapo-glucoside acts on the heart in the same way as digitalis. Requires further investigation

Mahonia nepalensis DC. Syn.-*Berberis nepalensis* Spreng (Vern.—Chiror Anrudana, P.-Amudanda, Nep.-Chatra)

Grows in the outer Himalayas from Ravi eastward to Khasia and Naga Hills also in the Nilgiris at an altitude of 5,000 ft. It contains umbellatine

Mallotus philippinensis Muell -Arg (Vern —Kambila, S -Rechanaka, H. & B.-Kamala, Bo -Shendri, M -Kapila) I.P.C

Grows throughout the plains of India and Ceylon, tropical parts of Asia and Australia, Bombay and Bengal. It is used as anthelmintic especially against tape worms. Fruit contains a brownish-red resin called rottlerin, an impure rottlerin, traces of a volatile oil, starch, sugar, tannins and oxalic acid. It has toxic action on frogs, tadpoles and worms. The drug has a marked cathartic and anthelmintic action in doses of 2 to 3 drams.

Melia azedarach Linn. (Vern.—Drek)

Grows wild in Sub-Himalayan tract at altitudes of 2,000-3,000 ft and is also cultivated. Root bark, fruit, flowers and leaves are considered hot, dry and to have deobstruent and resolvent properties.⁶ Juice of leaves used as anthelmintic, diuretic and emmenagogue. It contains a resin and the fruit a bitter substance. No systematic pharmacological studies or clinical trials have been carried out

Mentha arvensis Linn (Vern.—H.-Podina, B. & Bo.-Pudinah. M.-Pudina) The marsh mint B.P., B P.C, I.P.C. & I.P.L

Grows abundantly in northern and western Himalayas and in Kashmir at altitudes of 5,000 to 10,000 ft It is used as a flavouring agent and as carminative and stimulant Yields an oil similar to peppermint oil, to the extent of 0.18 to 0.2 per cent The oil contains 4 to 14 per cent. esters as menthyl acetate and 46% of free menthol Official in I.P.L

Moringa oleifera Lam Syn -*M pterygosperma* Gaertn. (Vern — Sanjna, Saonjna). I P C.

The plant is found wild in the Sub-Himalayan tract from the Chenab to Oudh and is also cultivated throughout India The root of this tree is extensively employed in intermittent fevers, epilepsy, hysteria, palsy, chronic rheumatism, dropsy etc The root contains physiologically active bases and an antibiotic substance pterygospermin

Myristica fragrans Houth (Vern —S -Jatifalam, H, B. & Bo.-Jayphal, M -Jadikkay). The nutmeg B P, B P C, I.P.L & I.P.C.

Grows in the Nilgiri hills and on the Malabar coast Oil is used in the manufacture of pharmacopoeial preparations and aperient pills to prevent griping Nutmegs yield 5 to 15 per cent of a volatile oil and 30 to 40 per cent of fats, also phytostirin, starch, amyloextrin, colouring matter and a saponin. It is largely used for flavouring and as a carminative.

Myrsine africana Linn (Vern Gugul, H. Chapra).

Grows in Afghanistan, salt range and the Himalayas from Kashmir to Nepal at altitudes from 1,000 to 8,000 ft. Berries used as anthelmintic and as a remedy for skin diseases They contain embellic acid 3 per cent. and quercitol 1 per cent No pharmacological work or clinical trials have been carried out

Oldenlandia auricularia Schum Syn -*Hedyotis auricularia* Linn. (Vern.-Gatta colla, B.-Muttia-lata)

Grows in Eastern Bengal and from Nepal, Sikkim and Khasia hills to Assam, Chittagong, Manipur and Burma. Also found in Kanara and Ceylon Used in the treatment of dysentery and diarrhoea Roots contain a fatty oil and glucosides, also reducing sugars, colouring matters, tannin, alkaloids auricularine, heydyotin and albumin No pharmacological work has been done.

Papaver somniferum Linn. (Vern —S -Ahiphena, H & B -Afim, Bo -Aphu, M -Postakatol). The Opium B.P., B P C, I P C & I P.L.

Cultivated in many parts of India especially in Utter Pradesh, Bihar and other parts of the world, China, Asia Minor, Turkey, Persia and Egypt. Dried inspissated juice obtained by incising the green fruit is opium which is used in the treatment of dysentery and diarrhoea as an anodyne sedative and euphoric Opium contains about 25 alkaloids, the important ones being morphine, codeine, thebaine, papaverine and narcotine In addition to these acetic, lactic, sulphuric and meconic acids, gummy and pectinous substances, albumin, wax, fat, resin, etc. also occur.

Opium is a sovereign remedy for relieving pain and producing sleep. Opium and morphine are habit forming drugs and produce addiction

Peganum harmala Linn (Vern --H. & Bo.-Hurmal, B.-Isband, M.-Shimai-azha-vanai-virai) I.P C

Grows wild all over N.W India, Sind, Punjab and Kashmir, Agra and Western Deccan Also found in Arabia, North Africa, Hungary and Spain. Seeds are used as alterative, aphrodisiac, lactagogue and anthelmintic. Contain alkaloids harmine, harmaline and harmalol to the extent of 4% The alkaloids have some anthelmintic action and depress respiration and heart Clinical trials revealed no effect in malaria. The seeds have been used to expel tape worms

Peucedanum graveolens Benth (Vern --S -Shatapuspī, B.-Soolpha, Bo -Balunt-shep, M -Satakuppi)

Grows in central and southern Europe and in tropical and sub-tropical countries. In India, it is cultivated as a cold weather crop. Used as a condiment, carminative, digestive and flavouring agent Largely used in soap manufacture as a perfume Yields 0.06 per cent. of an essential oil

Picroena quassioides Benth. (Vern Bharangi) I P C & I P L.

Found in the outer Himalayas from Kashmir to Nepal and also in China. Bark and leaves used as febrifuge and insecticide. Contains bitter principles quassin and neo-quassin and a bitter alkaloid 0.05 per cent. Quassia is a popular bitter

Picrorhiza kurrooa Benth (Vern --S -Katuka, H & B -Katkī, Bo -Kalikutkī, M -Katuka-rogaṇi). B P.C , I P C & I P.L

Grows in the North West Himalayas from Kashmir to Sikkim at altitudes of 5,000 to 11,000 ft. Used as a bitter tonic and stomachic. Contains 26.6 per cent. of a bitter substance and a glucoside Can be used in medicine as a bitter if properly standardised.

Pimpinella anisum Linn. (Vern.—S -Shetapuspa, H.-Saonf, B.-Muhuri, Bo.-Sonf, M.-Shombu). The Anise B P , B P.C. & I.P C.

Grows in Europe, Russia and Middle East and many other parts of the world In India, found in Uttar Pradesh, the Punjab and Orissa. Used to prevent flatulence and intestinal colic and as an aromatic. It contains an essential oil which is used in medicine to prevent flatulence, intestinal colic and as a flavouring agent.

Pinus roxburghii Sargent Syn.-*P longifolia* Roxb (Vern -Chir, S -Sarala) Chir Pine. B.P , I.P C. & I.P L

Grows in abundance in U.S.A. and France. In India, it is found in the temperate Himalayas from Kashmir to Burma at altitude from 2,000 ft. to 6,000 ft. The oil is used in the manufacture of paints and varnishes and colophony for plasters in medicine The oleoresin yields about 20 per cent. of turpentine oil and about 80 per cent. of residue named as colphony which is used in medicine for preparation of plasters and ointments. Oil of turpentine is used externally as a counter-irritant

and rubefacient. Small doses of oil are given internally in bronchitis and larger doses as an anthelmintic.

Piper betle Linn. Betel leaf. (Vern.-S.-tambula, H., B. & Bo.-Pan, M.-Vettilai) I P C.

It is cultivated in Madras, Central Provinces, Bengal, Orissa, Bombay, U. P. and Burma. Leaves are used as an aromatic, stimulant, carminative, astringent and aphrodisiac. Contain an essential oil, starch, sugar, tannins and diastases. No pharmacological studies have been carried out. Leaves with other substances such as betel nut, lime, catechu, etc. are widely chewed in India. They act as mental stimulant and digestive.

Piper cubeba (Vern.—S -Sugandha muricha, H., B. & Bo.-Kababchini, M -Val milaku) Cubebs. I P.C & B P.C

Found in Java, Sumatra and Malaya Archipelago and cultivated to a small extent in India in Mysore. Fruit is used as a condiment and in the treatment of genito-urinary diseases, cystitis, gonorrhoea and gleet. Contains an essential oil upto 10-15 per cent.

Pistacia integerrima Stew (Vern —Kakra, S -Karkata shingi, H & B.-Kakra shingi, M -Kakkata shingi). I P C

Is found in the sub-alpine Himalayas. It is used as a remedy for cough, phthisis and asthma in doses of 20 grs combined with aromatics. Galls contain an essential oil 1-21 per cent, a crystalline hydrocarbon 3.4 per cent, tannins and gum mastic 50 per cent. No systematic investigations have been carried out. Further clinical trials are needed.

Plantago ovata Forsk. Ispaghula (Vern.—H -Ispaghul, B -Isabgul, Bo.-Isabghol, M.-Ishappukol-virai). B P C, I P.C. & I P L.

Grows in the plains of the Punjab and Sind and low hills from Sutlej westwards. It is also cultivated in Bengal, Mysore and Coromandel Coast. Seeds are cooling and demulcent and are used in dysenteries, diarrhoea and other inflammatory conditions of digestive and urinary tract. Contain a fatty oil, albuminous matter and mucilage. Some species contain a glucoside named aucubin, which is pharmacologically inactive. The mucilage from seeds inhibits growth of common intestinal organisms in the gut. The colloidal nature of the mucilage has a remarkable power of adsorbing bacteria and toxins and is soothing. The seeds are prescribed in chronic bacillary and amoebic dysentery, chronic constipation and chronic diarrhoea in doses of 1—2 ounces soaked in water. The pericarp of seeds separated is more convenient for use and is taken in doses of 2 to 3 teaspoonsful once or twice daily.

Plumbago indica Linn Syn.-*P. rosea* Linn. (Vern.-S.-Chitraka, H., B. & Bo.-Lal chitra).

Commonly cultivated in gardens of India. Root is used as abortifacient and vesicant in rheumatic affections of joint and paralytic conditions. Contains, plumbagin, the active principle, amorphous brown pigment and a reducing sugar. No systematic pharmacological investigations have been carried out.

Plumbago zeylanica Linn. (Vern.—S.-Chitraka, H. & B.-Chita, Bo.-Chitaro, M.-Chittira)

Grows throughout India, particularly in Uttar Pradesh, Bengal and Southern India. Root is used in digestive disorders, piles, anasarca and skin diseases and as abortifacient. It contains an active principle named as plumbagin, which stimulates the central nervous system, heart and plain muscles in small doses. Larger doses have a depressant action. Systematic clinical trials are needed to establish its therapeutic value.

Podophyllum hexandrum Royle. (Vern.—H.-Papra). B.P., B.P.C., & I.P.C.

Grows in the temperate forests of the Himalayas from Sikkim to Kashmir at a height of 7,000 ft. Used as a drastic purgative and as a cholagogue. Contains a resin 8 to 13 per cent, flavonol, quercetin, starch and calcium oxalate. Podophyllum is irritant to the eyes and mucuous membrane. It is an active purgative and is administered in average doses of 0.01 gm.

P. paltatum contains podophylotoxin and α & β paltatins, the former has been shown to be present in *P. hexauctum*, but not paltatins. The resin is effective in destroying soft warts or condylomata.

Pongamia pinnata Merr. Syn -*P. glabra* Vent (Vern —Kanga, S., H. & Bo -Karanja, B.-Dahar karanja)

Grows all over India especially near the Coast, from Central and Eastern Himalayas to Ceylon. The seeds and leaves are used in treatment of skin diseases and rheumatism. Internally the seed oil is used as stomachic, cholagogue, febrifuge and expectorant. It is bitter fatty oil forming 27 to 36.4 per cent. of seeds containing traces of an essential oil. It also contains two crystalline compounds named as karanjin and pangamol. No systematic pharmacological investigations or clinical trials have been carried out.

Premna integrifolia Linn. (Vern.—S.-Ganikarika, H. & Bo.- Arni, B.-Ganiari, M -Munni-vayz)

Grows near the sea from Bombay to Malacca and in Ceylon. Root is used as laxative, carminative and stomachic. The decoction of whole plant is used in the treatment of fevers and rheumatism and neuralgia. Stem bark contains three alkaloids, premine, ganiarine, ganikarine and some unsaturated aromatic hydrocarbons. The alkaloids ganiarine and premine have a sympathomimetic action. Further work is indicated.

Pristimera indica Willd Syn.-*Hippocratea indica* Willd.

Grows wild in forests from Konkan to Madras in South Bengal and Assam. Paste from roots used as a remedy for respiratory troubles. Root contains dulcitol and an antibiotic pristimerin which is effective against gram positive organisms. Preliminary clinical trials revealed that pristimerin is effective against throat affections. Further investigations are in progress.

Psoralea corylifolia Linn. (Vern —S -Vakuchi, H. & B.-Babachi, Bo -Bobawachi, M.-Karpō-Karishi). I.P.C & I.P.L.

Grows all over the plains of India. The seeds are considered anthelmintic, diuretic, diaphoretic and a remedy against skin diseases and leprosy. They contain an essential oil, a fixed oil, a resin and traces of an alkaloidal substance. The oil has an irritant action on skin. It is

toxic to paramoecia and bacteria. Plain muscle is stimulated and arterioles are dilated. An oleoresinous extract made from seeds has beneficial effect in leucoderma when applied locally. Seeds are also taken internally in this affection.

Randia dumetorum Lamk. (Vern.—S.-Madan, H.-Maniphal, B.-Mempal, Bo.-Gelaphal, Maruk-kallan-kai).

Grows throughout India. Fruit is used as an emetic and aperient and as a substitute for ipecacuanha. Contains a small quantity of an alkaloidal substance, Randia saponin and a glucoside. Randia tannic acid and Randia fat are also present. The drug produces irritation of the mucous membrane and haemolysis both in vitro and in vivo. Frog's heart is arrested when perfused with drug solution in concentration of 1 in 50,000. No clinical trials have been carried out.

Rauwolfia canescens Linn

Grows wild in Howrah district near Calcutta in Bengal. Roots used to adulterate those of *R. serpentina*. Root bark stem bark, and leaves contain an alkaloidal body named as rauwolscine. The alkaloid is depressant to the cardio-vascular system. It is also sympatholytic in action and abolishes the pressor effects of adrenaline. Further work is in progress.

Rauwolfia serpentina Benth. (Vern.—S.-Sarpagandha, H.-Chota chand, B. & Bo.-Chandra, M.-Covannamilpori) B.P.C., I.P.C. & I.P.L.

Grows near the foot of hills from Moradabad to Sikkim, in Assam, Pegu, in Southern India along the Ghats, also occurs in Java and Malaya. Root is used as an antidote in snake bites and as a remedy for insomnia, hypochondria and insanity. It contains alkaloids, ajmaline, ajmalinine, serpentine and serpentinine. Extracts containing the total alkaloids, including serpentine, lower blood pressure in animals, ajmaline and serpentinine raise it. Recently a new alkaloid, Reserpine, has been isolated which has a marked sedative & hypotensive action. Doses of 20 to 30 grains of powdered root produce sedative effect and lowers blood pressure especially the diastolic. Considerable work is being carried out on this drug in Europe and America on account of blood pressure lowering properties and sedative effects.

Rheum emodi Wall (Vern.—H. & B.-Revandchini, Bo.-Iadaki-revanda chini, M.-Nattu-ireval-chinni) Indian Rubarb I.P.C. & I.P.L.

Grows wild in the Himalayas in Nepal and Sikkim at altitudes of 4,000 to 12,000 ft. Used in ailments of children and as a purgative. Contains derivatives of anthraquinone to the extent of 2.0 to 4.5 per cent gallic acid, tannin, starch, fat, dextrose and levalose, pectin and calcium oxalate.

Ricinus communis Linn (Vern.—S.-Eranda, H.-Arand, B.-Verenda, Bo.-Erendi, M.-Amanakham chedi) Castor seeds. B.P., B.P.C. & I.P.C.

Cultivated throughout India particularly in Madras, Bombay and Bengal. Seeds contain a fixed oil which consists of ricinoleate of glycerol with a small quantity of palmitin and stearin and a toxic substance ricin. Oil from seeds is used as a purgative.

Rosa damascena Mill. (Vern. — H.- Gulab ka phul, S.- Satapatni, B.- Golapphul, Bo.-Gul, M.-Golappu). The Rose. B.P. & B.P.C.

Chiefly cultivated in Bulgaria. In India it is cultivated in the Punjab, United Provinces, Bihar and Orissa. Rose water is used for eye lotions and the oil as a flavouring agent. Flowers contain a pleasant smelling essential oil used as perfume, and fruits contain Vit. C.

Santalum album Linn. (Vern — S -Swet-chandan, H.-Safedchandan, B.-Sadachandan, S -Mhandanak-kattai). B.P., B. P. C & I. P. C.

Grows wild and is cultivated in Mysore State, Coorg, Coimbatore and Southern parts of Mysore. Wood is used as bitter, cooling, astringent and useful in biliousness, fever and thirst. Paste made from wood as a cooling application for skin affections. The heart wood of the tree yields an essential oil 2.5 to 6 per cent which is used for flavouring purposes and as urinary disinfectant.

Saraca indica Linn. (The Asoka tree) Vern — S -Asoka, H, B. & B.-Asok, M -Asek) I.P.C.

Grows in Bengal, South India, Arakan, Tenasserim and is also cultivated in many parts. Bark is used as astringent, uterine sedative in menorrhagia and as a remedy for piles and dysentery. It contains tannins and catechol. Various fractions isolated from the bark produced no marked action on uterus but faith in its beneficial effects in uterine disorders persists.

Saussurea lappa Clarke Comp. The Costus (Vern — S -Kushtha, H.-Kut, B.-Pachak, Bo -Ouplate, M -Goshtam). I.P.C & I.P.L.

Grows in N.W. Himalayas especially on the slopes of mountains round the Kashmir valley upto an altitude of 13,000 ft. Root is used as hairwash to kill lice and as insect repellent, carminative, stimulant and antiseptic. Root contains an essential oil, 1.5 per cent., an alkaloid named as sausserine, resin 60 per cent, traces of a bitter substance, tannin, inulin 18 per cent, a fixed oil and potassium nitrate. Leaves contain only the alkaloid. The essential oil has strong antiseptic and disinfectant properties. The circulatory system is stimulated and smooth muscle of bronchi and intestines is relaxed. The alkaloid produces relaxation of bronchioles and has a general depressant action in involuntary muscles. The heart is stimulated. An alcoholic extract of root in 1/2 to 2 dram doses relieves asthmatic attacks.

Scilla indica Baker (Vern.—H. & B -Suphadie khus, Bo.-Bhui-kanda, M -Shirunari-vengayam).

Grows frequently in sandy places near the sea in the Deccan Peninsula from the Concan and Nagpur southwards. Used as expectorant, cardiac stimulant and diuretic. Squill contains a number of cardiac glycosides. Clinical trials have revealed that it has beneficial effect on patients suffering from bronchial catarrh and chronic bronchitis.

Semecarpus anacardium Linn. The marking nut tree (Vern — S.- Bhallatamu, H & B.-Bhela, Bo.-Biba, M.-Shayrang)

Grows in Sub-Himalayan tract from Sutlej eastwards down to Assam at an altitude of 3,500 ft. Juice is used in preparation of a varnish, fruit is used as a substitute for marking, counter irritant in rheumatism, sprains

and leprotic nodules Internally juice is used in syphilis, scrofulous affections, piles and nervous debility. The pericarp of the fruit contains 20 per cent. of an oil, which gives bhilawanol and other compounds No systematic investigations have been carried out to determine its beneficial effects in treatment of disease.

Sida cordifolia Linn. (Vern —Kharetī, S -Bala, H -Kungyī, B.-Brela, Bo -Chikana, M.-Chiribenda). I P C.

Grows throughout tropical and sub-tropical India and Ceylon. Root is used as cooling, stomachic, tonic, febrifuge and diuretic The seeds are considered aphrodisiac and are also used in the treatment of cystitis, rheumatism and spermatorrhoea. The seeds contain fatty oil, and an alkaloid (0.085 per cent) recognised as ephedrine, which probably is responsible for its cardiac stimulant effects.

Skimmia laureola Sich & Zucc (Vern -Bassu, Nep.-Chumlani, P.-Ner)

Grows throughout temperate Himalayas from Kashmir to Khasia hills The leaves are used as an incense and burnt near smallpox patients Leaves contain an essential oil with pleasant smell and an alkaloidal substance. No systematic pharmacological studies have been carried out. The essential oil could be used as perfume in soap and cosmetic industry.

Sonneratia acida Linn (Vern -B -Archaka)

Grows in Bengal and the Deccan Peninsula. Used as a poultice in sprains and swellings Juice is used for stopping haemorrhage Contains colouring matter, archin and archinin and a crystalline compound archicine. No pharmacological studies have been carried out

Stephania glabra Roxb. Miers (Vern -Pusha)

Grows in tropical and temperate Himalayas ascending to an altitude of 7,000 ft from Sindh eastwards to Khasia hills and Pegu Root is used as a remedy against phthisis, asthma, dysentery, fevers and intestinal complaints. Tubers contain three alkaloids, gindarine, gindaricine and gindarinine. No pharmacological and clinical studies have been carried out.

Strychnos nux-vomica Linn. (Vern —S -Visha-mushti, H -Kuchla, B -Kuchla, Bo -Kajra, M.-Yetti) B P., B P.C, I P C & I P L

Grows wild throughout tropical India upto an altitude of 4,000 ft in Orissa, Madras, Cochin, Travancore and the Coromandal Coast Seeds are used in the treatment of dyspepsia and diseases of the nervous system Fruit, bark, leaves and wood contain strychnine, brucine, vomicine, etc Seeds contain 1.53 to 3.42 per cent. of the total alkaloids of which half is strychnine

Swertia chirata Ham. (Vern. — S -Kirata, H -Charayatah, B -Chireta, Bo.-Chiraita, M.-Nila-Vembu) Chiretta. B.P C., I.P C & I.P.L.

Grows abundantly in the temperate Himalayas from Kashmir to Bhutan and Khasia Range between altitudes of 4,000 to 10,000 ft. Used as a bitter tonic, stomachic febrifuge and anthelmintic. Contains a bitter glucoside chiratin, resins, tannin and ash 4 to 8%.

Symplocos racemosa Roxb. (The Lodh tree) Vern —S -Lodhra, H. B. & Bo -Lodh, M.-Ludduga). I.P.C.

Grows in plains and lower hills of Bengal, Assam and Burma. It is also found in dry forests of Chhota Nagpur plateau. Bark and leaves are used for dyeing purposes. The bark is used as astringent in bowel complaints, eye diseases and ulcers. Bark contains three alkaloids, loturine 0.24 per cent. collutrine 0.02 per cent and loturdine 0.06 per cent. No systematic pharmacological investigations or clinical trials have been carried out.

Terminalia arjuna W. and A. (Vern.—S & Bo -Arjuna, H & B -Arjun, M.-Vellai-maruda-maram). I P C.

Grows throughout the Sub-Himalayan tracts of United Provinces and in the Deccan, Southern Bihar, Chhota Nagpur, Burma and Ceylon. Bark is used as a cardiac tonic in heart diseases and contains large quantities of calcium salts, about 12 per cent. tannins, an organic acid, an ester and some colouring matters and sugar. It is reputed to have a stimulant action on heart. Clinical trials however showed no beneficial effects in heart disease. Further investigations needed.

Thalictrum foliolosum D C. (Vern —H -Pinjari, P -Gurbani, Bo.-Mamiran).

Grows in Khasia hills between altitudes of 4,000 ft to 6,000 ft. It is used as a bitter tonic and laxative. It is a cheap but valuable substitute of Mamira used in preparation of eye solves. It contains two alkaloids berberine and thalictrine. No proper investigations have been carried to study its efficacy in conjunctive affections.

Thevetia peruviana Schum Syn -*T. nerifolia* Steud. See poisonous plants of India (Vern -H & Bo -Pila-kaner, B -Kolka-phul M -Pachch-ai-alari).

Grows in plains all over India, but originally a native of west Indies. Seeds are used as abortifacient and as poison for man and cattle. They contain 57 per cent of a colourless oil, which yields glucoside thevetin which is not toxic to unicellular organisms. The drug is toxic to higher animals and is not likely to be of therapeutic value because of powerful toxic effect on heart muscle.

Tinospora cordifolia Miers (Vern — Giloe, S -Guduchi, H & B -Gulancha, Bo -Gulwail, Bo -Gulwail, M -Sindilkodi) I P C & I P L

Grows in Uttar Pradesh. It is used in treatment of rheumatism, hyperacidity and dyspepsia and as a tonic, antiperiodic and diuretic. Stems contain glucoside giloin and non-glycosidic bitter, gilenin and gilosterol. No Pharmacological investigations or clinical trials have been carried out.

Toddalia asiatica Lam. Syn -*T. aculeata* Pers. (Vern —S -Kanchana, H -Kanj, B.-Koda-todali, Bo.-Jun-li-kali-mirchi, M Milkaranai).

Grows in the Nilgiris and sub-tropical Himalayas from Kumaon to Bhutan upto an altitude of 3,000 ft. Root bark is considered to be antimalarial. The plant contains two alkaloids, todaline and todalinine. Leaves contain an essential oil and root bark contains an essential oil;

Vitex peduncularis Wall. (Vern —H.-Nagbail, B.-Goda, M.-Navaladi).
I.P.C.

Grows in Bihar, Eastern Bengal and Madhya Pradesh. The leaves and roots are used in the treatment of malarial and black water fevers. Leaves contain a glycoside. Clinical trials with infusion of leaves showed no beneficial effect in malaria or black water fever.

Withania somnifera Dunal (Vern —S., B. & Bo -Ashwagandha, H -Asgandh, M -Amku lang-kalang) I.P.C.

Grows throughout the drier parts of India, Baluchistan and Ceylon. Leaves and roots are used in the treatment of tumours, tuberculous glands, bronchitis, rheumatism and dyspepsia. The plant contains pot. nitrate, tannin, colouring matter, glucose and an alkaloid. It has a mild sedative effect on animals. Deserves further study.

Xanthium strumarium Linn (Vern —S -Aristha, H Chhota-gokru, B -Bon-okra, Bo -Shankeshvara, M -Marlu-mutta) I P C

Grows throughout the hotter parts of India upto an altitude of 7,000 ft. The whole plant is used as diaphoretic and sedative. Root is used as bitter tonic and in the treatment of cancer and strumous diseases. It contains fat 38.6 per cent., albuminoids 36.6 per cent and a glycoside Xanthostrumanin 1.3 per cent. The glucoside is physiologically inactive. No clinical trials have been carried out.

Zingiber officinale Rosc. (Vern —S.-Adrakam, H -Adrak, B -Ada, M.-Inji) B.P & I.P C

Cultivated in Cochin, Bengal, Bombay, Kumaon and many other parts of India. Used as a condiment in food and as carminative and stimulant of the gastro-intestinal tract. Contains 0.25 per cent. of a volatile oil of light yellow colour, resinous matter, starch and mucilage.

CHAPTER VII

I. TOXICOLOGICAL ASPECTS OF INDIAN PLANTS

The second line of basic research has been in connection with the Poisonous Plants of India including the group of insecticides, insect repellent etc. Here we will briefly refer to the present position with regard to the toxicological aspects of Indian Plants.

(I) CRYPTOGRAMS OR THE FLOWERLESS PLANTS

The toxicological aspects of the Cryptogams are little known so far as India is concerned

(a) **Bacteria.** The bacteria are among the simplest form of plant life and are met with universally. The majority of them are harmless, but some are injurious to man and animals. They produce deleterious effects in two ways — Firstly as parasites, when they derive nourishment from living animals and many of them produce, within the body, toxins which are harmful. Secondly many saprophytic bacteria produce poisonous substances, especially such as those occurring in putrid flesh, fish and other decaying organic matter. It is not our intention to give details of toxic properties of this group. These are fully discussed in books on bacteriology. Bacteria are a class by themselves and although they belong to vegetable kingdom do not come under our purview here.

On the other hand a number of organisms of this group produce antibiotic substances which are of very great value in the treatment of disease. Streptomycin has been derived from *Streptomyces griseus* (*Actinomyces griseus*), Aureomycin from *Streptomyces auriofaciens*, Terramycin from *Streptomyces rimosus*, Chloromycetin from *Streptomyces venezuelae*, Bacitracin from strains of *B. subtilis*, etc.

A large amount of work is now being done in this country on the isolation of antibiotic substances from various soil and other micro organisms found in this country.

(b) **Algae** The algæ that cause poisoning are mostly those which are found in stagnant waters. The normally offensive odour may be sufficient to indicate their presence, but only a microscopic examination can determine just what the forms of algæ present may be. Blue-green algæ, as a group, are perhaps the most pronounced in their toxic effect. Parker and other workers have shown that when odours in water are pronounced, these microscopic organisms are present in considerable numbers. He claims that of the organisms which produce objectionable and deleterious qualities in water, the microscopic ones are the more important and very few cases have been observed in which really serious trouble in water supplies could be attributed directly to the growth of larger plants. In any study of the algæ from this point of view, however, account must be taken of the products of decomposition by the associated bacteria, since poisoning may be produced by the toxins produced by bacteria rather than by the algæ.

Certain algæ, such as *Microcystis flos-aquae* (Witttr.) Kirch, *Aphanizomenon flos-aquae* (Linn) Ralfs. and species of *Anabaena*, etc. form on the surface of water what is generally called water bloom. The presence of water bloom on the surface of lakes, ponds, and other open sheets of water is distasteful to bathers and obnoxious to those living in the vicinity. Livestock compelled to drink water containing water bloom are reported to have suffered from poisoning. In Minnesota, (U.S.A.) during recent years, horses, cattle, sheep and donkeys have died in large numbers on the shores of lakes where water bloom is present. All the above mentioned algæ forming water bloom have been recorded in various parts of India but no work has been done in connection with their toxic effects. According to Dr Bhardawaja of the Benaras Hindu University, water blooms containing these species occur commonly on the surface of many temple tanks in different parts of India. Of the other possibly harmful algæ may be mentioned species of *Nodularia*, *Clathrocystis*, *Nostoc*, *Oscillatoria*, *Pandorina*, and *Volvox* when present in large numbers.

The question of growth of algæ in water reservoirs leads us to a very important public health problem. Although in India very little information is available about the contamination of the water supplies with the group of toxic algæ, the important question of checking their growth in the reservoirs of water supplies needs study. One of the essentials of the algal growth is light. This growth may, therefore, be prevented, or at any rate considerably reduced, by covering up the reservoirs and cutting off sunlight. Most of the reservoirs for the supply of water to both animals and man, however, in India are generally not covered and are often largely contaminated with algal growth. The removal of organic matter by keeping the source of water supply in as pure a state as possible will not doubt keep down the algal growth. It must be understood, however, that nearly all water contains sufficient organic matter for the growth of algæ, especially the water coming from water-sheds. Growth of algæ can also be successfully prevented by additions of copper sulphate in dilutions of one in five millions or even more. Such dilutions do not render the water deleterious to man and animals. The problem of toxic algæ is important and deserves the attention of workers in this field. It is also possible that some of these algæ may contain active principles having therapeutic properties in the treatment of disease. This aspect has yet to be investigated.

(c) **Fungi.** (i) Some fungi live on the skin and mucous membranes of man and animals and cause various, diseases, e.g. ringworm, thrush, etc.

(ii) There are others which attack foodstuffs and among these may be mentioned —

(1) **Smuts.** Many of these are destructive parasites which invade plants of vital economic importance on account of their food value, such as oats, millet and other cereals. Some are supposed to be poisonous if taken in large quantities, and others are said to produce irritation of the mucous membranes. There is difference of opinion with regard to the injurious effects produced by particular kinds of smut and hardly any authentic information is available regarding those occurring in India. The subject deserves the attention of mycologists.

(2) **Rusts.** Annual recurrence of the outbreaks of rust attacks of cereals in India, especially those attacking wheat, is of great economic importance to the country. These, especially the uredo stage, produce

inflammation of the mucous membrane of the mouth and nose. The dust coming from the infested straw when the grain is thrashed is stated to cause serious disturbances of the respiratory tract of man and animals. Very little information is available about the Indian strains.

(3) **Ergot**, which grows on rye, is a well known example of a fungus which produces highly poisonous substances. Some species of this fungus grow on wild grasses in India and Ergot itself has been grown in this country for medicinal purposes.

(4) The poisonous nature of the seeds of darnel (*Lolium temulentum* Linn.), a grass and annual weed cultivated especially in northern India, is believed to be due to a fungus, and cases of poisoning due to admixture of the seeds with wheat grains are not infrequently reported in India and abroad. Cases of death among livestock have also been reported.

(5) **Moulds**. Very variable data are available as regards the poisonous effects of mouldy food stuffs in India, but there appears to be little doubt that the presence of certain species may occasionally produce harmful effects in man and animals. Species of *Mucor*, *Aspergillus*, *Penicillium* and *Fusarium*, etc. deserve special investigation in this connection. It appears, however, that there is an appreciable difference in the susceptibility of different species of animals to the effects of mouldy food-stuffs. In general it has been stated that horses, dogs and pigs are more susceptible than ruminants and poultry, while in other animals the case may be the reverse. Very little information is available about the toxicity of moulds occurring in India and the problem requires a thorough investigation because of its great economic importance. In the meantime it would be safer to consider all fungus-infected food-stuffs as deleterious. Acute poisoning with the moulds is rarely met with and if they are taken in small quantities there is hardly any danger. Mouldy food should however be avoided. It is worthy of note that some of these fungi are the source of powerful antibiotic substances such as penicillin from *Penicillium notatum*, fumigaen from *Aspergillus fumigatus*, calvicin from *Aspergillus clavatus*, etc. A large amount of work is now being done in this country in connection with antibiotics from local strains of these organisms.

(iii) **Mushrooms or fleshy fungi**. The third group of the poisonous fungi belongs to the mushroom class. A number of these are edible and many occurring in India are indiscriminately eaten by people and if properly cooked few produce serious toxic effects. Cases of fungus poisoning, however, are not infrequently met with, particularly in the hills. Unfortunately very little information is available about the poisonous fungi growing in this country and in spite of numerous cases of poisoning, little attention has been paid to the subject.

Stropharia semiglobata (Batsch) Quel. from Khasia hills, *Hypholoma fasciculare* (Huds.) Fr. from Darjeeling and Simla and *Lactarius vellereus* Fr. from Sikkim are regarded as poisonous. There is also evidence on record that there exists in Bengal a fungus which closely resembles an edible variety but which contains amanitine or muscarine, the poisonous principle of the foreign *Amanita muscaria* Pers. There are probably many more poisonous species that have really been incriminated as poisonous, but on the whole their number may be small and indeed if properly cooked

only a few are dangerous. If washed in water and macerated in vinegar before cooking, and if eaten with plenty of bread there is almost no danger in most cases. The safest method, however is to learn to recognize the edible species and never to eat a fungus until its identity is certain.

Some of the foreign poisonous fungi, e.g. *Lepiota cristata* Quel. *Volvaria gloiocephala* Gill, *Amanita muscaria* Pers. and *Amanita phalloides* Secr. are well known. The last-mentioned is responsible for perhaps 90 per cent. of the deaths caused by fungus poisoning in Europe, Great Britain and U.S.A. During the World Wars, when food scarcity became acute in Germany and Austria, poisoning from fungi appreciably increased.

According to Ford there are four main types of mushroom intoxication —(1) Gastro-intestinal type in which the attack ceases when the stomach is emptied (2) General catharsis which is painless. (3) Violent vomiting and pain but no involvement outside the gastro-intestinal tract. (4) Choleraform type producing widespread degeneration of cells. With regard to edible fungi it may be stated here that the nutrition value of mushrooms is small, their chief value being their flavour and the feeling of satiation they produce. The following is a list of mushrooms met with in India :—

Agaricus campestris Linn.—The Mushroom; (Sans.—Chattrak; Kash—Manskhel, Beng.—Banger chhata, Sant—Ot: Bomb.—Alombe). Generally in damp debris throughout India during rainy season; universally eaten fresh or dried.

Amito pers. Kurrum.

A. ostreatus Jacq (Cutch & Bomb.—Phanasa-alambe, or vulgarly phansamba) Grows upon stumps of old jack-trees (phanas).

Cantharellus cibarius Fr Kashmir, Peshawar, Mussoorie.

Collybia albuminosa (Berk) Betch. (Syn -*Lepiota albuminosa* Berk). (Beng—Durga chhata) Bengal, C P, and Berar Grows from inside the termites, nests, eaten with relish.

Coprinus comatus (Battara) Fr The Mushroom; Hindi. & Punj.-Khumbi, Khumb) Punjab, Uttar Pradesh and several other parts of India. Eaten fresh or dried Collected during rainy season.

Entoloma microcarpum Berk & Broome. (Beng -Wee-chhata). Bengal. Grows on the surface of outer crust of termites' nests, commonly eaten by villagers.

Fistulina hepatica Fr. Darjeeling.

Helvella crispa Fr. Common in Afghanistan

Hirneola polytricha Mont. (Syn.-*Exidea polytricha* Mont.). Belgium, Poona, Dharwar, Nidungayam, Malabar and Burma.

Hydnum caralloides Scop Darjeeling 7,500 ft., Chitral, N.W.F.P. (common) and Afghanistan. In crevices of old tree-trunks collected during August; dried in the sun and largely used.

H. repandum Linn. Mussoorie, Uttar Pradesh

Lactarius deliciosus Fr. Sikkim.

Lentinus subnudus Berk Common in Bengal, Kadala and Bombay.
On dead branches of logs Eaten by Kholes fresh and young.

Lepiota mastoidea Fr. Bengal.

L. procera (Scop) Sacc Saharanpur.

Lycoperdon sp. Puff-balls Bengal, Kashmir and many parts of Western Himalayas.

Malanogaster durissimus Cooke Truffle. Simla (abundant), Kangra.
Occasionally eaten.

Morchella esculenta Pers. The Morell, (Punj-Guchhian) Plains, kana kach hills. Fleshy fungus abundant in Kashmir, Chamba and many parts of Northern Punjab It appears on hills as snow melts in early spring. It is dried and is eaten with much relish.

Pleurotus cretaceus Massee (Vern -Dhingri). Peshawar and Madhya Pradesh. On wood.

P. fimbriatus Bolt C P. and Berar

Polyporus squamosus (Huds) Fr Darjeeling, 7,500 ft., Pangl, N.W. Himalayas. On dead wood.

Truffles. Stewart describes some being found in Kashmir Badhwar has recently collected some blackish-brown ones from the Kagan valley locally known as 'usri'. They are highly flavoured and their presence in the soil is discovered by the villagers by smell in September-October when they are said to develop the flavour best Goats are also said to dig out some during grazing and eat them. (Also see *Melanogaster durissimus* above).

Volvaria diplasia Berk & Broome (Beng -Pawal-chhatta) Bengal, Burma.

V. terrestria Berk & Broome. (Beng. - Poal-chhattea) Bengal
Grows on heaps of waste paddy straw

In addition Stewart mentions another species as being freely eaten in the Punjab, which is known as 'shirian' in the Jhelum and 'batbakri' in the Kair valley It is a thin, flat ragged-looking fungus, yellow above and with white gills below, which is found on dead trees in various parts of the Punjab Himalayas at altitudes of 8,000 to 8,500 ft. He also mentions an 'underground mushroom' of doubtful species found near Multan called 'boinphal'.

So far as the Poisonous Mushrooms are concerned little information is available. The following is a list of mushrooms which are known to be poisonous :-

POISONOUS FUNGI

Very little information is available regarding the Indian poisonous fungi. From time to time cases of fungus poisoning are reported but, it is to be regretted that little or no attention has been paid to the subject.

Stropharia semiglobata (Batsch) Quel from Khasia Hills

Hypholoma fasciculare (Huds). Fr. from Darjeeling and Simla and *Lactarius vellerens* Fr from Sikkim, are regarded as poisonous. There is also evidence on record that there exists in Bengal a fungus which closely resembles an edible form but which contains amanitine or muscarine, the poisonous principle of *Amanita muscaria* by eating which, symptoms closely resembling those of intoxication rapidly ensure. Furthermore, *Mucor* has been regarded as a harmful in India since ages, and the pickles and all edible stuff attacked by it are not thought fit for eating.

There are, however, some foreign fungi which are definitely reported to be poisonous.

Amanita phalloides The death cap is responsible for perhaps 90 per cent of the deaths caused by fungus poisoning in Europe, England and U S A. It is the most dangerous fungus known and very small quantities will cause intense suffering and often death. There are indeed, several other species of the genus that are very poisonous, e g , *Amanita muscaria* - fly agaric and *A. pantheriana* - warted agaric, etc which are intensely poisonous.

Lepiota cristata - crested agaric, and several other small species of *Lepiota* are regarded with suspicion and should be avoided. *Volvaria gloiocephala* - glutinous agaric and its allied mushrooms have always been regarded as poisonous, but there is recent evidence that they may be eaten without ill effects.

Psalliota xanthoderma - yellow staining mushroom has caused illness in some cases.

(d) **Lichens**. Very little is known about these symbiotic organisms which consist of algal cells enveloped by the mycelium of the fungus forming a felted mass. Although this group is not to be regarded as a serious menace to livestock, cases of poisoning due to *Parmelia* and *Cretaria* species, etc are mentioned in foreign literature. *Parmelia molluscula* has been said to effect sheep and cattle, producing lack of coordination of the hind limbs. In more severe cases the animal lies down and is unable to move either its front or hind limbs. Little or no information is available about lichens in India and even their systematic botany has not been sufficiently worked out.

(e) **Bryophyta** (Liverworts and Mosses). This is the least-known group of plants from the view-point of poisoning and there is little to be said about it.

(f) **Pteridophyta** (Vascular Cryptogams). This group includes ferns and allied plants but unfortunately little or no work has been done in India with regard to their toxicity. Greshoff and others have reported

the presence of hydrocyanic acid in a number of ferns, especially when young. References to the supposed poisonous properties of the bracken fern (*Pteris aquilina*) have appeared in the literature for a long time, and Stockman in great Britain showed that it is poisonous to cattle when eaten in considerable quantities. The plant is found in India. *Aspidium filix-mas*, the male fern, is suspected of being poisonous. The roots are used in medicine and large quantities of it produce haemorrhagic gastro-enteritis, tremors, weakness, stupor coma, acute nephritis, and cystitis. Six drachms of the oleoresin have proved fatal in man and three ounces in the cow. This fern is not found in India, but there are several other foreign species of *Aspidium* which are also suspected of being poisonous. The present authors have examined Indian representatives of these plants. Some foreign species of *Osmunda*, *Davallia* and *Adiantum* are also suspected of being poisonous and medicinally active. The following ferns growing in the North Western Himalayas have been examined and found to be active as anthelmintics.

<i>Name of Ferns</i>	<i>Locality</i>	<i>Active principles</i>		
<i>Dryopteris odontoloma</i>	Kashmir & Mussoorie	Up to B P & U S P standard		
<i>D. barbigera</i>	Kashmir	do	do	do
<i>D. marginata</i>	Kashmir	do	do	do
<i>D. blanfordii</i>	Chamba	do	do	do
<i>D. schimperianum</i>	Mussoorie	do	do	do
<i>D. calcarata</i>	Mussoorie	Not up to standard		
<i>D. marginata</i>	Mussoorie	Up to standard		

Some of the foreign species of *Equisetum* (horsetail) have long been recognized in foreign countries as injurious to cattle and horses. They produce an intoxication in which the animals stagger about and wander aimlessly. There is no information available in India with regard to the Indian horsetail *Equisetum arvense*, but several European and American workers are convinced that it is definitely poisonous to horses, while others hold a contrary opinion. This plant grows commonly in certain places in India where it might be a menace to livestock.

CHAPTER VIII

II. TOXICOLOGICAL ASPECTS OF PHANEROGAMS (THE FLOWERING PLANTS)

After having given a very brief survey regarding the toxicological aspects of the Cryptogamic flora we will now take up the Phanerogams or the flowering plants. Economically this is the most important group both for man and animals from the point of view of everyday necessities of life, e.g., food, medicines, etc. It is probably on account of this that more information is available with regard to this group.

From a toxicological point of view the Phanerogams may be divided into two main groups.

I PLANTS POISONOUS TO MAN AND LIVESTOCK.

(a) **Medicinal Plants** — Primitive man in the quest for food must have come across plants containing poisonous principles by accident and by experience must soon have learned to avoid them. He even made use of them for the purpose of fighting against his enemies and for procuring his food by killing animals with them. On the whole, our knowledge is fairly well advanced so far as the relationship of poisonous plants have been used for criminal purposes from ancient times but the majority of them are used as medicinal agents for the amelioration of human suffering. It is well known that many plants that are harmful to life in large quantities, produce remarkably beneficial effects in small regulated doses in disease conditions. From the economic point of view, the abundance of this group of plants in our midst is of very great importance in as much as it provides us with medicinal agents of every description, not only sufficient for our own use but also for purposes of export. We have already given brief review of the work done on this group.

(b) **Poisonous to livestock.** The second important aspect of these plants is in connection with other countries, our knowledge is very meagre. In India, there are hundreds of plants that are intimately connected with the food supply of the bovine population such an essential factor in connection with agriculture and food supply. The fodder required for this livestock amounts to about fifty million maunds daily (excluding the concentrates that are in use). Even in its present unsatisfactory condition, the cattle industry contributes very largely to the annual agricultural income of this vast country. Its welfare is therefore of the utmost importance. There is no doubt that the mortality among cattle through poisoning in India is very large. Unfortunately no figures are available of the loss suffered on this account but it must be enormous.

Even in a country like the United States of America where knowledge with regard to Poisonous Plants is well advanced as compared to India, great losses are incurred every year through the poisoning of cattle. The loss occurring in this sub-continent can thus be imagined.

Though the number of plants which are known to have markedly poisonous properties is perhaps small as compared with the total species included in the Indian floras, there are many which are of common occurrence and which no doubt produce serious losses by death or illness among sheep, cattle and other domestic animals. The toxic effects produced may be indicated by reduction in the yield of milk, the milk may become unpalatable through excretion in it of toxic products, or it may even become unfit for consumption. Some time even the meat of these animals becomes poisonous.

It may be stated here that animals do not instinctively select toxic plants as forage, that all classes of livestock are not necessarily equally susceptible to the same poisonous plants, that not all poisonous plants are dangerous from their initial appearance up to maturity and that in some the animals do acquire a depraved appetite for harmful plants, especially when the fodder supply is scarce, a condition which is of frequent occurrence in many parts of India. The losses in many cases may be avoided by increasing our knowledge about these plants.

Prevention. The question arises as to what should be done to prevent poisoning by plants. The adage 'prevention is better than cure' is applicable to the problem of plant poisoning with just as much force as in other spheres. Often cases are brought to notice when the symptoms have developed and the poison has already circulated in the blood stream and has done irreparable damage to the system. Increased knowledge of the poisonous plants is the first step in this direction and this is sure to have an effect in decreasing fatalities among human beings and livestock. It is for this reason that Indian Council of Agricultural Research and Indian Council of Medical Research have encouraged research on this group. Keeping the animals always as far as possible from dangerous areas and exercising special care during periods of drought are likely to decrease the mortality amongst livestock. Eradication of poisonous plants is a difficult matter, involving an enormous amount of labour and capital, but wherever and whenever possible it should be resorted to. This depends upon the habits of the particular plant. Such plants may be annual, biennial or perennial herbs, or shrubs or trees. Annuals should be pulled out or dug out before seeding and biennials may be dealt with as the annuals. Perennials are propagated both by the seeds and by the underground organs, such as tubers, rootstocks, bulbs, etc and may be dug out if not deeply rooted. Shrubs are woody perennials and should be cut down or dug out. Cutting down of lower branches of trees within the reach of animals or children is advocated.

The indiscriminate importation of ornamental plants has recently increased the number of poisonous plants in India. Some of these do not find much competition in their adopted home and are likely to spread in this country at an enormously rapid pace. The time is now ripe to agitate for a law prohibiting the importation of poisonous plants for gardens or to take measures to forbid the cultivation of those already introduced. The foodstuff dealers should make sure that adulteration is not practised either with poisonous plants or with plants whose properties are doubtful. Recent work in connection with the causation of epidemic dropsy has shown that in some epidemics, mustard oil adulterated with 'katakari' oil from the seeds of *Argemone mexicana* Linn, the Mexican poppy or 'Shialkata', was the cause of the outbreak.

Food Poisons. There is a number of common articles of food which produce poisoning in man and animals and a brief reference may be made to these.

(1) **Khesari dal, *Lathyrus sativus* Linn**, an important article of diet in man and animals, has been responsible for a large number of cases of poisoning under certain conditions in man, cattle, sheep, pigs, horses, pigeons, ducks etc. Examples of lathyrism in man in the form of spastic paralysis are commonly seen every day in the streets of Calcutta and towns in Bihar and its toxic effects in horses and cattle are well known. Moderate amounts of this pulse can be taken with impunity, but large amounts, especially to the exclusion of other fodders or foods, produce poisoning.

(2) Grasses (Gramineae) form an important part of the food of animals. Some of these develop dangerously large quantities of hydrocyanic acid under certain climatic and soil conditions especially at times of drought or when the plants are wilting, stunted or young. Unfortunately our knowledge of Indian grasses in this connection is meagre and it is not possible to estimate the losses in livestock from this source.

The common Jowar (*Sorghum vulgare* Pers), the Indian millet, is largely cultivated as fodder for cattle and also for human food. It has caused serious outbreaks of poisoning among livestock when wilted or stunted under drought conditions. *Sorghum halepense* Pers, a tall perennial grass, with creeping rhizomes and numerous suckers, known as Johnson grass, grows all over India under the name of 'baru' in Hindi and 'Kala-mucha' in Bengali. It has been responsible for serious losses among livestock during recent years in the North-Western parts of India. It has been stated that the amount of hydrocyanic acid in these plants decreases with the age of the plant but never entirely disappears. The points to be remembered about these grasses are that they are dangerous during wilting and under conditions of draught, that the younger and more succulent ones are often more likely to contain lethal doses of hydrocyanic acid and that if well dried, these plants are generally not dangerous. The toxicity in the case of cyanogenetic compounds depends on the quantities of hydrocyanic acid liberated, and according to the amount and speed at which they are eaten. Often such large quantities are given that the animal will die before any veterinary aid can be given. The only remedy is prevention. The problem of poisonous grasses is of great economic importance in certain parts of India where rains often fail and drought conditions prevail.

The workers of the Drug Research Laboratory have been engaged in carrying a survey of grasses occurring in the North Western Himalayan Region. About 150 different species have been collected and are now being studied with regard to their nutritive values and possible poisonous properties. A number of poisonous grasses have been discovered which the cattle of these parts instinctively avoid, but new-comer gets poisoned. *Stipa siberica* commonly growing on hills round Kashmir Valley is an example and there are others.

(3) The linseed plant, *Linum usitatissimum* Linn., contains a cyanogenetic glucoside, the maximum amount of which is reached very early in the development of the plant and finally disappears, except in the seed, which

3. **Adonis aestivalis** Linn , *A. chrysocyathus* H. f & T.
These are poisonous to animals and act as poison to heart.
4. **Anemone obtusiloba** D. Don. (Vern - P.-Rattanjog)
It is a vesicant and when taken internally produces vomiting and purging, drying alters properties.
5. **Aquilegia vulgaris** Linn
It is poisonous to animals.
6. **Caltha palustris** Linn.
It is acrid and poisonous and deaths among horses are reported.
7. **Cimicifuga foetida** Linn (Vern - P -Jiunti)
It is a heart depressant and acts as insect repellent.
8. **Clematis gouriana** Roxb , *C graveolens* Lindl , *C napaulensis* DC. (P.-Oandak), *C orientalis* Linn , *C triloba* Heyne (Vern - S.- Laghukarni, H & Bo- Moravela) *C wightiana* Wall.
These are irritant and produce blisters, such properties are altered by drying
9. **Delphinium brunonianum** Royle (Vern - P -Laskar), *D. caeruleum* Jacq (Vern - P.-Dhakangu), *D. elatum* Linn , *D. vestitum* Wall
These are cardiac and respiratory depressants, acrid in taste, insecticidal and poisonous to animals
10. **Nigella sativa** Linn. (S -Krishnajiraka, H & B.-Kalajira, Bo - Kalenjire, M -Karun-shiragam)
Produces abortion in larger doses.
11. **Paeonia emodi** Wall Vern.- H.-Udsalap, P - Mamekh).
It is said to have narcotic properties
12. **Ranunculus arvensis** Linn (Vern - P - Chambul), *R. falcatus* Linn , *R. laetus* Wall , *R lingua* Linn , *R pensylvanicus* Linn. f , *R. sceleratus* Linn (Vern - Pers.- Kabiraj).
These are vesicant and poisonous to livestock when fresh, drying alters these properties.

Magnoliaceae

(Magnolia and Champa Family)

Shikimin, illicin, essential oils.

13. **Illicium griffithii** Hk. f & T., *I anisatum* Linn., (*-I religiosum* Sieb. & Zucc.) (Vern.- H -Anasphal, Bo.-Badian, M.-Anashuppu)

This is the star anise of China (*I verum* Hook.f.) which was imported into India sometimes adulterated with *I religiosum* , it has produced deaths. The latter is respiratory and cardiac poison. Indian *I. griffithii* is also referred to as being poisonous

Annonaceae

(Custard apple Family)

Resin, alkaloid, etc.

14. **Annona reticulata** Linn (Vern - H.-Lona, B -Nona, Bo.-Ramphal, M -Ramsita), *A squamosa* Linn (Vern - S.- Gandhagatra, H - Sitaphal, B -Ata, M - Sitapalam).

The seeds are intensely irritant to the conjunctiva; applied locally they act as abortifacient They have insecticidal properties. Roots act as drastic purgative.

Menispermaceae

(Moonseed Family)

15. **Anamirta cocculus** (Linn) W. & A (Vern - S -Kakaphala, H. & B - Kakmarı, M -Kakkay-kolli-vıraı)

It is a convulsant poison to animals. It acts as an insecticide and it is also used to poison fish and cattle.

- 16 **Pachygone ovata** (Poir) Miers

It is an insecticide and piscicide

Berberidaceae

(Berberry Family)

Berberine, podophyllum resin

- 17 **Berberis aristata** DC (Vern -S -Daru harıdra, H -Dar-hald (and probably few more species)

It is poisonous to lower animals and is a fish poison

- 18 **Podophyllum hexandrum** Royle (*P emodi* Wall. ex Hk f et T)

It contains a resin which is a drastic purgative and is irritant to the mucous membranes

Papaveraceae

(Poppy Family)

Morphine, codeine, papaverine, berberine, argemone oil, etc

- 19 **Argemone mexicana** Linn (Vern - S -Srıgala-kantaka, H & B - Sıalkanta, M -Bırama-dandu)

The oil expressed from its seeds is occasionally mixed with mustard oil which has been held responsible for producing symptoms resembling epidemic dropsy

- 20 **Meconopsis aculeata** Royle (Vern - Sımla - Kanta), *M. napaulensis* DC.

The roots are considered to have narcotic properties.

21. **Papaver dubium** Linn., *P. nudicaule* Linn , *P. rhoeas* Linn. (Vern - S -Rakra-posta, H -Lalpost, Bo.-Janglimudrika, M.- Shivappu-postaka chedi), *P. somniferum* Linn (Vern.- S.-Ahiphena, H. & B.-Afim, Bo -Aphu, M.-Postakatol)

All species yield opium more or less but *P. somniferum* is the chief source. Opium has been used for suicidal purposes.

Cruciferae

(Mustard Family)

Glucocides on contact with water produce vesicant active essential oils

- 22 **Brassica cernua** (Thunb) Forbes et Hemsl, *B integrifolia* (West) O E Schulz, *B juncea* (Linn) (Czernjaew et cosson (rai), *B napus* Linn with four varieties (Toria, sarson), *B nigra* (Linn.) Koch (black mustard).

Seeds of these are vesicant, and mustard cakes if fed in large quantities and over prolonged periods is harmful to cattle. Sarson cake is safest, mixture with rai or black or white mustard dangerous.

- 23 **Lepidium draba** Linn (Vern - Afgh - Bijindak)

It is a fish poison

- 24 **Sinapis alba** Linn (white mustard)

It is discussed under Brassica

Capparidaceae

(Caper Family)

Essential Oils

- 25 **Capparis decidua** Edg (*C aphylla* Roth) (Vern - S - Karira, H - Karer, P Karia, Bo - Kari, M Karyal)

It acts as a vesicant

- 26 **Cleome felina** Linn f *C viscosa* Linn, (Vern - S - Swarnakshira)

It acts as a vesicant

- 27 **Gynandropsis gynandra** (Linn) Merr (*G pentaphylla* DC)

It is an insecticide and a piscicide. It has vesicant properties

Bixaceae

(Chaulmoogra family)

Cyanogenetic glucoside, chaulmoogra oil.

- 28 **Gynocardia odorata** R Br (Vern - H., B. & Bo - Chaulmoogra)

The fruit acts as a piscicide.

- 29 **Hydnocarpus kurzii** (King) Warb. (*-Taraktogenos kurzii* King), *H. laurifolia* (Dennst.) Sleumer (*-H. wightiana* Bl)

The fruit acts as a piscicide and the seed oil is gastro-intestinal irritant.

Polygalaceae

(Milkwort Family)

Saponins.

30. **Polygala chinensis** Linn. Vern -H -Meradu, Bo -Negli), *P. crotalariaoides* Buch -Ham (Vern -Santh -Lil kathi), *P. telephioides* Willd.

It has an acrid taste and is an emetic. It has expectorant properties.

Caryophyllaceae

(Carnation Family)

Saponins.

31. **Saponaria vaccaria** Linn (Vern - H.-Musna, B.-Sabuni) and probably some others of the family.

It is acrid and its toxicity is partially removed by boiling

Hypericaceae

(St John's wort Family)

Balsamic resinous juice

- 32 **Hypericum perforatum** Linn (Vern -H & P -Basant)

It is poisonous to animals, especially horses if taken in excess, usually it is not eaten

Guttiferae

(Gamboge Family)

Gum resins

- 33 **Calophyllum inophyllum** Linn (Vern -S -Punnaga, H. Sultana, champa, B -Punnag, Bo -Undi, M -Punnagam).

It is a fish poison.

- 34 **Garcinia morella** Desrouss, and probably others

It yields a gum resin which is a violent gastro-intestinal irritant

Ternstroemiaceae

(Tea Family)

Caffeine, theophylline

- 35 **Thea sinensis** Linn

Its excessive indulgence only is harmful.

Malvaceae

(Cotton family)

Gossypol, resin, ephedrine, pseudo-ephedrine.

36. **Gossypium** species.

The root bark is emmenagogue and used as abortifacient; occasionally harmful effects of cotton seed cakes on animals have been reported.

37. **Malva parviflora** Linn. (Vern -H. Panirak)

It is reported to have produced narcotic poisoning in animals.

38. **Sida rhombifolia** Linn (Vern - S -Atibala, H & B -Swet-berela, M -Athiballachetu)

The ripe capsules are reported to be fatal to fowls.

Linaceae

(Flax Family)

Cyanogenetic compounds, cocaine

39. **Erythroxylum coca** Lam

It is a central nervous stimulant, sensory nerve endings-paralysant, addiction to it is harmful

40. **Linum usitatissimum** Linn (Vern - S -Atasi, H & B -Tisi, Bo -Alasi, M -Alshiviral).

The young plants are known to produce deaths in animals, sometimes seed cakes also harmful

Zygophyllaceae

(Bean-caper and Guaiacum Family)

Harmine, harmaline, harmalol, peganine, essential oils, saponins, resins.

41. **Peganum harmala** Linn. (Vern.-H & Bo -Hurmali, B -Isband, M -Shimai-azha-vanai-virai)

It is an insecticide, narcotic, nauseant and emetic It is used as abortifacient It is a protoplasmic poison and paralyzes skeletal and cardiac muscles of frog

42. **Tribulus terrestris** Linn. (Vern -S -Gokshura, H -Chotagokhru, B -Gokhuri, Bo -Lahana-gokhru, M.-Nirunji)

It causes the disease goeldikkop (dikgeel) in South Africa in small stock which is characterized by oedema of head, fever and jaundice.

Rutaceae

(Rue Family)

Essential oils, rutin, skimmianine, saponins, resins, etc.

43. **Acronychia pedunculata** (Linn) Miq (-*A laurifolia* Bl)
It is a fish poison.

44. **Ruta graveolens** Linn var *angustifolia* Hk f., (Vern - S.-Somalata, H.-Sadab, B.-Ermul, B -Satap, M.-Arvada), *R. tuberculata* Forsk.

These are acro-narcotic poisons which are rubefacient. The oil and herb frequently used to produce criminal abortion.

- 45 **Skimmia laureola** Sieb & Zucc ex-Walp (Vern - Nep -Chum-lan, P -Ner).

It is reported to be poisonous to sheep and goats.

46. **Zanthoxylum alatum** Roxb (Vern - S -Tumburu, H.-Tejmal, B.-Nepali dhania) (Probably some more species)

It is a fish poison.

Simaroubaceae

(Bitter-bark Family)

Essential oils, saponins, resins, bitter substances

- 47 **Ailanthus altissima** (Mill) Swingle (—*A glandulosa* Desf)

It is nauseant and nervous system depressant Accumulation of its leaves in well water is reported to produce chronic gastritis

48. **Balanites aegyptiaca** (Linn) Delile (—*B roxburghii* Planch) (Vern -S -Ingudi, H & B -Hingan, Bo -Hinger, M.-Najunda)

It is believed to be a fish poison and a purgative

- 49 **Brucea amarissima** (Lour) Merr - (*B sumatrana* Roxb)

Its seeds produce nausea, vomiting, abdominal pain and purging

- 50 **Picrasma napalensis** Benn

It is stated to be used as larvicide in Sikkim

Meliaceae

(Neem and mahogany Family)

Bitter substances, bitter oil, saponins

51. **Azadirachta indica** A Juss

It is parasiticial and the leaves are used as insect repellent

52. **Melia azedarach** Linn

Its berries are especially poisonous to man and animals, they are narcotic and gastro-intestinal irritant

- 53 **Walsura piscidia** Roxb (Bo & M -Walsura).

It is a dangerous emmenagogue and a violent emetic. It is largely used as a fish poison.

Celastraceae

(Spindle-tree Family)

Alkaloid, essential oil, resin.

54. **Elaeodendron glaucum** Pers. (Vern -H.-Bakra, Bo.-Bhuta-pala, M -Selupa.)

It is emetic, overdoses fatal.

Sapindaceae

(Soap-nut Family)

Saponins, cyanogenetic compounds

- 55 **Cardiospermum halicacabum** Linn (Vern -S -Karnaspota, H.-Kanaphata, M -Mooda cotton).

Its leaves are emetic and rubefacient

- 56 **Dodonaea viscosa** Linn (Vern -H -Aliar, Bo -Bandurgi, M -Virali)

It is fish poison, it is deleterious to camels

- 57 **Harpullia cupanioides** Roxb.

It is a fish poison

- 58 **Melianthus major** Linn.

It produces acute diarrhoea, salivation and colic, honey from flowers stated to be poisonous.

- 59 **Sapindus mukorossi** Gaertn. (Vern -S -Phenila, H B & Bo -Ritha), *S trifolatus* Linn (Vern -S -Phenila, H B. & Bo -Ritha, M -Ponnan-kotta)

Both are fish poisons, emetic, purgative and are used for procuring abortion

- 60 **Schleichera oleosa** (Lour) Merr. (*S tryuga* Willd.) (Vern -H -Kosum, M -Pu-maram).

The oil is occasionally mixed with mustard oil or ghee It produces irritant poisoning and the seeds are used as insecticide.

Anacardiaceae

(Cashew and mango Family)

Toxic phenolic compounds, toxic resin.

- 61 **Anacardium occidentale** Linn. (Vern -H. & Bo.-Kaju, B.-Hijli badam, M -Mundiri-ka).

The pericarp contains powerfully vesicant juice, used to preserve floors, wood, books, etc. from white ants, tar from bark also vesicant.

62. **Holigarna arnotiana** Hook. f. (Vern -Bo -Bibu), *H. ferruginea* March, *H grahami* (Wight) Hook f, *H longifolia* Buch. Ham ex-Roxb. (Vern - B.-Barola, Bo -Hulugiri).

The juice is vesicant although not equally powerful in all species.

63. **Rhus insignis** Hook f., *R. punjabensis* J. L. Stewart, *R. succedanea* (Vern -S -Karkata sringi, H & B.-Kakrasingi, Bo -Taka-dasingi), *R. wallichii* Hook. f. (Vern -Nep.-Ghosi, H -Akoria.)

Dreaded by local people; even smoke from burning wood is considered poisonous, its juice is vesicant

- 64 **Semecarpus anacardium** Linn f (Vern -S-Bhallatamu, H. & B-Bhela, Bo-Biba, M-Shayrang), *S travancoricus* Bedd

The pericarp contains vesicant juice It is used in some parts as abortifacient.

Coriariaceae

(Coriaria Family)

Coriamyrtin, tutin in foreign species.

65. **Coriaria nepalensis** Wall

It is stated to be narcotic, the foreign species are very poisonous acting like picrotoxin and producing convulsions.

Moringaceae

(Horse-radish Family)

Essential oils, alkaloid, moringine, moringinine

- 66 **Moringa oleifera** Lamk (—*M pterygosperma* Gaertn) (Vern -S - Sobhanjana, H -Sajnah, B -Sojna, Bo-Sujna, M -Murunga1)

The fresh root bark is said to be vesicant and is used to procure abortion Moringinine acts on sympathetic nervous system.

Leguminosae

(Pea Family)

Alkaloids, glucosides, saponins, cyanogenetic compounds, rotenone, toxic albumin, bitter substances ; globulins

67. **Abrus precatorius** Linn (Vern -S & Bo -Gunja H -Gaungchi B.-Kunch, M -Gunduman1)

It is specially a blood poison and is used to poison cattle and to procure abortion.

- 68 **Acacia pennata** Willd (H.-Biswul, Kumaon -Agla, Nep -Arfu)
It is a fish poison

- 69 **Albizzia procera** Benth (Vern -H -Safed siris, B -Kori, Bo - Kina1 tihiri, M -Konda vaghe)
It is a fish poison

- 70 **Butea monosperma** (Lam) O Ktze (—*B frondosa* Koen ex-Roxb) (Vern - S -Kinsuk, H & B -Palas)

Its seeds are insecticide and painful if taken internally

- 71 **Caesalpinia nuga** Ait (Vern -M -Kakumullu)
It is a fish poison.

72. **Canavalia virosa** W. & A. (*C. ensiformis* DC. var *virosa* Baker. (Vern .-M -Kattuvalari).

Its fruit is stated to be poisonous.

- 73 **Cassia absus** Linn. (Vern -H -Chaksu, M -Karun Kanam), *C. acutifolia* Delile, *C. alata* Linn, *C. angustifolia* Vahl, *C. fistula* Linn, *C. obovata* Collad (Vern -Surati sonnamukai).

These are all purgatives which are irritant in large doses *C. absus* seeds are irritant application to eyes, and *C. alata* is a fish poison

- 74 **Clitoria ternatea** Linn (Vern -S -Aparajita, H. & B -Aparajit, M -Kakkanan)

The roots are powerful cathartic like Jalap, not a safe medicine

- 75 **Cytisus scoparius** Linn

The plant is not eaten by cattle It is an emetic and a cathartic

- 76 **Dalbergia stipulacea** Roxb.

It is a fish poison.

- 77 **Derris elliptica** Benth (Vern -Malay -Tubah), *D. scandens* Benth (Vern -B -Noalata, P -Gunj, M -Nala tige), *D. uliginosa* Benth. (Vern -B -Panlata, Bo -Kirtana), (possibly *D. ferruginea* Benth)

These are all fish poisons and *D. elliptica* is a powerful insecticide.

78. **Entada phaseoloides** (Linn) Merr (—*E. scandens* Benth) (Vern -H -Chian, B -Gilagach, Bo -Gardal).

It is a fish poison

- 79 **Lathyrus aphaca** Linn, *L. sativus* Linn (Vern -S -Tripuri, H. & B -Khesari, Bo -Lakh)

It is a food and fodder *L. sativus* if taken in large amounts and over prolonged period produces lathyrism in man and animals Ripe seeds of *L. aphaca* are stated to be narcotic if taken in excess

- 80 **Melilotus alba** Desr.

It is stated to be poisonous to cattle.

- 81 **Milletia auriculata** Baker, *M. pachycarpa* Benth., *M. piscidia* Wight.

These are fish poisons *M. auriculata* is an insecticide.

- 82 **Mundulea suberosa** Benth.

It is a fish poison.

- 83 **Ougenia dalbergioides** Benth. (Vern - S.-Tinisasegandun, H.-Sandan, B.-Tinis, Bo Tiwas, M-Tella-motuku).

It is a fish poison.

84. **Phaseolus lunatus** Linn.

The coloured variety sometimes exhibits poisonous properties if eaten.

85. **Pithecellobium bigeminum** Mart (Vern -H. & Bo -Kachlora)

It is a fish poison The seeds are stated to be eaten in Burma but sometimes produce severe poisoning

86 **Pongamia pinnata** (Linn) Merr (—*P. glabra* Vent) (Vern - S., H. & Bo -Karanja, B Dahar karanja, M -Pungammaram)

It is piscicide and insecticide.

87 **Sophora mollis** R. grah var. *Hydaspidis* Baker, *S. tomentosa* Linn.

The Seeds of *S. mollis* have insecticidal properties and the leaves of *S. tomentosa* are powerfully emetic and cathartic in large doses.

88 **Tephrosia candida** Linn, *T. purpurea* Pers (Vern - S -Sara-punkha, B -Bon-nil, H & Bo -Sarphankha, M -Kolluk-kay-velai)

These are fish poisons Some foreign species are insecticides Some species of *Tephrosia* in India are likely to prove of value as insecticides

89. **Trifolium repens** Linn

It is highly prized fodder in Europe In the Himalayas poisoning has been reported in horses

90 **Vicia sativa** Linn. (Vern -H -Ankra, B.-Ankari)

It is suspected to be the cause of lathyrism.

Rosaceae

(Rose family)

Cyanogenetic glucosides, phloridzin

91 **Prunus amygdalus** Batsch. (bitter variety). (Vern -H, B. & Bo -Badam, M -Vadam-Kottai), *P. armeniaca* Linn. (Vern -H -Khubani, P -Gurdli), *P. avium* Linn, *P. cerasus* Linn, *P. mahaleb* Linn (Vern -S, Priyangu), *P. padus* Linn. (Vern -H -Jamana, P -Jamma), *P. persica* Stokes, (Vern -H -Aru), *P. puddum* Roxb, (Vern -S.-Padmaka, H -Paddam, Bo -Padma-kasta), *P. undulata* Buch -Ham

The seeds are poisonous and the leaves of many species are said to be dangerous to live-stock when wilted; harmless when on the plant, suspicious when dried.

92. **Pygeum gardneri** Hook. f.

The seeds are fish poison.

93. **Pyrus aucuparia** Linn., *P. malus* Linn.

The bark of *P. aucuparia* is irritant to the alimentary tract and wilting leaves of other species are occasionally poisonous to animals browsing upon them.

94. **Rubus moluccanus** Gaertn (Vern.-Kumaon.-Katson)

The leaves are reported as powerful emmenagogue and abortifacient.

Crassulaceae

(Life-Plant-Family)

Glucosides-in foreign species.

95 **Kalanchoe spathulata** DC. (Vern:-H.-Tatara).

The expressed juice of bitter variety is a drastic purgative and is poisonous to goats. It is not eaten by cattle. Leaves are said to be insecticidal.

Droseraceae

(Sundew Family)

Organic acids, proteolytic enzymes

96 **Drosera peltata** Sm var. *lunata* Clarke, (Vern - H -Mukajali, P Chitra), *D. spathulata* Labill. (*D. burmanni* Vahl)

These are rubefacient Some Australian species are reported to be injurious to sheep.

Combretaceae

(Myrobolan Family)

Tannins

97 **Terminalia bellerica** Roxb (Vern -S -Bahira, H. & B.-Bahera, Bo -Behaira, M.- Vallaimurdu), *T chebula* Retz. (Vern - S & B -Haritaki, H -Harara, Bo - Hirda, M - Kadukkay-pu)

T bellerica is reported to be fish poison. The kernel is stated to be poisonous and cases are reported where narcotism followed nausea and vomiting Some varieties of *T chebula* are drastic purgatives.

Myrtaceae

(Myrtle and jamun Family)

Saponins, essential oils, tannins.

98 **Barringtonia acutangula** Gaertn (Vern - S. - Dhatriphal, B -Hijal, H -Hijal, Bo -Samudraphala, M -Samutrapullam), *B asiatica* Kurz (*B speciosa* Forst), *B racemosa* Bl (Vern - S -Samudrapad, H -Norvishee, B -Samudraphal, M -Samudra).

These are believed fish poisons

99 **Careya arborea** Roxb. (Vern -S , H & B.-Kumbhi).

It is a fish poison The inner bark rubbed on shoes keeps off leeches.

100 **Eucalyptus globulus** Labill (Vern - M -Karpura maram)

The essential oil contained is an important ingredient of insecticides, internally gastro-intestinal irritant.

- 101 **Melaleuca leucadendron** Linn. (Vern - H., B & Bo.-Nim, M - Vembu)

Its essential oil is an irritant and a mosquito repellent.

Lythraceae

(Henna and pomegranate Family)

Acrid principle

- 102 **Ammannia baccifera** Linn (Vern - S - Agnigarva, H - Jangli-mendi, M - Nirumel-neruppu) *A. senegalensis* Lamk (Vern - P.-Fauglimehandi)

These are vesicant, taken internally they cause great pain.

- 103 **Lagerstroemia indica** Linn, *L. speciosa* (Linn) Pers (-*L. flosreginae* Retz) (Vern - S - Arjuna, H & B - Jarul, B - Taman, M Kodali)

The bark and leaves are purgative, seeds of the former species are narcotic

Samydaceae

(Casearia Family)

- 104 **Casearia graveolens** Dalz (Vern - H - Chilli, Bo - Naro), *C. tomentosa* Roxb (Vern - H - Chillara)

The pounded fruit is used as a fish poison

Caricaceae

(Papaw Family)

Carpaine, carposide, caricin in seeds yielding essential oil on hydrolysis, papain

- 105 **Carica papaya** Linn (Vern - H - papaya, B - Papey, Bo.-Papai, M.- Pappayi)

The seeds are believed to have powerful emmenagogue properties and are used as abortifacient The juice of unripe fruit is acrid or even vesicant

Passifloraceae

(Passion-flower Family)

Hydrocyanic acid, saponins

106. **Adenia (modecca) palmata** Engl., *A. wrightiana* Engl.

The roots and fruits are poisonous. Deaths from eating fruits of *A. palmata* are reported

Cucurbitaceae

(Cucumber Family)

Bitter substances, such as colocynthin, alkaloids, glucosides.

107. **Citrullus colocynthis** Schrad, (Vern - S - Indra varuni, H. & Bo.-Indrayan, B.-Makhal, M -Peyt-tumatti) *C vulgaris* Schrad. (Bitter variety).

The fruit is purgative, *C colocynthis* is a drastic purgative and has produced fatal results, the dust when dry is very irritating to eyes and nostrils.

108. **Corallocarpus epigaeus** Benth. & Hook f.

Its fruit is a drastic purgative

109. **Cucumis sativus** Linn (Bitter variety) (Vern -S -Sukasa, H - Khira, B -Sasa, Bo -Kankri, M -Mulluvellari), *C trigonus* Roxb (Vern -S -Vishala, H -Bislambhi, M -Hattut-tumatti)

The fruit is a purgative, *C trigonus* excessively so

- 110 **Lagenaria siceraria** Standl (*L vulgaris* Seringe) (Wild variety)

It is drastic purgative Cases have been reported where beer kept in bottle gourd produced poisoning

- 111 **Luffa acutangula** Roxb. var *amara* C B Clarke (Vern -S -Koshataki, H & Bo -Torai, B -Jhinge, M -Pikumkai), *L aegyptiaca* Mill. ex. Hook f (Vern -S -Rajkoshataki, H -Ghiatarui, B -Dhundul, Bo -Ghosali, M -Guttibira) (Wild variety), *L echinata* Roxb (Vern -S -Koshataki, H -Kukarlata, B -Ghosalata, Bo -Kukarwele, M -Panibira).

The fruit of *L. acutangula* var *amara* is violently emetic and purgative and is not eaten. Others are also purgative

- 112 **Momordica balsamina** Linn (Vern -Bo -Kurelo-jangro), *M charantia* Linn (Vern -S -Sushavi H -Karela, B -Karala, Bo -Karla, M -Pavakkachedi), *M tuberosa* Cogn (—*M cymbalaria* Fenzl) (Vern -Bo -Kadavanchi)

Fruit of *M balsamina* is fatal to dogs, death occurs from violent vomiting and purging produced by the juice of this plant. *M charantia*, roots are used as abortifacient Decoction of roots of *M. tuberosa* are used as abortifacient

113. **Trichosanthes bracteata** Voigt (—*T palmata* Roxb) (Vern -S -Mahakal, H -Lal-indrayan, B.-Makal, Bo -Kaundal, M -Korattai) *T. cucumerina* Linn. (H -Jangli-chichonda, S -Patola, B -Bonpatol, Bo -Ranparul, M.-Pudel), *T dioica* Roxb (Vern -S -Patola, H.-Parvar, B.-Potal, Bo -Potala, M -Kombupudalai)

The roots are powerful cathartic Fruit of *T cucumerina* is never eaten, because of powerful cathartic action Fruit of *T. bracteata* is used as cattle poison and to destroy crows

- 114 **Zanonia indica** Linn. (Vern -S.-Dirghapattra, H -Chirpoti, Bo.-Chiraputi, M.-Penarvalli).

The fruit is very acrid and a powerful cathartic

123. **Sambucus ebulus** Linn , (Vern -P —Mushkiara), *S. nigra* Linn

Both are strongly purgative *S. ebulus* has foetid smell when bruised and is not eaten by cattle, poisoning amongst boys and fowls reported

Rubiaceae

(Madder and coffee Family)

Quinine, quinidine, cinchonine, cinchonidine, caffeine, emetine, cephaeline, ipecacuanhin, essential oils, saponins.

- 124 **Adina cordifolia** Benth & Hook f (Vern —S.-Dharakadamba, H -Hardu, B -Keli kadam, M -Manja kadambe)

Its juice is used as insecticide

- 125 **Cinchona calisaya** Wedd and var. *ledgeriana* Howard, *C. officinalis* Linn f , *C. succirubra* Pavon

The cinchona alkaloids are general protoplasmic poisons and parasiticide The plants are believed fish poisons.

- 126 **Coffea arabica** Linn (Vern —H -Coffee, B.-Kafi)

Its excessive indulgence produces harmful effects and chronic poisoning

- 127 **Cephaelis ipecacuanha** Stokes

It is emetic and irritant and a powerful cardiac depressant

128. **Randia dumetorum** Lamk , (Vern —S -Madan, H -Mainphal, B -Menphal, Bo -Gelaphal, M -Maruk-kallan-kai), *R. uliginosa* DC (Vern —S -Pindaluka, H -Pindalu, B -Piralu, Bo -Pendari, M - Wagata)

These are believed fish poisons *R. dumetorum* is used to preserve grain from attacks of insects, it is also used as abortifacient.

Compositae

(Sun-flower Family)

Essential oils, artemisin, santonin, bitter substances (absinthin lactucin, etc), saponins, resin, senecio alkaloids, xanthostrumarin, pyrethrins.

129. **Anthemis cotula** Linn.

It is an undesirable food for livestock, it is acrid and vesicant.

130. **Artemisia absinthium** Linn. (Vern —H. & Dec -Vilayatı afsantin), *A. maritima* Linn (Vern —S -Gadadhar, H -Kirmala, Bo -Kiramani owa), *A. vulgaris* Linn (S -Nagadamani, H -Nagadouna, B -Nagdona).

The essential oil from *A. absinthium* is a violent narcotic poison producing convulsions *A. maritima* is an irritant poison when given in large doses, and fatal cases are reported *A. vulgaris* produces epileptiform spasms in man and is also reported as fish poison.

131. **Centratherum anthelminticum** O. Ktze. (- *Vernonia anthelmintica* Willd).

It is used as insecticide and insect repellent.

132. **Chrysanthemum cinerariæfolium** Vis., *C. coccineum* Willd., *C. roseum* Adam.

All are reputed insecticides

133. **Erigeron canadensis** Linn.

It is an irritant.

- 134 **Eupatorium odoratum** Linn.

It is stated to be a fish poison. *E urticifolium* L. f. of foreign countries produces acidosis and trembles in sheep and cattle

135. **Gnaphalium luteo-album** Linn (Vern -P -Balraksha).

It is suspected to produce livestock-poisoning in South Africa.

136. **Inula graveolens** Desf.

It is suspected to be poisonous to livestock.

137. **Lactuca tatarica** C A Meyer var *tibetica* C B Clarke

It is occasionally browsed by sheep, sometimes injurious

138. **Saussurea lappa** C.B Clarke (Vern —S.-Kushtha, H -Kut, B.-Pachak, Bo.-Ouplate, M.-Goshtam).

The roots are used to protect woollen fabrics against insects.

139. **Senecio species** (*S vulgaris* Linn introduced plant)

The Indian species should be studied in Ragwort poisoning due to several species of this plant is well known in foreign countries, some species produce hepatic cirrhosis

140. **Sphaeranthus indicus** Linn (Vern —S.-Munditika, H & Bo -Gorakmundi, B.-Murmura, M -Kottak).

It is believed to be a fish poison.

141. **Xanthium strumarium** Linn. (Vern —S.-Arishta, H.-Chhota-gokru, B -Bon-okra, Bo.-Shankesh vara, M -Marlu-mutta)

It is reported to be poisonous to cattle and pigs in America and Australia.

Campanulaceae

(Bell-flower Family)

Alkaloids.

142. **Lobelia excelsa** Leschen, *L. nicotianifolia* Heyne. (Vern.—Bo.-Dhavala, M.-Rattu papillay).

It is irritant to nose and deaths have been reported in man. Its action resembles nicotine, except that more burning pain in the stomach is produced. It is used as substitute for datura.

Ericaceae

(Rhododendron Family)

Andromedotoxin, ericolin, Essential oils

143. Gaultheria fragrantissima Wall.

It is an irritant poison and deaths are reported from its use as abortifacient

144 Pieris ovalifolia D. Don.

It is poisonous to goats and is an insecticide.

145 Rhododendron anthopogon D Don , (Vern —Kash.-Tazak-tsun), *R. arboreum* Sm (Vern -P —Ardawal), *R. barbatum* Wall (Vern —Nep -Guras), *R. campanulatum* D Don (Vern -H -Cheraelu Kash -Gaggar), *R. cinnabarinum* Hook f (Vern. --Nep -Bulu), *R. falconeri* Hook f (Vern —Nep -Kurlinga), *R. setosum* D. Don (Vern - Bhutia -Tsallu).

These are probably all poisonous to live-stock; some are reported to be fish poisons and honey from some is reported to be poisonous.

Plumbaginaceae

(Plumbago Family)

Plumbagin

146 Plumbago indica Linn (-*P. zeylanica* Linn) (Vern:-S —Chitraka H & B -Chita, Bo -Chitaro, M -Chittira), *P. rosea* Linn. (Vern —S -Chitraka, H., B. & Bo -Lal chitra, M.-Chittur-mol).

These are strong irritants externally and internally and are used to procure abortion.

Primulaceae

(Prime-rose Family)

Saponins

147. Anagallis arvensis Linn. (Vern —H.-Jonkhmar)

It produces gastro-enteritis in dogs and horses. It is used to poison fish and expel leeches from nostrils of animals.

148 Cyclamen persicum Miller (Vern --Ind. Baz.-Bankhur-1-Miryam).

It is believed to be a fish poison.

149. Primula reticulata Wall. (Vern -Kumaon.—Bishcopra).

It is stated to be poisonous to cattle.

Myrsinaceae

(Ardisia Family)

Saponins.

- 150 **Maesa indica** Wall. (Vern — M -Kırithı) Leaves are stated to be fish poison.

Sapotaceae

(Sapodilla and mohwa Family)

Saponins

- 151 **Madhuca indica** J F Gmel *M* (*Bassia*) *latifolia* (Roxb.) Macbride, *M. longifolia* (Linn) Macbride.

The residual cake is used as fish poison and is said to have insecticidal properties *Mohwa meal* is used to kill worms on lawns

Ebenaceae

(Ebony Family)

Bitter substances

- 152 **Diospyros ebenum** Koenig (Vern — H & Bo -Tendu, M -Acha), *D. montana* Roxb (Vern -S --Tumala, H -Lohari, B -Ban-gal, Bo Kundu, M -Muchi-tankı), *D paniculata* Dalz (Vern —S -Thinduka, M -Karınthuvarı)

All these are believed to be fish poisons.

Salvadoraceae

(Salvadora Family)

- 153 **Salvadora oleoides** Dcne (Vern —H , S. & B -Pilu, Bo.-Kankhina, M -Ughaiputtaı), *S persica* Linn (Vern —S -Pilu, H & B -Chota-pilu, Bo -Pilvu, M -Ughaiputtaı)

The root bark is vesicant

Apocynaceae

(Dog-bane and Oleander Family)

Kurchı and rauwolfia alkaloids, glucosides, e g cerberin, karabin, nerin, neriodorin, oleandrin, I-strophanthin, thevetin, etc., bitter substances

- 154 **Allamanda cathartica** Linn. (Vern —Bo -Jaharı sontakka)
It is a hydragogue cathartic
155. **Cerbera manghas** Linn (-*C odollam* Gaertn) (Vern.—B.-Dhakur, M -Kataralı)

The green fruit is used to poison dogs. The seeds are irritant poison and the plant is a fish poison

- 156 **Ervatamia dichotoma** (Roxb) Blatter (-*Tabernaemontana dichotoma* Roxb)

The seeds are powerfully narcotic and poisonous.

- 157 **Holarrhena antidysenterica** Wall (Vern — S -Kutaja, H Karchı, B -Kurchı, Bo -Pandhrakura, M -Kashappu-vetpalarışı)

It is not browsed by cattle and goats, it is an anthelmintic, kurchicine is a general protoplasmic poison.

158. **Lochnera pusilla** K Schum (-*Vinca pusilla* Murr.), *L rosea* (Linn) Reichb (-*Vinca rosea* Linn)

These are cardiac poisons and *L pusilla* is regarded as poisonous to cattle.

- 159 **Melodinus monogynous** Roxb (Vern —B -Sadul kou)
It is a fish poison

- 160 **Nerium-indicum** Mill. (*N odorum* Soland) (Vern —S -Karavi, H -Karber, B -Karabi, Bo -Kanhera, M Alari)

It is very poisonous and is used for suicidal purposes and to procure abortion, depresses nervous system and heart

- 161 **Plumeria acuminata** Ait (-*P acutifolia* Poir) (Vern —S -Kshira champa, H & Bo -Khair Champa, B -Gobar champa, M -Vadaganneru)

Its milk is rubefacient and is used to procure abortion, internally it acts as purgative and is poisonous

- 162 **Rauwolfia serpentina** Benth ex Kurz (Vern —S -Sarpagandha, H -Chota chand, B & Bo -Chandra, M -Covannamilpori)

It is hypnotic and a fish poison

- 163 **Thevetia peruviana** (Pers) Merr (-*T nerifolia* Juss) (Vern —H & Bo-Pila-kaner, B -Kolkaphul, M -Pachch-a1-alar1)

All parts especially seeds are very poisonous It is used to poison cattle and produces violent vomiting and purging
Action on heart like digitalis It is also a fish poison

Asclepiadaceae

(Milk-weed Family)

Tylopharine, glucosides, bitter substances, resins, saponins, etc

- 164 **Asclepias curassavica** Linn (Vern —H -Kakatundi, Bo-Karki)

It is a fish poison, emetic and cathartic

- 165 **Calotropis gigantea** R Br (Vern —S -Arka, H -Ak, B & Bo -Akanda, M -Erukku), *C procera* R Br (Vern -S Alarka, H -Madar, P -Shakar-al-lighal, Bo -Mandara, M -Vellerku)

Milk of these plants is drastic purgative and caustic Stated to be used for suicidal and homicidal purposes and as an abortifacient and cattle poison

- 166 **Cryptostegia grandiflora** R Br (Vern —Bo -Vilayati-vakhandi, M.-Palai).

A fatal case from leaves is reported in which persistent vomiting was observed

167. **Cynanchum arnottianum** Wight, *C. vincetoxicum* Pers.

C. arnottianum is used as insecticide. *C. vincetoxicum* is not eaten by cattle and is regarded poisonous Its root is emetic.

- 168 **Sarcostemma acidum** (Roxb.) Voigt (-*S. brevistigma* W. & A.)
(Vern —S & Bo -Soma, H. & B -Somlata, M.-Kondapala).

Stated to have insecticidal properties.

- 169 **Secamone emetica** R. Br. (Vern —B -Shada-buri).

The roots are acrid and the plant is powerfully emetic.

- 170 **Tylophora indica** (Burm f) Merr (-*T. asthmatica* Wight and Arn), (Vern -H. & B -Antamul, Bo -Anthamul, M.-Nay-palai),
T. fasciculata Buch.-Ham (Vern -Bo -Bhuidari)

Fatal cases are reported in man, it is emetic and *T. fasciculata* is used as rat poison.

Loganiaceae

(Nux-vomica Family)

Strychnine, brucine, etc.

- 171 **Strychnos colubrina** Linn (Vern —H & B -Kuchila-lata, Bo -Goagarilakei, M -Nagamusadi), *S nuxvomica* Linn. (Vern —S -Visha-mushti, H -Kuchla, B -Kuchila, Bo.-Kajra, M -Yetti)

These are poisonous *S nux-vomica* seeds are used as fish poison and are source of strychnine, one of the deadliest poisons known Suicidal and homicidal cases are recorded. It is employed to kill dogs, rodent, etc

Boraginaceae

(Borage and Sebestan Family)

Alkaloids.

- 172 **Heliotropium eichwaldii** Steud (Vern —H & P -Nilkattei, Kash.-Chirghas), *H indicum* Linn (Vern —S -Hastisunda, H. & B -Hatisura, Bo -Burundi, M -Tel-kodukk1)

These are suspected to be poisonous

Convolvulaceae

(Convolvulin, pharbitin, terpithin, terpethein, cucutalin, resin.

- 173 **Calonyction muricatum** (Linn.) G Don (-*Ipomoea muricata* Jacq)

See Ipomoea.

- 174 **Convolvulus arvensis** Linn (Vern —H -Hiranpadi, Bo -Hiranpag, M -Naranji), *C scammonia* Linn.

The roots are strongly purgative

175. **Cuscuta reflexa** Roxb. (Vern —S Amaravela, H.-Akasbel, B -Algusi, Bo.-Nirmuli, M -Sitamapurgonalu).

It is nauseant and emetic, it is used to procure abortion.

- 176 **Ipomoea reptans** (Linn) Poir (-*I aquatica* Forsk) (Vern —S -Kalambi, B -Kalmisak, Bo.-Nalichi baji, M -Sarkarei-valli),
I. nil Roth (—*I. hederacea* Jacq.) (Vern —H., B. & Bo -Kaladana, M -Jirkivirai), *I purga* Heyne,

These are strongly purgative and irritant poisons in over-doses.

177. **Operculina turpethum** (Linn) Manso (—*Ipomoea turpethum* R. Br.)

See Ipomoea

Solanaceae—

(Datura and nightshade Family)

Hyoscyamine, hyoscine, atropine

- 178 **Atropa belladonna** Linn. (Vern —H -Sag-angur, B -Yebruj, Bo.-Girbuti)

Fatal cases of poisoning are reported, dryness of mouth and throat, dilation of pupils and delirium characteristic features

- 179 **Capsicum annum** Linn. (Vern —H. & P -Mirch), *C frutescens* Linn., *C. minimum* Roxb.

The seeds are gastro-intestinal irritant and are used for torturing purposes

180. **Datura fastuosa** Linn. (Vern —S -Krishna dhatura, H & B -Kala Dhatura, M -Karu umattai), *D metel* Linn , *D. stramonium* Linn (Vern —B.-Sada dhutura, P -Tattu dattura, M -Umatai).

These are commonly used by criminals for stupefying their victims , symptoms resemble those of Atropa

181. **Hyoscyamus muticus** Linn , *H. niger* Linn (Vern —S -Parasi-kaya, H -Khurasani ajvayan, B.-Khurasani ajowan, Bo -Khurasani-owa, M -Khurasani-yomam), *H. pusillus* Linn , *H. reticulatus* Linn.

Cases of livestock and children poisoning are on record, action like Atropa.

- 182 **Lycium barbarum** Linn. (Vern —Baluchi -Koh-tor)

It is reported to be poisonous to livestock

- 183 **Mandragora caulescens** Clarke

It is suspected to be poisonous

- 184 **Nicandra physaloides** Gærtn.

It is considered to be insecticide

- 185 **Nicotiana rustica** Linn. (Vern —H & B -Vilayeti tamaku, P -Kakkar tamaku), *N. tabacum* Linn (Vern —H -Tamaku, B -Tamak., Bo -Tambaku, M.-Pugai-ilai)

These are insecticide and are also used to ward off leeches fatal cases reported among human beings and livestock.

186. **Physochlaina praealta** Miers (Vern —P -Nandru)

It is reported to be poisonous.

187. **Scopolia anomala** (Linn et Otto) Airy-Shaw. (*S. lurida* Dunal).

It is poisonous and action is like Atropa.

- 188 **Solanum dulcamara** Linn (Vern — S Kakmachı, P -Ruba-barık)
S. incanum Linn , (-*S coagulans* Forsk), *S. nigrum* Linn (un-ripe berries) (Vern — S & B -Kakmachı, H -Makoı, Bo -Mako, M -Manattak kalı), *S spirale* Roxb (Vern — H -Mungas kajur, Bagua), *S. tuberosum* Linn (Sprouting).

The cases of poisoning among human beings and animals are reported, some fatal These are gastro-intestinal irritant , occasionally associated with atropa-like symptoms

- 189 **Withania somnifera** Dunal (Vern — S , B & Bo -Ashwagandha, H -Asgandh, M -Amku-lang-kalang)

It is reported to be used as abortifacient and as an insecticide, it is also stated to be hypnotic

Scrophulariaceae

(Mimulus and Digitalis Family)

Digitalin, digitonin, digitoxin, gitalin, gitonin, etc , saponin, bitter substance

- 190 **Digitalis purpurea** Linn

It is a cardiac poison and fatal cases due to eating of plant are reported in India

- 191 **Verbascum thapsus** Linn

It is a fish poison and the seeds are narcotic

Bignoniaceae

(Bignonia Family)

- 192 **Dolichandrone falcata** Seem. (Vern.-H — Hawar, Bo -Manchingı, M -Kadatathıe)

It is fish poison reputed to be abortifacient.

Pedaliaceae

(Sesamum Family)

Sesamol (a phenolic substance), seasmolin

- 193 **Sesamum indicum** Linn (-*S orrientalis* Linn) (Vern.—S , H , B , & Bo -Tıl, M -Yellucheddie)

Seed cakes are commonly fed to cattle in India but it is stated to be toxic to livestock in Europe producing colic, tremors, dyspnoea and distention.

Verbenaceae

(Verbena and teak Family)

194. **Callicarpa longifolia** Lamk var *lanceolaria* C. B. Clarke.

It is a fish poison

- 195 **Duranta repens** Linn (*D plumieri* Jacq)

It is very bitter and believed to be poisonous to livestock, but generally refused

196. **Lantana aculeata** Linn (-*L camara* Linn) (Vern -Bo.—Vhaneri, M -Aripu)

The reports about being poisonous to livestock received from the Punjab and Assam Government Departments described it to be abortifacient.

- 196A **Stachytarpheta jamaicensis** (Linn) Vahl var. *indica* Lam (-*S indica* Vahl.)

It is described as abortifacient

- 197 **Verbena officinalis** Linn (Vern.—P -Pamukh)

It is stated to be irritant poison

Labiataeae

(Mint and sage Family)

Essential oils, saponins

198. **Eremostachys acanthocalyx** Boiss, *E vicaryi* Benth (Vern —P.-Gurgunna)

E acanthocalyx is stated to be poisonous and *E vicaryi* is used as a fish poison

- 199 **Lamium amplexicaule** Linn

It is regarded as injurious in America

- 200 **Pogostemon heyneanus** Benth (*P patchouli* F B I, non Pelletier) (Vern --H -Pacholi)

The leaves are used as insecticide.

Chenopodiaceae

(Spinach and beet Family)

Essential oils, saponins, salsoline, oxalic acid

- 201 **Chenopodium ambrosioides** Linn, *C botrys* Linn

Anthelmintic against hook-worm and round worm Fatal poisoning is on record

- 202 **Haloxylon recurvum** Benge ex Boiss, *H salicornicum* Bunge ex Boiss.

These are stated to be poisonous but *H. recurvum* is a favourite food of camels

- 203 **Salicornid brachiata** Roxb (Oomarie keeray) Ash is stated to be abortifacient

204. **Salsola kali** Linn.

It is suspected to be poisonous but a feeding test with half dried plants in flowering stage was negative

- 205 **Suaeda fruticosa** Forsk

It is stated to be poisonous.

Phytolaccaceae

(Phytolacca Family)

Bitter substances.

- 206 **Phytolacca latbenia** (Buch-Ham) H Walt. (*P. acinosa* Hook f., B I non-Roxb.) (Vern —H.-Matazor)

It is stated to be poisonous if eaten raw, but it is edible when cooked.

Polygonaceae

(Buck-wheat and rhubarb Family)

Rutin, essential oils, anthra-quinone derivatives, oxalic acid, oxalates.

207. **Fagopyrum esculentum** Moench, *F. tataricum* Gaertn.

It is commonly eaten but under certain conditions, not properly understood at present, produces eruptions and urticaria.

- 208 **Polygonum aviculare** Linn. (Vern —S -Nisomali, H -Bannatia, B -Machutie), *P flaccidum* Meissn, *P hydropiper* Linn. (B -Packur-mul), *P orientale* Linn., *P persicaria* Linn., *P. tomentosum* Willd

P hydropiper is biting to a degree that no animal will eat it. Acrid, emetic, vesicant, insecticidal and piscicidal properties to varying degree are strongly suspected.

- 209 **Rheum emodi** Wall (Vern —H. & B -Revandchini, Bo.-Ladaki-revanda chini, M -Nattu-ireval-chinni) and probably some other species

Its leaves and petioles are edible but the latter is responsible for occasional poisoning

- 210 **Rumex acetosa** Linn, *R acetosella* Linn (Vern —S. Chutrika, B -Chukapalam)

Oxalic acid poisoning is produced if eaten in excess.

Aristolochiaceae

(Birth-wort Family)

Aristolochin, glucoside, essential oils, bitter substance.

211. **Aristolochia bracteata** Retz. (Vern —S.-Dhumrapatra, H.-Kirmar, M -Adutina-palai), *A. indica* Linn (Vern —S -Rudra-jata, H. & B -Isharmul)

These are nauseous and bitter, emmenagogue and abortifacient. *A bracteata* is insecticide

Piperaceae

(Pepper Family)

Essential oils, piperine, piperovatine

212. **Piper sp.**

Harmful effects of *P. betle* Linn. and *P. nigrum* Linn. are well known.

Myristicaceae

(Nutmeg Family)

Essential oil (with myristicin), saponins.

- 213 **Myristica fragrans** Houtt., *M. malabarica* Lamk., (Vern —Bo.-Ramphal), possibly some others also.

These are narcotic and occasional cases of poisoning are reported

Lauraceae

(Laurel Family)

Essential oils

214. **Cassytha filiformis** Linn

It is stated to be used as insecticide

- 215 **Cinnamomum camphora** F Nees (imported).

It is protective against moths, counterirritant, systemically stimulates then depresses and paralyses central nervous system.

Thymeliaceae

(Mezereum Family)

Saponins

- 216 **Daphne cannabina** Wall , *D. oleoides* Schreb

These are severe gastro-intestinal irritants. Camels do not eat *D. oleoides*

217. **Edgeworthia gardneri** Meissn.

It is a fish poison

218. **Lasiosiphon eriocephalus** Dcne

Dust from dried plant very irritant and is not eaten by livestock It is a fish poison

219. **Wikstroemia viridiflora** Meissn. (*W. indica* C. A Mey, var. *viridiflora* Hook. f)

It is a fish poison.

Loranthaceae

(Mistletoe Family)

220. **Viscum album** Linn and possibly others.

The poisonous properties are probably acquired if growing on poisonous hosts, e g. *Strychnos nux-vomica*.

Euphorbiaceae

(Croton and eastor oil Family)

Cyanogenetic compounds, saponins, crotonoside, ricinine, essential oils, euphorbon, phenolic substance, resins, toxalbumins

221. **Andrachne cordifolia** Muell. Arg. (Vern.—P.Gurguli).

It is reported to be cattle poison and African species are used as insecticide.

- 222 **Baliospermum montanum** Muell, Arg (-*B axillare* Blume)
(Vern —S, H, & B -Danti, Bo -Dantimul, M.-Naga-danti).

Its seeds and oil are drastic purgative and the seeds in over-doses are acro-narcotic poison

- 223 **Buxus sempervirens** Linn. (Vern —Kash -Chikri, P -Papri)

It is stated to be fatal to camels, cattle and goats are probably immune

- 224 **Chrozophora rottleri** A Juss ex Spreng (-*C tinctoria* Hook f in part) (Vern —H -Subali, P -Kukronda)

It is emetic and cathartic and animals avoid it

- 225 **Cleistanthus collinus** Benth. & Hook f (Vern —M -Nachuta)

It is used as fish poison and occasionally as human poison
An extract of it is a violent gastro intestinal irritant.

- 226 **Croton oblongifolius** Roxb, *C tiglium* Linn (Vern —S -Kanakaphala, H -Jamalgota, B-Joypal, Bo.-Geyapal, M -Nervalam)

The seeds especially and the oil also is a drastic purgative and is reported to be poisonous The seeds are stated to be used as insectide and piscicide .

- 227 **Euphorbia acaulis** Roxb, *E antiquorum* Linn (Vern —S -Vajrakantaka, H -Tridhara-sehund, B -Tekata si, Bo -Naraseja, M -Shadhurakkali), *E cattimandoo* W Elliot, *E helioscopia* Linn, *E hirta* Linn, *E hypericifolia*, Linn (Vern —Bo -Nayeti, P -Hazardana), *E nerifolia* Linn (Vern —S -Snuhi, H -Sehund, B -Mansa-si, Bo.-Minguta, M -Ilaikal, *E. nivulea* Buch Ham, (S patta karie, B -Si, Bo -Newrang, M -Aku-jemudu), *E peplus* Linn *E pilosa* Linn., *E rothiana* Spreng, *E royleana* Boiss (Vern —H. & P -Shakar pitan), *E thomsoniana* Boiss, *E thymifolia* Linn (Vern —S -Racta-vinda-chada, H -Chhoti dudhi, B -Dudiya, Bo -Nayeti, M -Sittrapaladi), *E tirucalli* Linn (Vern:—H.-Sehud, B -Lankasi, Bo -Shera, M -Kombu-Kali), *E trigona* Haw.

Acrid and vesicant juice is found in most species and some are used as abortifacient when applied locally *E antiquorum*, *E. nerifolia*, *E royleana*, *E tirucalli* are fish poisons and *E. antiquorum*, and *E thymifolia* are stated to be used as insecticides, some are poisonous to livestock

228. **Excoecaria agallocha** Linn (Vern —B -Gangwa, B -Geva, M -Chilla)

Its fresh sap is extremely acrid and causes intolerable pain if it gets into the eye The woodcutters have suffered and call it blinding tree It is also fish poison

- 229 **Fluggea leucopyrus** Willd, *F virosa* Baill (*F microcarpa* Bl)
(Vern —H.-Dalme, B Pandharphali)

It is a fish poison and is used to destroy worms in sores.

230. **Hura crepitans** Linn.

Its seeds and oil are violent purgative, milky juice very irritant

231 **Jatropha curcas** Linn, *J glandulifera* Roxb, *J gossypifolia* Linn, *J multifida* Linn

These are violent purgative like *Croton* sp and *J curcas* is a fish poison

232 **Manihot esculenta** Crantz (*M utilisima* Pohl.) (Vern —Baz - Cassarva M -Maravuli)

Its fresh tubers are extremely poisonous and cassava or tapioca meal is specially prepared

233 **Phyllanthus urinaria** Linn (Vern —S -Tamara valli, H & B - Hazarmani, M -Shiyappunelli)

It is stated to be a fish poison

234 **Ricinus communis** Linn (Vern —S -Eranda, H -Arand, B - Verenda, Bo -Erendi, M -Amanakham chedi)

The seeds produce violent gastro-enteritis, subcutaneously very poisonous The oil is stated to be an active poison for flies

Plant is a fish poison

235 **Sapium indicum** Willd (Vern —B -Hurua, Bo -Hurna), *S -insigne* Trinen (Vern —H -Khinna, Bo -Dudla)

S indicum juice is narcotic poison The fruit is extremely nauseous and seeds are fish poison *S insigne* juice is vesicant

236 **Tragia bicolor** Miq, *T involucrata* Linn (with varieties) (Vern —S -Vrischikali, H -Barhanta, B -Bichuti, Bo -Kanchkuri, M -Kanchuri-vayr)

These are stinging nettles

Urticaceae

(Nettle, hemp and mulberry Family)

α - β & γ -antiarin, saponin, resin containing cannabindol (toxic), formic acid

237 **Antiaris toxicaria** Lesch O (Vern —Bo -Chandla, M.- Nettavil, Sing -Riti, Burma -Hmyaseik)

Its sap is used as an arrow poison, it is a powerful heart poison

238 **Cannabis sativa** Linn. (Vern —S -Ganjika, H, B & Bo-Ganja, M -Bhangı)

The preparations of bhang, charas and ganja are well known in India, excessive indulgence produces physical and mental injury. Stated to be used as a fish poison in Bengal It is spread on beds to drive away bugs.

239. **Ficus sp.**

Some species contain acrid juice, according to Watt fruit of *F bengalensis* is poisonous to horses.

240. **Fleurya interrypta** Gaud.

Stings.

241. **Girardinia leschenaultiana** Dcne , *G zeylanica* Dcne.

These are stinging nettles.

242. **Laportea crenulata** Gaud., *L terminalis* Wight

These are stinging nettles

243 **Urtica dioica** Linn , (Vern —H & P -Bichu), *U. hyperborea* Jacq.,
U parviflora Roxb., *U. pilulifera* Linn.

These are stinging nettles.

Juglandaceae

(Walnut Family)

244. **Juglans regia** Linn (Vern —S -Akshota, H. & B -Akhroot, Bo -Akroda, M -Akrottu).

The rind of unripe fruit is stated to be fish poison in Jaunsar and Tehri Garhwal

Myricaceae

(Sweet-gale Family)

Essential oils, myricelin).

245. **Myrica nagi** Thunb. (Vern—S.-Katphala, H., B & Bo -Kaiphai, M.-Marudam-pattai).

Bark is stated to be used as fish poison in Khasia hills.

Gnetaceae

(Gnetum Family)

Saponins, bitter substance

246 **Gnetum scandens** Roxb.

It is a fish poison

Coniferae

(Pine Family)

Essential oils, taxine, taxicatin.

247 Several members especially *Taxus baccata* Linn.

Most members possess toxic essential oils and poisoning due to the use of Juniper oil as abortifacient is reported. Deaths in man and animals due to eating the berries and leaves of *T. baccata* are reported and the seeds are very poisonous.

Iridaceae

(Iris Family).

Saponins, picrocin (bitter substance), essential oils.

248. **Crocus sativus** Linn. (Vern.—S -Kumkuma, H., B & Bo -Jafran, M -Kungumapu)

Bulbs are toxic to young animals and stigmas in overdoses are narcotic poison. It is used as abortifacient

Amaryllidaceae

(Amaryllis and Agave Family)

Saponin, Lycorine, tazetidine.

- 249 **Agave americana** Linn (Vern —H -Kantala, B -Jungli anarash)

It is stated as fish poison and it is also stated to be toxic to livestock under field conditions. The wall paper impregnated with expressed juice is said to be proof against white-ants

- 250 **Cirium asiaticum** Linn Vern —S -Vishamandala, H Pindar B -Bara-kanur, Bo -Nagdowan, M -Vishomangil), *C. lotifolium* Linn (Vern —H & B Sukh-darsan)

- 251 **Narcissus tazetta** Linn (Vern —P Nargis)

Its bulbous roots are emetic and purgative, and produce irritant poisoning in overdoses

Taccaceae

252. **Tacca pinnatifida** Forst (Vern —Bo -Diva, M -Karachunai)

The tuber is intensely bitter, acrid and poisonous when fresh, yields nutritious starch by maceration and repeated washing.

Bromeliaceae

(Pine-apple Family)

- 253 **Ananas comosus** Merr (—*A sativus* Schult) (Vern —H.-Anannas, B.-Anaras, M -Anasha pazham).

The juice of leaves and unripe fruit is purgative and is sometimes used as abortifacient.

Dioscoreaceae

(Yam Family)

Dioscorine, glucoside (toxic).

- 254 **Dioscorea bulbifera** Linn. *D hispida* Dennst, (—*D. daemona* Roxb), *D prazeri* Prain & Burk (—*D deltoidea* Wall.).

Tubers are very acrid but in most cases boiling makes them edible

Liliaceae

(Lily Family)

Imperialine, colchicine, methyl-colchicine, saponine, borbaloin, emodin, sicaloin, resin, essential oils, etc.

255. **Allium sativum** Linn (Vern —S -Lasuna, H. & Bo. Lasan, B.-Rasun, M.-Vallaipundu)

Its essential oil is very irritant and pungent, produces irritant poisoning in excess. It is also stimulant, narcotic and anthelmintic.

256 **Aloe species.**

Insipisated juice 'Mushabbar' of commerce is a powerful drastic purgative and fatal cases are reported from its use. It has been used to procure abortion

257. **Colchicum luteum** Baker (Vern —Surinjan)

Resembles closely the foreign *C autumnale* which is poisonous and produces gastro-intestinal irritation The Indian species is also poisonous

258 **Fritillaria imperialis** Linn.

The bulbs are toxic when fresh and are said to act as heart poison.

259 **Gloriosa superba** Linn (Vern —S -Sukra puspita, H -Kalihari, B -Bishlanguli, Bo.-Karianag, M -Agnishikha)

The root is stated to be sometimes used for suicidal purposes and as abortifacient, it is an acronarcotic poison The juice of leaves is stated to be used to destroy lice in hair

260 **Scilla iddica** Baker

..

261. **Urginia coromandeliana** Hook. f., *U indica* Kunth (Vern —S.-Vana-palandam, H & B.-Jangli piyaz, Bo.-Jangli-kanda, M -Nari-vengayam)

The bulbs are irritant poison The foreign species *U. scilla* is a fish poison, as also is the Indian representative

Juncaceae

(Rush Family)

262 **Juncus effusus** Linn

It is suspected to be poisonous to livestock in South Africa This and other species in India are worth investigating.

Palmaceae

(Palm Family)

Arecaïne, arecolidine, arecoline, guvacine, guvacoline, saponins

263 **Areca catechu** Linn (Vern —S -Gubak, H & B -Supari, Bo -Sopari, M Kamugu)

Young and undried nut when chewed in excess gives rise to temporary giddiness It also produces gripping and strong intestinal irritation, resulting in loose motions.

264. **Arenga obtusifolia** Mart.

Juice from fruit is used in Malaya to poison enemies and *A obtusifolia* is stated to be used as fish poison.

265 **Corypha umbraculifera** Linn. (Vern —S.-Alpayushi, M.-Tali-panai)

The fruit is stated to be used as fish poison.

266 **Wallichia disticha** T Anders.

Watt states that berries and perhaps the leaves are irritant to skin

Araceae

(Aroid Family)

Calcium oxalate (acicular crystals), bitter substance, sharp acrid substance, essential oil (alkaloid and saponin in foreign plant)

267 **Acorus calamus** Linn (Vern —H & B.-Bach, Bo -Vaj, M Vashambu), *A gramineus* Soland

Roots are stated to be used as effective isecticides and insectifuge. Doubtful case reported when *A calamus* proved poisonous to camels during the Afghan Campaign. The rhizome is used in medicine but in overdoses produces violent and persistent emesis.

268 **Alocasia indica** Schott (Vern —S.-Manaka, H -Mankenda, B -Mankachu), *A montana* Schott, *A odora* (Roxb) C Koch (-*A macrorhiza* Schott)

Fresh tubers are acrid and irritant

269 **Amorphonphallus campanulatus** (Roxb) Bl *A lyratus* Engl, *A sylvaticus* (Roxb) Kunth (*Sunantherias sylvatica* Schott)

Fresh tubers are acrid and irritant and the seeds are intensely acrid. Seeds of *A sylvaticus*, like *Plesmonium*, and fruit are intensely acrid.

270 **Arisaema speciosum** Mart (Vern —P -Kiralu), *A tortuosum* Schott (Vern —P -Samp-ki-kumb)

The tubers are poisonous and have insecticidal properties. The fruit is also probably poisonous.

271 **Homalomena rubescens** Kunth

It is stated to be poisonous

272 **Lagenandra ovata** (Linn) Thw (-*L toxicaria* Dalz) (Vern:—Bo -Rukh-ali, M.-Maravara Tsjembu)

It is stated to be very poisonous and has insecticidal properties.

273. **Plesmonium margaritiferum** Schott

Its crushed seeds produce local anaesthesia and it is used as a cure for toothache

274 **Sauromatum guttatum** Schott.

Tubers are regarded as very poisonous.

275 **Steudnera virosa** (Kunth) Prain (-*Colocasia virosa* Kunth.).

It is believed to be poisonous.

276 **Thomsonia nepalensis** Wall

It is acrid when fresh.

277. **Typhonium trilobatum** Linn. Schott. (Vern.—B -Ghet-kachu, M.-Karunaikkizhangu).

Fresh tubers are exceedingly acrid

Cyperaceae

(Sedge Family)

Essential oil.

278 **Carex cernua** Boott

It is said to be one of the causes of 'vlei' poisoning in cattle in South Africa.

279. **Cyperus longus** Linn

It is regarded as poisonous in South Africa.

280. **Scirpus corymbosus** Heyne

See *Carex cernua*.

Gramineae

(Grass Family)

Cyanogenetic glucosides, hydrocyanic acid, temuline, saponins, oxalic acid, selenium protein (toxic)

281 **Avena fatua** Linn (Vern —H -Kuljud), *A sativa* Linn

It is good fodder but occasionally deleterious probably on account of 'hair balls' that are developed in the stomach

282 **Bambusa bambos** Druce (-*B. arundinacea* Willd) (Vern:—S -Vansa, B & H -Bans, Bo.-Mandgay, M -Mangal).

Fresh young shoots are stated to have insecticidal properties

283. **Dendrocalamus strictus** (Roxb.) Nees (Vern:—H -Bans kaban, B -Karail, Bo -Bas, M.-Kanka).

The leaves are stated to be used to procure abortion

284 **Lolium perenne** Linn., *L. temulentum* Linn (H -Machni)

Several cases of poisoning, mostly non-fatal in man and animals, from eating the seeds of *L temulentum* have been recorded Gastro-intestinal irritation and severe nervous symptoms are reported.

285. **Panicum maximum** Jacq

It is suspected to be responsible for the production of 'Dikoor', a disease affecting young sheep in Africa.

286. **Paspalum scrobiculatum** Linn (Vern —S.-Kodrava, H.-Kodo, B -Kodoa, dhan, Bo -Kodra, M -Kiraruga).

The 'kodra' poisoning is very similar to *L temulentum* poisoning, the animals suffer much more than men. The animals should be prevented from grazing the crop when ripening.

- 287 **Sorghum halepense** (Linn) Pers (Vern —H -Baru, B -Kalamucha), *S saccharatum* Pers (Vern —H & Bo -Deo-dhan, M -Tellajonna), *S vulgare* Pers (Vern —S Javanala, H, B & Bo -Jowar, M -Cholam)

These are good fodder but occasional poisoning is reported with stunted growth, under drought conditions. The frosted leaves, or second growth is dangerous.

- 288 **Stipa sp** (some)

It is believed to be poisonous and mechanical action of 'seeds' may not be overlooked.

289. **Triticum aestivum** Linn (Vern —S -Godhum, H -Gehun, Bo -Gam, M -Godumai)

Under certain conditions it becomes deleterious fodder.

- 290 **Zea mays** Linn (Vern —S -Yavanala, H. & B -Bhutta, Bo.-Makai, M -Makka-schalam)

The pollen from it is stated to be a possible cause of hay fever. It is said to be occasionally responsible for deleterious effects, as yet not fully understood.

CHAPTER IX

We have studied the literature on Indigenous Systems of Medicine as practiced in India with a view to find out the uses to which different medicinal plants have been put in the treatment of various important diseases. For instance certain plants have the reputation of being effective in the treatment of bowel diseases particularly dysentery and cholera. Others are said to be useful in the treatment of prolonged fevers such as the group of enteric fever. Certain other plants are alleged to be effective in the treatment of tuberculosis (pulmonary). Yet another group is considered to have action on genital organs and emenagogue and abortifacient properties are attributed to them. Investigation on some of these plants has been carried out and brief summaries of the results achieved are given. Others marked with an asterisk have not been investigated so far and will form a field for research for the investigators.

Besides these certain plants are considered to have insecticidal and insect-repellent properties and another allied group is toxic to fishes. These, as has been already indicated, are a very important group from point of view of control of insect and other pests which do incalculable harm to man, livestock and agricultural crops.

The object of giving these lists is to bring these to the notice of those who are interested in indigenous drugs. To the lay public these will be of general interest, as they will find many names of common plants growing at their very door which have been considered effective in the treatment of common ailments and are household remedies. For the research worker these lists will open up a field for investigation and will give an idea as to the avenues in which investigations might be directed.

Indian Medicinal Plants Alleged to be useful in Tuberculosis in the Indian Indigenous Medicine.

- 1 ABRUS PRECATORIUS Linn
- 2 ADHATODA VASICA Nees
3. AEGLE MARMELLOS Corr
- 4 ALANGIUM SALVIFOLIUM (Linn) Wang Syn - *A. lamarckii* Thwaites.
- *5 ALBIZZIA JULIBRISSIN Durazz, (B - Kalkora, H - Lal siris), outer Himalaya from the Indus to Sikkim, ascending upto 6 000 to 7,000 ft
- *6 ALBIZZIA LEBBECK Benth, (S - Shirisha, H, B & Bo - Siris), throughout India, ascends to 4,000 ft in the Himalayas usually planted
- *7 ALLIUM CEPA Linn, (S - Palandu, H - Piyaz, B - Piyaj, Bo.-Kanda), extensively cultivated all over India.
- 8 ALLIUM SATIVUM Linn.

- *9. ALLIUM SCHOENOPRASUM Linn. , Kashmir to Kumaon 8,000 ft to 11,000 ft , grown as a garden crop
- 10. ALPINIA GALANGA Swartz
- 11. ASCLEPIAS CURASSAVICA Linn
- *12 ASPARAGUS RACEMOSUS Willd , (S & B -Shatamuli, H -Shatawar, Bo -Satavari) Himalayas, from Kashmir eastwards
- 13. AZADIRACHTA INDICA A Juss , Syn *Melia azadirachta* Linn
- *14 BAUHINIA RACEMOSA Lam , (S -Svetakanchan, H -Kanchnal) , throughout India.
- *15 BAUHINIA VARIEGATA Linn , (S -Kovidara, H -Kachnar, B -Rakta kanchan) , sub-Himalayan tract from Indus eastwards, dry forest of Eastern, Central & South India
- 16 BUTEA MONOSPERMA (Lam.) Kuntze
- *17 CANSCORA DECUSSATA Schult , (S -Sankhapuspi, H -Sankhaphuli) , throughout India upto 4,000 ft , grows in moist situations
- *18 CAPPARIS SPINOSA Linn , (S -Kakadani, H & P -Kabra) , plains between the Indus & Jhelum, Salt Range, Chamba, Kumaon, Nepal, Deccan, W Ghats, Baluchistan, Waziristan
- 19 CASSIA FISTULA Linn
- *20 CASSIA TORA Linn Syn -*C obtusifolia* Linn , (S -Chakramards, H & Bo -Chakunda) , throughout India
- *21 CHIONACHNE KOENIGII (Spreng) Thw *Polytoca barbata* Stapf (B -Gurgur, H -Kansa, S -Kanda) , throughout India in hot and damp parts
- 22 CIMICIFUGA FOETIDA Linn , (P -Junti)
- 23 CITRULLUS COLOCYNTHIS Schrad
- *24 CLERODENDRON SERRATUM (Linn) Moon , (H -Barnagi, S & Bo -Bharangi), more or less throughout India
- 25 CLITORIA TERNATEA Linn.
- 26 COCCINIA CORDIFOLIA Cogn. Syn -*C indica* W & A , *Cephalandra indica* Naud
- *27. COCCULUS HIRSUTUS Diels , (B -Hayer, Bo -Vasanvel, H.-Jamti-kibel), from the foot of the Himalayas to S India
- *28 COCOS NUCIFERA Linn , (H -Nariyal, Bo -Narel, M -Tanba), cultivated in the hot damp regions of India especially near the sea
- *29 COMMIPHORA AGALLOCHA Engler (B , Bo & H -Guggul), Assam, Sylhet, Bengal, Madhya Pradesh
- 30. COMMIPHORA MUKUL Engl , Syn.-*Balsamodendron mukul* Hook. ex Stock, Bellary, Mysore, Khandesh, Kathiawar, Rajasthan, Sind, Baluchistan.
- *31. CORIANDRUM SATIVUM Linn , (S.-Dhanyaka, H. & B -Dhania); extensively cultivated throughout India.

- *32. *CRATAEVA NURVALA* Ham ; (H -Barun, S -Varuna) almost all over India, wild or cultivated, often found along streams
- *33. *CRESSA CRETICA* Linn , (H & B -Rudranti, Bo -Khardi), throughout India
- *34. *CUMINUM CYMINUM* Linn , (S Jiraka, H -Jira, Zira, Tam.-Shiragam); cultivated throughout India (except Bengal & Assam),
- *35. *CURCUMA ANGUSTIFOLIA* Roxb , (H & B -Tikhur Tam -Kua), outer ranges of Central Himalayas, Bihar, Bengal
- *36. *CURCUMA ZEDOARIA* Bose
- *37. *DIOSPYROS MELANOXYLON* Roxb , (H -Kendu, B Kend, Bo -Temru), deciduous forests of the Madhya Pradesh, Bihar, W. Peninsula, Chota Nagpur.
- 38. *DRYNARIA QUERCIFOLIA* J Sm
- 39. *ELETTARIA CARDAMOMUM* Maton
- *40. *EMBELIA TSJERIAM-COTTAM* A Dc (H -Bayabirang, Bo -Barbatti) Ceylon, Malabar Coast, Sylhet and Assam
- *41. *EULOPHIA NUDA* Lindl , (S -Manya, H -Goruma, H -Badbar) Tropical Himalaya, Nepal to Sikkim, Assam, Khasia Hills, Chota, Nagpur, Manipur, W Peninsula
- *42. *FRITILLARIA CIRRHOSA* Don Prodr , Central & E Himalayas, Sikkim, 11,000 to 16,000 ft
- *43. *FRITILLARIA ROYLEI* Hook , Kashmir to Kumaon, 8,000 to 13,000 ft
- *44. *GERANIUM ROBERTIANUM* Linn , Kashmir to Garhwal
- 45. *GLYCYRRHIZA GLABRA* Linn
- *46. *GMELINA ARBOREA* Linn , (S -Gumbhari, H -Kambari), throughout India
- *47. *GOSSYPIUM ARBOREUM* Linn , (H -Nurma, P -Papasa), Indian gardens.
- *48. *GRAPTOPHYLLUM PICTUM* (L) Griff , (M.-Ysjudemaram), Indian gardens
- *49. *GREWIA ASIATICA* Linn (S -Parusha, H. & B.-Phalsa) , extensively cultivated throughout India
- *50. *HAUCLEA SESSILIFOLIA* Roxb
- *51. *HYSSOPUS OFFICINALIS* Linn ; (Zufah-Yabis), Kashmir to Kumaon, 9,000 to 11,000 ft
- *52. *INDIGOFERA TINCTORIA* Linn , (S -Nilika, H & Bo.-Nil) , widely cultivated in many parts of India
- *53. *JASMINUM AURICULATUM* Vahl , (S. & Tel , Magadhi) ; Deccan, Carnatic, W. Peninsula.
- *54. *LACTUCA SERRIOLA* Linn. Syn *L scariola* Linn. (H -Kahoo, B.-Salad); W Himalaya, 6,000-12,000 ft.

- *55 LEEA AEQUATA Linn , (S , H. & B -Kakajangha), Sikkim, Himalaya, Assam, East Bengal, Sylhet, Andamans.
- 56. LINUM USITATISSIMUM Linn
- *57. LITSEA CHINENSIS Lam (H -Garbijaur, B -Kukarchita), throughout the hotter parts of India
- *58. LORANTHUS ASPER Lour
- 59. LUFFA ACUTANGULA var *amara* Clarke
- 60 LUFFA ECHINATA Roxb Host Beng
- 61 LUVUNGA SCANDENS Ham.
- *62 MACHILUS MACRANTHA Nees. , (Tam -Kolamavu, Mal -Uravu) , W Peninsula
- *63 MARTYNIA ANNUA Linn , (H -Bichu, B -Baghnoki, Tel -Garudamukku) , naturalized in India
- *64 MELISSA PARVIFLORA Benth , (H -Bihlotan), Garhwal to Sikkim & Mishmi, Khasia hills
- *65 MIMUSOPS HEXANDRA Roxb , (S -Rajadani, H -Khirmi, M -Palla) , upper Gangetic plain, Madhya Bharat, S India
- *66 MOMORDICA DIOICA Roxb , (S -Vahisa, M -Palupaghel kalung) , throughout India, ascending to 5,000 ft in Himalayas
- 67 MORINGA OLEIFERA Lam
- 68 MUSA SAPIENTUM O Kuntze.
- *69 OCHNA PUMILA Ham ex D Don , (Santh -Champabaha), Kumaon to Sikkim, Bihar, Chota Nagpur
- *70. OLDENLANDIA UMBELLATA Linn , (M.-Saya, H -Chirval) , Orissa, Bengal, Deccan, Circars, Carnatic
- 71. OPERUCINA TURPETHUM (Linn) Silva Manso.
- *72 PHASEOLUS ACONITIFOLIUS Jacq (H -Mat, Tam -Tulkapyre), Himalayas to Ceylon
- *73. PHASEOLUS TRILOBUS Art , (H. & B.-Mugani) , Himalayas to Ceylon and Burma
- *74 PIPER CHABA Hunter (H B & Bo.-Pan) , cultivated in various parts of India
- *75 PIPER LONGUM Linn., (S -Pippali, H.-Pipal), hotter states of India.
- 76. PINUS ROXBURGHII Sargent, Syn.-*P longifolia* Roxb.
- 77 PISTACIA INTEGERRIMA Stew.
- *78 PISTIA STRATIOTES Linn , (H -Jalqumbhi, Bo -Prashni), throughout India in still sweet waters
- *79 PLANTAGO MAJOR Linn , (H -Lahuriya, Bo Bartang), Assam, Khasia hills, W Ghats, Konkan, Nilgiris, Pulneys, Baluchistan
- 80. PLUMBAGO ZEYLANICA Linn.

- *81 RHODODENDRON CAMPANULATUM Wall. ex G Don., (H.-Cherailu, Kash -Gaggar), alpine regions from Kashmir to Bhutan
- 82. RHUS SUCCEDANEA Linn
- *83 SANSEVIERIA ROXBURGHIANA Schult , (B.-Murba, Bo -Morwa), Coromandel coast
- *84 SANTALOIDES MINUS Schellenb. Syn -*Rourea santaloides* W & A , (B -Vitaraka, Bo -Vardara), Konkan, S M Country, Kanara to Travancore
- *85 SAPINDUS TRIFOLIATUS Linn (S -Phemila H , B & Bo -Ritha).
- *86. SESBANIA SESBAN (Linn) Merr Syn -*S aegyptiaca* Pers , (H & B -Jayanti), plains from the Himalayas to Ceylon, ascending upto 4,000 ft in north-west
- 87. SIDA CORDIFOLIA Linn
- 88 SIDA RHOMBIFOLIA Linn
- *89 SOLANUM TRILOBATUM Linn , (S -Alarka, M -Tudovullay), Gujerat, Deccan, S M Country, N Circars, Carnatic
- *90. SPHAERANTHUS INDICUS Linn , (S -Munditika, H & Bo -Gorakmundi), throughout India, ascending the Himalayas upto 5,000 ft. from Kumaon to Sikkim
- *91. SPONDIAS PINNATA Kurz , (S -Amrataka, H , B & Bo -Amra), from the Chenab eastwards, Salt Range, Andamans, W Indian Peninsula
- 92 STEPHANIA GLABRA Roxb
- *93. STREBLUS ASPER Lour , (H. Siora, B -Sheora), drier parts of India, Travancore, Andamans
- *94. TERAMNUS LABIALIS Spreng , (S -Masha-Parui, H -Mashparui), throughout India
- *95. URTICA DIOICA Linn., (H & P -Bichu), N W. Himalaya, from Kashmir and the Salt Range to Simla, 8,000-10,700 ft.
- *96 VANDA SPATHULATA Spreng , (Mal -Ponnam-penmaraiva), W Peninsula from Malabar to Travancore
- *97. VATERIA INDICA Linn , (S -Ajakarua, H -Safed damar), Western India, from N. Kanara to Travancore, chiefly in evergreen forests, Coorg.
- 98 VERBASCUM THAPSUS Linn
- *99 VERNONIA CINEREA Less., (S & H -Sahadevi, B -Kukseem), throughout India, ascending to 8,000 ft in the Himalaya Khasia and Peninsula mountains
- *100. VERNONIA ROXBURGHII Less, Upper Gangetic Plain, Kumaon, Bengal, Central & W. India.
- 101 WITHANIA SOMNIFERA Dunal.
- *102. ZIZYPHUS JUJUBA Lam., (S -Badari, H.-Baer, B -Kul), indigenous and naturalized throughout India, in the Outer Himalaya upto 4,500 ft.

Indian Plants Considered to be useful in Dysentery in Indian Indigenous Medicine.

- I ABRUS PRECATORIUS Linn.
- *2 ABUTILON INDICUM Sw , (H -Kanghi, B.-Potari), throughout the hotter parts of India
- *3 ABUTILON THEOPHARSTIC Medic, (S -Jaya, Bo -Nahani khapat), N W India, Sind, Kashmir, Bengal
- *4 ACACIA ARABICA Willd , (H -Kikar, B -Babla), naturalized in all parts of India, indigenous to Sind and the Deccan.
- *5 ACACIA CATECHU Willd , (H -Khadira, H -Khair), Punjab, N. W. Himalaya, Central India, Bihar, Ganjam, Konkan, Deccan
- *6 ACACIA FERRUGINEA DC , (Nep -Khour, Bo -Ker), Gujerat, Berar, Circars, Deccan, Konkan, Carnatic, W Ghats
- 7 ACORUS CALAMUS Linn
- *8 ADENSONIA DIGITATA Linn , (H & Bo -Gorakh-amli), occasionally cultivated in some parts of Uttar Pradesh, Bihar, Bombay & Madras
- 9 ADHATODA VASICA Nees
- *10 ADIANTUM LUNULATUM Burm , (H & B -Kali-Jhant, Bo -Hansrai), throughout N India in moist places, South India
- 11 ADINA CORDIFOLIA Benth
- 12 AEGLE MARMELOS Corr.
- *13 AGARICUS OSTREATUS (Jacq) Fries., (Bo -Phanasa-alambe)
- *14 AGERATUM CONYZOIDES Linn ; (B -Dochunty, Bo -Osari), throughout India upto 5,000 ft
- 15 AILANTHUS ALTISSIMA (Mill) Swingle, Syn — *A glandulosa* Desf ; (English-Ailanto).
- *16 AILANTHUS EXCELSA Roxb , (H -Mahanimb, Tam -Peruppi), Bihar, Chota Nagpur, Madhya Pradesh, Ganjam, Vizagapatnam, Deccan
- *17 AILANTHUS MALABARICA DC , (Bo -Guggula-dhup Tel -Maddi-palu), Konkan, Deccan, W Ghats.
- 18 ALBIZZIA LEBBECK Benth
- 19 ALSTONIA SCHOLARIS R Br.
- *20 ALTHAEA ROSEA Cav , Indian Gardens
- *21 AMARANTHUS TRICOLOR Linn Syn — *A gangeticus* Linn.; (H -Lalsag, S -Marisha), cultivated throughout India
- *22 AMOMUM XANTHIODES Wall., (H -Ilayechi, Tam.-Elam), Tavoy, Tenasserim.
- 23 ANDROGRAPHIS PANICULATA Nees, (S.—Bhunimba, H -Kiryat).
- *24. ANNONA MURICATA Linn ; (S -Mullanjakka, Tam.-Mullu-chitta); grown to a small extent in Assam.

25. *ANNONA RETICULATA* Linn
- *26. *ANTHOCEPHALUS INDICUS* A. Rich Syn - *A. cadamba* Miq S. & H. - Kadamba), sub-Himalayan tract from Nepal to Burma, Circars and W. Ghats.
27. *ANTIARIS TOXICARIA* Leschen
- *28. *ASPARAGUS ADSCENDENS* Roxb, (H - Safed musli, Gharwal-Jhirna), Punjab & the Himalayas upto 5,300 ft
29. *ASPARAGUS RACEMOSUS* Willd
- *30. *ASTERACANTHA LONGIFOLIA* Nees, (H - Tal-makhana, S - Kakil-akshya) throughout India in moist places
- *31. *AVERRHOA CARAMBOLA* Linn, (H - Karmal, M - Tamarta), gardens in hotter parts of India as far north as Lahore
32. *BALANITES AEGYPTIACA* Del Syn - *B. roxburghii* Planch.
- *33. *BARRINGTONIA ACUTANGULA* (Linn) Gaertn (B - Hijal, H - Hijjal), common in the sub-Himalayan tracts east of the Madhya Pradesh, Jumna, Bihar, Orissa, Bengal, Assam, S. India
- *34. *BASELLA RUBRA* Linn, (S - Potaki, H Lalbachlu), throughout India, wild & cultivated
35. *BAUHINIA RACEMOSA* Lam
- *36. *BAUHINIA TOMENTOSA* Linn, (S - Asw. mantaka, H - Kachnar, Bo - Asundro, M - Kanchnine), N W States, Circars, Carnatic, often cultivated in other parts of India
37. *BAUHINIA VARIEGATA* Linn
- *38. *BERGENIA LIGULATA* (Wall) Engl. (B - Patharchuri, Bo Pashanbheda), temperate Himalayas, from Kashmir to Bhutan, between 7,000-10,000 ft Khasia hills
- *39. *BIDENS TRIPARTITA* Linn, Central & W Himalaya from Nepal to Kashmir, W Tibet
- *40. *BOSWELLIA SERRATA* Roxb, (S - Shallaki, H & B - Luban), Madhya Pradesh, Deccan, Bihar, Orissa, Rajasthan, Madhya Bharat, Gujerat
- *41. *BOTRYCHIUM LUNARIA* Sew, (English-Moonwort), common from Kashmir to Sikkim ascending upto 13,000 ft, extending to Korakoram Range
42. *BRASSICA CERNUA* (Thunb) Forbes & Hemsley
43. *BRUCEA AMARISSIMA* (Lour) Merr Syn - *B. sumatrana* Roxb.
44. *BUTEA MONOSPERMA* O Kuntze.
- *45. *CAESALPINIA JAYABO* Maza; (Arab - Bunduk, Tanu.-Kalarisikkodi), Indian Peninsular region
- *46. *CALAMUS ROTANG* Linn, (S.-Vetasa, H., B & Bo.-Bet), Madhya Pradesh, Deccan, Carnatic.
47. *CALOTROPIS GIGANTEA* R. Br.

48. CARAPA MOLUCEENSIS Lam.
49. CARUM BULBOCASTANUM Koch (H.-Kalajira, Kash -Guniyan); Kashmir, Baluchistan
50. CARUM CARVI Linn
- *51. CASSIA AURICULATA Linn., (H & B -Taruar, Tam.-Avaram), wild in the dry regions of Madhya Pradesh, W Indian Peninsula, Rajasthan, cultivated in other parts of India.
52. CASSIA FISTULA Linn
53. CASSIA TORA Linn
54. CASSYTHA FILIFORMIS Linn
- *55. CASUARINA EQUISETIFOLIA Linn ; (H -Jang'isaru, B -Belatijau) , from Chittagong Southwards
56. CEDRELA TOONA Roxb
- *57. CELSIA COROMANDELIANA Vahl (S -Kulahala, Bo -Kolhal), throughout India
58. CENTELLA ASIATICA (Linn.) Urban Syn —*Hydrocotyle asiatica* Linn.
- *59. CEROPEGIA TUBEROSA Roxb.; (P -Galot, Bo -Khappar kadu, M -Manda), Konkan, Deccan, S. M. Country Circars.
- *60. CINNAMOMUM INERS Reinw , (Bo.-Tikhī, H -Janglī darchinī), ever-green forests of W Ghats, Carnatic, Sheoroy & Kollimalai
61. CLITORIA TERNATEA Linn
- *62. CORALLOCARPUS EPIGAEUS Benth ex Hook f , (S -Patalagaruda) , Punjab, Sind, Gujerat, Rajputana, Deccan, Carnatic.
- *63. CORCHORUS CAPSULARIS Linn , (S -Kalasaka, H. & B.-Narcha) , throughout the hotter parts of India
- *64. CORCHORUS FASCICULARIS Lam , (B.-Bilnalita, Bo -Hirankhorī) , throughout the hotter parts of India
65. CUMINUM CYMINUM Linn
- *66. CYDONIA OBLONGA Mill Syn -*C vulgaris* Pers. , (H -Bihī, S.-Amritphala) , cultivated in Punjab, Kashmir, Nilgiris
- *67. CYLISTA SCARIOSA Roxb. (Bo -Ranghevada) , Madhya Pradesh, West & South India.
68. CYNODON DACTYLON Pers (H., B & Bo -Dhub) , throughout India ascending to 8,000 ft. in the Himalaya
- *69. CYPERUS ROTUNDUS Linn , (S & Bo -Musta, B & H -Mutha) , throughout India, common in waste grounds, gardens, road sides, in open spots & upto an elevation of 6,000 ft
- *70. CYPERUS SCARIOSUS R. Br ; (S -Nagar mustaka, H & B -Nagar Motha) damp places in Bengal, Uttar Pradesh, eastern & southern parts of India.

- 71 DAUCUS CAROTA Linn
- *72 DESCURAINIA SOPHIA Linn , (H.-Khubkallana), Kashmir to Kumaon upto 14,000 ft , E Himalaya, Salt Range Peshawar, Baluchistan
- *73 DESMODIUM GANGETICUM DC , Outer Himalaya upto 5,000 ft. & throughout India
74. DIOSPYROS MELANOXYLON Roxb
- *75 DIOSPYROS PEREGRINA Gurkeim , (S -Tinduka, H & B -Gab, Bo -Tendu) , throughout India
- *76. ELAEOCARPUS SERRATUS Linn ; (B -Julpai, M -Olang-karai) W Indian Peninsula
- 77 ERIGERON CANADENSIS Linn
- 78 EUPHORBIA ANTIQUORUM Linn
- 79 EUPHORBIA HIRTA Linn
80. EUPHORBIA HYPERICIFOLIA Linn
81. EUPHORBIA THYMIFOLIA Linn
- *82 EVOLVULUS ALSINOIDES Linn , (S -Vishnugandhi, H -Sankha pushpi) Tropical and sub-tropical regions of India
83. FAGONIA CRETICA Linn
- *84 FERONIA LIMONIA (Linn) Swingle, Syn —*F. elephantum* Corr , (Kapittha, H -Kavitha) ,indigenous in S India, cultivated in many parts of India.
- *85 FICUS BENGALENSIS Linn , (S -Vata, H -Bor, B -Bar) , sub-Himalayan tract, W Peninsula, planted elsewhere
- *86. FICUS HETEROPHYLLA Linn , (S -Trayamana, B -Bhui-dumur) , throughout hotter parts of India
- *87. FICUS HISPIDA Linn , (S -Kakadumbura, H -Konea-dumbar) more or less throughout India
- *88. FICUS RACEMOSA Linn Syn —*F. glomerata* Roxb , (S -Udumbara, H -Gulari) , throughout India
- *89. FIMBRISTYLIS JUNCIFORMIS Kunth , (Santh -Bindimuthi), Madhya Pradesh, W Peninsula.
90. FLEMINGIA TUBEROSA Dalz (Bo -Birmova), Konkan
91. FOENICULUM CAPILLACEUM Gilb , Syn —*F. vulgare* Gærtn
92. GARCINIA MANGOSTANA Linn , (H., B & Bo -Mangustan), cultivated in Madras State, Nilgiris
- 93 GASTROCHILUS PANDURATA Ridley's , Konkan, Andamans
94. GOSSYPIUM HERBACEUM Linn. , (H., B & Bo -Kapas, S -Karpas), cultivated in N. W Frontier region of Pakistan, Baluchistan
95. GREWIA HIRUSTA Vahl.
96. GREWIA MICROCOS Linn.

- 97 *HELICTERES ISORA* Linn
- 98 *HEMIDESMUS INDICUS* R Br.
- *99 *HIBISCUS CANNABINUS* Linn. , (S -Nali, H -Patsan, Bo -Ambari) ;
generally cultivated.
- 100 *HOLARRHENA ANTIDYSENTERICA* Wall.
101. *HOLARRHENA MITIS* R Br (Sinhalese-kiriwolla) , endemic in
Ceylon
102. **HYDROCOTYLE JAVANICA* Thunb
- *103 *HYPERICUM JAPONICUM* Thunb , Khasia Hills, Assam, E & W.
Peninsula
- *104 *INDIGOFERA OBLONGIFOLIA* Forsk. , (S -Jhilla) , throughout the
Indian plains, Baluchistan
- *105 *IXORA NIGRICANS* Br , (Tam -Mashagani) , E. & W. Indian
Peninsula
- *106 *JATEORHIZA PALMATA* (Lam) Miers, (Bo -Colombo, Tam -Kolu-
mbu) cultivated in some parts of India
- 107 *JATROPHA CURCAS* Linn
- 108 *JUGLANS REGIA* Linn
- *109 *JUSSIAEA SUFFRUTICOSA* Linn (S -Bhallava-anga, H -Banlaunga) ;
throughout India
- *110 *JUSTICIA GENDARUSSA* Burm (S -Nila-nirgundi, H -Nili-nargandi) ,
cultivated throughout India
- *111 *LEEA INDICA* Merrill , (H & B -Kakurjiwan, Bo -Karkani) ,
throughout India, Andamans
- *112 *LENS CULINARIS* Medic Syn.—*L esculenta* Moench , (B -Masuri, H.
& S -Masur) , cold weather crop throughout India
- 113 *LITSEA CHINENSIS* Lam
- 114 *LUFFA ACUTANGULA* var *amara* Clarke
- *115 *MALVA ROTUNDIFOLIA* Linn , (H & Bo -Khubazi) , Sind, Balu-
chistan, Waziristan, plains of N India, ascending to 10,000 ft ,
Kumaon.
- *116 *MANGIFERA INDICA* Linn , (H B & Bo -Amb, S -Amva), indigenous
in Sikim, Assam, Khasia hills, Khandesh, along the W. Ghats,
cultivated in the tropical regions
- *117 *MELASTOMA MALABATHRICUM* Linn , Burma Myetpye, Tel -Pattudu),
throughout India except the Desert
- *118. *MENTHA LONGIFOLIA* Hunds Syn — *M -sylvestris* Linn. (Bo , H & S -
Pudina), W. Himalaya, 4,000 to 12,000 ft
- *119 *MESUA FERREA* Linn (S , H. & B -Nagkeshar, Bo -Nagchampa),
Mountains of E Himalaya & E Bengal, Assam, ever green rain
forests of N. Kanara & S. Konkan, forests of W Ghats, Andamans.

- *120. *MIMOSA PUDICA* Linn.; (S -Lajja, H. & Bo.-Lajalu); natural:rel more or less throughout India
- *121. *MIMUSSOPS ELENGI* Linn., (S , H & B -Bakul Bo -Borasali), Andamans, W Peninsula, Khandala Ghats, Circars
- *122. *MORINDA CITRIFOLIA* Linn , (H & B -Ach, Bo-Aal), indigenous in the Darjeeling Terai and Andamans and along the Konkan coast, cultivated largely in India
- *123. *MORINDA UMBELLATA* Linn., (Bo -Al, S -Pitadaru), Khasia, hills, Madras State, E Ghats , W Ghats
- 124. *MUSA SAPIENTUM* O Kuntze
- *125. *MUCUNA PRURITA* Hook , (S -Atmagupta, H -Kiwach), Punjab plain, from base of Himalayas to Ceylon & Burma.
- *126. *MURRAYA KOENIGII* Spreng , (S -Sourabhi nimba, H -Katnim), Konkan, W Ghats, Deccan, S M Country, Madras State, along the foot of Himalaya from Kumaon to Sikkim, upto 5,000 ft., Bengal
- *127. *MURRAYA PANICULATA* Jack, (H Marchula, B -Kamini), outer Himalaya from the Jumna eastwards, ascending to 4,500 ft , Assam Satpura Range, Peninsula
- 128. *MYRICA NAGI* Thunb , (S -Katphala, H , B & Bo -Kaiphali)
- *129. *MYRTUS COMMUNIS* Linn , (H -Vilayiti mehdi, B -Sutr-sowa), indigenous from the Mediterranean to N W Himalaya, gardens, throughout India
- *130. *NANNORHOPS RITCHIEANA* H Wendl , (H Mazri), Punjab, Sind, Waziristan, Baluchistan
- *131. *NEPETA ELLIPTICA* Royle ex Benth , (P -Tukhmalanga), W. temperate Himalaya, from Kashmir to Garhwal, 6,000 to 8,000 ft.
- *132. *NYMPHAEA ALBA* Linn , (Kash -Brimposh, Bo -Pandharen-kamal), Kashmir lakes
- *133. *NYMPHAEA PUBESCENS* Willd , (M -Alli), all over India in the warmer parts
- *134. *OCIMUM AMERICANUM* Linn Syn -*O canum* Sims , (H. & B.-Kala tulshi), plains and lower hills of India
- *135. *OCIMUM BASILICUM* Linn , (S -Munjariki), indigenous on the lower hills of the Punjab, cultivated throughout the greater parts of India
- 136. *OLDEN LANDIA AURICULARIA* K Schum.
- *137. *OROXYLUM INDICUM* Vent , (S.-Syonaka, H -Arlu, B - Sona), throughout India, except in the Western drier area
- *138. *OXALIS CORNICULATA* Linn., (S -Amlika, H. & B -Amrul); nearly all regions throughout the warmer parts of India, in the Himalayas upto 8,000 ft.

- *139. *PAVONIA ODORATA* Willd., (S -Harivera, B -Bala, Bo.-Kala vala)
N W. India Bundelkhand, Sind, Baluchistan, Rajasthan, Bengal,
Konkan, S M Country, Deccan
- 140. *PEUCEDANUM GRAVEOLENS* Benth
- 141. *PHASEOLUS TRILOBUS* Ait
- *142. *PHYLLANTHUS EMBLICA* Linn , (S -Dhatraphala, H -Aoula, Bo
Amla), throughout tropical India, wild or planted
- *143. *PHYLLANTHUS NIRURI* Linn , (S -Bhumya-amalaki, H.-Jar-amlal),
throughout the hotter parts of India from the Punjab to Assam and
southwards to Travancore
- 144. *PHYLLANTHUS URINARIA* Linn
- 145. *PIPER LONGUM* Linn
- *146. *PIPER NIGRUM* Linn , (S -Maricha, H -Golmirch, B -Golmarich), culti-
vated in hot damp parts of India
- 147. *PISTACIA INTEGERRIMA* Stew
- 148. *PISTIA STRATIOTES* Linn
- *149. *PLANTAGO CILIATA* Desf , Punjab hills, Sind, Baluchistan
- 150. *PLANTAGO MAJOR* Linn.
- 151. *PLANTAGO OVATA* Forsk
- *152. *PLATANUS ORIENTALIS* Linn (P & Kash -Buin), N W Himalaya,
5,000 to 8,000 ft., cultivated only
- 153. *PLUMBAGO ZEYLANICA* Linn
- 154. *PSIDIUM GUAJAVA* Linn
- *155. *PULICARIA DYSENTERICA* Gaertn Kashmir at 5,000-6,000 ft
- *156. *PUNICA GRANATUM* Linn , (S -Darimba, H -Anar) wild in Salt range
and in the Himalayas from 3 000 to 6,000 ft , also cultivated in
many parts of India.
- 157. *RHEUM EMODI* Wall
- *158. *RUBIA CORDIFOLIA* Linn , (S & B -Manjistha, H.-Manith, Bo -
Manjit), throughout India in hilly districts
- *159. *RUMEX SCUTATUS* Linn., (English-French sorrel), W Himalaya
- *160. *RUMEX VASICARIUS* Linn , (S -Chukra, H , B & Bo -Chuka), indig-
enous to W Punjab, the Salt Range & Trans-Indus Hills, also
cultivated.
- *161. *SALIX ALBA* Linn , (P -Bis, Kash -Vuir), cultivated in N. W.
Himalaya
- 162. *SAPINDUS TRIFOLIATUS* Linn.
- 163. *SARACA INDICA* Linn.
- *164. *SCINDAPUS OFFICINALIS* Schott., (H & B.-Gajapipal. Bo -Thora-
pimple), from Sikkim eastwards, Bengal, Chittagong, Andamans.

165. SEMECARPUS ANACARDIUM Linn f
- *166. SHOREA ROBUSTA Gaertn f, (S, H, B & Bo Sal), Punjab, along the Sub-Himalayan tract to Assam, Garo Hills, Orissa, Khasia hills, Jaintia hills, Madhya Pradesh
- *167. SIDA SPINOSA Linn.; (S -Nagabala, H -Gulsakari, B -Bonmethi), throughout the hotter parts of India from N W. India to Ceylon.
- *168. SMILAX PROLIFERA Roxb., (H -Ram dataum), Kumaon, Nepal Sylhet, Bengal, Bihar, Deccan, Peninsula
- *169. SOLANUM NIGRUM Linn, (S & B -Kakmachi, Bo -Mako), throughout India, upto 9,000 ft in the W Himalayas
- *170. SOYMIDA FEBRIFUGA A Juss, (S Rohuna, H, B & Bo -Rohan), dry forests of W Peninsula extending to Merwar & Chota Nagpur
- *171 SPILANTHES ACMELLA Murr, (Bo & P -Akarkara), throughout India, upto 5,000 ft in the Himalayas & other mountains
- *172. STACHYTARPHETA JAMAICENSIS Vahl var *indica* Lam Syn —*S indica* Vahl, (M -Simainayuruvi), tropical India from the Punjab & Sylhet to Travancore
- 173 STRYCHNOS NUX-VOMICA Linn
- 174 SYMPLOCOS RACEMOSA Roxb
- *175 SYZYGIUM CUMINI (Linn) Skeels Syn —*Eugenia jambolana* Lam, (H & B -Jam), throughout India
- *176 SYZYGIUM JAMBOS (Linn) Alst Syn —*Eugenia jambos* Linn, (H & Bo -Gulabjaman), Sikkim Terai
- *177 SYZYGIUM OPERCULATUM Gamble, Syn —*Eugenia operculata* Roxb, (H -Rai-jaman, S -Bhumbi jambu), sub-Himalayan tract, common in Savannahs in Sal forests, Bihar, Assam, Sylhet, Chittagong
- *178 TECTONA GRANDIS Linn f, (S -Saka, H & B -Segun), Konkan, W Ghats of Bombay & Madras States, Deccan, Carnatic, Madhya Bharat, Circars.
179. TERMINALIA ARJUNA W & A
- 180 TERMINALIA CHEBULA Retz
181. TERMINALIA CITRINA Flenning
- *182 TRIUMPHETA BARTRAMIA Linn, (H -Chikti, B -Bun-akora), throughout tropical & sub-tropical India, ascending to 4,000, ft in the Himalaya
- 183 TYLOPHORA ASTHMATICA Wight & Arn.
- *184. TYPHA ELEPHANTINA Roxb., (B -Hegla, Bo -Ramban, Kash -Pitz), marshes from N. W. India to Assam & Southwards, Indus Delta.
- *185 URARIA LAGOPOIDES DC, (Bo.-Dowla, H. Pithvan); Tropical zone, Nepal, Chota Nagpur Bengal to Ava.

- 186 VATERIA INDICA Linn
- '187 WOODFORDIA FRUTICOSA Kurz , (S -Dhatakī, H & B -Dhai), throughout India, Baluchistan
- '188 WRIGHTIA TINCTORIA R Br , (S -Asita kutanja, H -Mitha indarjou, Bo -Indrajau), Rajasthan, Madhya Pradesh, Deccan, Konkan, S M Country, Circars, W Ghats of Madras State

Indian Plants considered to be Useful in Cholera and Prolonged Fevers such as the Enteric Group

- 1 ACONITUM FEROX Wall
- *2 ARTABOTRYS SUAVEOLENS Blume, Chittagong, Sylhet
- *3. BLUMEA LACERA DC , (B —Kukursunga, Bo —Nimrudī), throughout plains of India, upto 2,000 ft
- *4 CAPPARIS ZEYLANICA Linn , (H -Ardanda, B -Kalu-kera), Indian gardens
- *5 CAPSICUM FRUTESCENS Linn Syn.-*C minimum* Roxb , cultivated, but not extensively
- *6 CARAPA MOLUCCENSIS Lam , (B -Pussar), Littoral forests of Bengal and Andamans
- 7 CASSIA ANGUSTIFOLIA Vahl
- *8. DESMODIUM GANGETICUM DC (S & Bo -Shal Parnī, H -Sarivan), outer himalayas and throughout India
- *9 DRYNARIA QUERCIFOLIA J Sm (Bo -Basingh, S -Ashvakatri), throughout plains of India, on trees or rocks
- *10. ELAEOCARPUS TUBERCULATUS Roxb (S Rudraksha, M -Rutthraṣham), Western Indian Peninsula
- *11. ERYCIBE PANICULATA Roxb , (Santh -Kari, Tam -Unankodī), throughout India
- *12. ERYTHROXYLUM MONOGYNUM Roxb (M Devadarum), Madras, N Circars dry evergreen forests of Deccan and Carnatic, W Ghats, dry hill forests of Travancore
- 13 EUPHORBIA HELIOSCOPIA Linn
- *14. FAGONIA CRETICA Linn. (H -Damahan, P -Dama); Deccan, W Khandesh, Cutch, Sind, Baluchistan, Waziristan, W Rajasthan, Upper gangetic plain, Punjab
- *15. FLACOURTIA INDICA Merr Syn -*F ramontchi* L' Herit , (B Bincha); Sub-Himalayan tract, Upper Gangetic Plain, common in Peninsula, W. Ghats, Forests of the N Circars and Deccan upto 3,000 ft
- *16. GREWIA HIRUSTA Vahl. Symb (Bo -Gowali H Kukarbicha), Sub-Himalayan tract upto 4,500 ft. from the Indus eastwards, Salt Range, Bihar, Orissa.

- *17 GREWIA MICROCOS Linn (M -Kottei), E. Bengal, Assam, W. Peninsula, Mysore.
- 18. KALANCHOE SPATHULATA DC.
- *19 LODOICEA MALDIVICA Pers. Syn -*L seychellarum* Labill. (H & Bo -Darya-kanariyal), cultivated in India
- *20 MARRUBIUM VULGARE Linn , (H -Pahari gandana), Kashmir from 5,000 to 8,000 ft Baluchistan etc.
- *21. MOMORDICA CHARANTIA Linn. (H -Karela, B -Karala, Bo -Karla), cultivated throughout India
- *22. MUSA PARADISIACA Linn var *sapientum* Kuntze Syn -*M sapientum* Linn. (S -Ramoha, H & Bo -Kela), commonly cultivated.
- *23. OLDENLANDIA AURICULARIA K Schum , (B -Muttia-lata, Bo.-Dapoli), almost all over India
- *24 POINCIANA PULCHERIMA Linn Syn -*Caesalpinia pulcherima* Swartz (B -Krishnachura, S & Tam -Ratna-gandhi), Gardens throughout India.
- *25. PSIDIUM GUAJAVA Linn. (H -Amrud, Bo -Perala), cultivated & naturalized throughout India
- 26. SAPINDUS TRIFOLIATUS Linn
- 27 SCHLEICHERA OLEOSA (Lour) Merr Syn -*S trijuga* Willd
- *28 SCHWEINFURTHIA SPHEROCARPA Br , (S H & Bo -Sanipat), Rajasthan Desert, Sind, Baluchistan
- *29 SOLANUM MELONGENA Linn , (S.-Bartaku, H -Baigun), widely cultivated in India
- * 30. SOPHORA TOMENTOSA Linn , (Burm -Thimbawmagyi), Andaman & Nicobar Islands, very occasionally in Indian gardens.
- 31. STYRCHNOS NUX-VOMICA Linn
- 32 TERMINALIA CHEBULA Retz.
- *33 TERMINALIA CITRINA Fleming , (H -Harira, B -Haritaki), Assam, Dacca, Mymensingh
- *34 TERMINALIA PANICULATA Roth , (Bo -Kindal, M -Pekarakai; Western regions of the Peninsula from Bombay through Kanara & Malabar to Travancore, upto 2,000 ft., Coorg, Nilgiris, Anamalais Cuddapah, Ballary
- 35 TRACHYSPERMUM AMMI (Linn) Sprague, Syn.-*Carum copticum* Benth.
- *36. ZANTHOXYLUM BUDRUNGA Wall., (Bazinali, Bo -Tessul, H -Badrang), Konkan, Deccan, S. M. Country, N Kanara, W Ghats in S Kanara, Mysore, Malabar, Annamalais and Travancore at low elevations, Orissa, Sylhet, Khasia Hills, Chittagong.

Indian Plants considered to have Emmenagogue and Abortifacient Properties

1. ABRUS PRECATORIUS Linn.
2. ALOE BARBADENSIS Mill.
3. ANANAS COMOSUS Linn
4. ANNONA SQUAMOSA Linn.
5. APIUM GRAVEOLENS Linn.
6. ARECA CATECHU Linn
7. ARISTOLOCHIA BRACTEATA Retz.
8. ARISTOLOCHIA INDICA Linn
9. ARTEMISIA VULGARIS Linn.
10. CALOTROPIS GIGANTEA (Linn.) Dryand (*C. gigantea* R. Br.)
11. CALOTROPIS PROCERA (Linn.) Dryand (*C. procera* R. Br.)
12. CARICA PAPAYA Linn.
13. CELASTRUS PANICULATUS Willd
14. CINCHONA CALISAYA Wedd
15. CINNAMOMUM CAMPHORA Nees & Eberm
16. CITRULLUS COLOCYNTHIS Schrid
17. CROCUS SATIVUS Linn
18. CUCUMIS TRIGONUS Roxb.
19. CUSCUTA REFLEXA Roxb.
20. DAUCUS CAROTA Linn.
21. DOLICHANDRONE FALCATA Seem.
22. EUPHORBIA TIRUCALLI Linn.
23. EXCOECARIA AGALLOCHA Linn.
24. GARCINIA MORELLA Desr.
25. GLORIOSA SUPERBA Linn.
26. GOSSYPIUM HERBACEUM Linn.
- *27. LEPIDIUM SATIVUM Linn, (S -Chandrasuru, H -Chansaur, B -Halim), cultivated throughout India.
28. MOMORDICA CHARANTIA Linn.
- *29. MOMORDICA TUBEROSA Cogn. (*M. cymbalaria* Fenzl ex Naud), Bo.-Kadavanchi), Western parts of India
30. MORINGA OLEIFERA Lam. (*M. pterygosperma* Geratn.)
31. NERIUM INDICUM Mill. (*N. odorum* Soland.)

- 32 NIGELLA SATIVA Linn.
- 33 PEGANUM HARMALA Linn.
34. PLUMBAGO INDICA Linn (*P rosea* Linn)
35. PLUMBAGO ZEYLANICA Linn
- *36. PLUMERIA RUBRA Linn var *acutifolia* Bailey (*P acutifolia* Poir, *P. acuminata* Ait), (S -Kahira champa, H & Bo -Khair champa), cultivated throughout India, naturalized in many parts
- 37 RANDIA DUMETORUM Lam
38. RUBUS MOLUCCANUS Linn
- 39 RUTA GRAVEOLENS Linn
- 40 SALICORNIA BRACHIATA Roxb
- 41 SAPINDUS TRIFOLIATUS Linn
- 42 SEMECARPUS ANACARDIUM Linn f.
- 43 SESAMUM ORIENTALE Linn (*S. indicum* Linn)
- 44 STACHYTARPHITA JAMAICENSIS (Linn), Vahl var *indica* H J. Lam. (*S. indica* Vahl)
45. TAXUS BACCATA Linn
46. THEVETIA PERUVIANA (Pers) Merr (*T. nervifolia* Juss ex Steud).
47. TRIANTHEMA PENTANDRA Linn
- 48 TRIANTHEMA PORTULACASTRUM Linn (*T monogyna* Linn)
- *49 URENA LOBATA Linn (H -Bachata, B -Bonokra), hotter parts of India, a weed of waste places, forest clearing and roadsides, common in bambo and mango clumps of Bengal
- 50 WITHANIA somnifera Dun

Indian Plants liable to produce Dermatitis

- 1 ABROMA AUGUSTA Linn
- 2 AILANTHUS ALTISSIMA (Mill) Swingle (Syn *A glandulosa* Desf)
- 3 ANACARDIUM OCCIDENTALE Linn.
- 4 ANAGALLIS ARVENSIS Linn.
- 5 ANTHEMIS COTULA Linn.
- 6 APIUM GRAVEOLENS Linn
- 7 ARISAEMA SPECIOSUM (Wall.) Mart.
- *8 ARISAEMA TORTUOSUM (Wall.) Schott , (P.-Don, Nep -Birbanka), from Simla to Sikkim & Bhutan at 8,000 ft , Khasia hills, Manipur, Ranchi, W Ghats
- *9. ASPARAGUS OFFICINALIS Linn., (P , H. & Kash.-Allipalli), from Kashmir to Bhutan, Khasia Hills, Assam.

10. CALOTROPIS GIGANTEA (Linn) Dryand.
11. CALOTROPIS PROCERA (Linn) Dryand.
- 12 CANNABIS SATIVA Linn
- *13. CISSUS SETOSA Roxb , (H -Karmal, Bo -Khaj-goli-cha-vel), Deccan,
W. Ghats of Madras State
- 14 DATURA STRAMONIUM Linn
15. DAUCUS CAROTA Linn
- *16. DELPHINIUM AJACIS Linn., commonly cultivated in gardens.
- *17. DICTAMNUS ALBUS Linn , from Kashmir to Kunawar.
- 18 ERIGERON CANADENSIS Linn
- 19 EUPHORBIA ACAULIS Roxb
20. EUPHORBIA ANTIQUORUM Linn.
- 21 EUPHORBIA CATTIMANDO W Elliot (*E trigona* Fl. Brit. Ind .
in part)
22. EUPHORBIA HELIOSCOPIA Linn
23. EUPHORBIA NERIIFOLIA Linn.
- 24 EUPHORBIA NIVULEA Buch Ham.
25. EUPHORBIA PEPLUS Linn
- 26 EUPHORBIA ROTHIANA Spreng.
- 27 EUPHORBIA ROYLEANA Boiss
- *28 EUPHORBIA THOMSONIANA Boiss , (Kash -Hirtiz), Kashmir 10,000 to
12,000 ft.
29. EUPHORBIA TIRUCALLI Linn
- 30 EUPHORBIA TRIGONA Haw
- 31 EXCOECARIA AGALLOCHA Linn.
- 32 FAGOPYRUM ESCULENTUM Moench.
- 33 FLEURYA INTERRUPTA Gaudich.
- *34. GINKGO BILOBA Linn., (English-Maiden-hair-tree), rarely cultivated
in gardens
- *35 GIRARDINIA HETEROPHYLLA Decne., Mount Abu, Chota Nagpur,
W Peninsula.
36. HEDERA HELIX Linn.
- *37. HIPPOMANE MANCINELLA Linn , (English-Manchineal tree), introduced
into Indian gardens.
38. HOLIGARNA ARNOTTIANA Hook. f.
39. HOLIGARNA GRAHAMII (Wight) Hook. f.

40. *HOLIGARNA LONGIFOLIA* Buch. Ham ex. Roxb.
41. *HUMULUS LUPULUS* Linn
42. *HYPERICUM PERFORATUM* Linn
43. *LAPORTEA CRENULATA* Gaudich
- *44. *LAPORTEA TERMINALIS* Wight , sub-tropical Himalayas from Kumaon to Mishmi at 4,000 to 8,000 ft., Madhya Pradesh, W Ghats of Madras State.
- *45. *LASIOSIPHON ERIOCEPHALUS* Decne , (Bo -Rametha) W Ghats of Bombay and Madras States also Nilgiris
- *46. *LEONURUS CARDIACA* Linn , Kumaon, Kashmir, Hazara, Kurrum valley.
- *47. *LOBELIA EXCELSA* Lesch , W Ghats of S India, Nilgiris, Pulney, Travancore above 6,000 ft
- *48. *LOBELIA NICOTIANIFOLIA* Heyne (B & H -Nala, Bo -Dhavala), W. Ghats from Bombay to Travancore at 3,000 to 7,000 ft., Konkan, Deccan, Nilgiris, Malabar
- *49. *MUCUNA ATROPURPUREA* DC , plains of Western India
- *50. *MUCUNA GIGANTEA* DC , (Malay -Kaku-vali), a littoral species found on the Indian Coast
- *51. *MUCUNA HIRSUTA* Wight & Arn , plains of Western India.
- *52. *MUCUNA MONOSPERMA* DC , (Bo , Sonogaravi, M -Thelu-Kodi), E Himalaya, Khasia Hills, Assam, Chittagong, Konkan, S M Country.
53. *MUCUNA PRURITA* Hook. (*M pruriens* Fl. Brit Ind., non DC.)
54. *NERIUM OLEANDER* Linn.
55. *PODOPHYLLUM HEXANDRUM* Royle (Syn — *P. emodi* Wall. ex. Hook, f & Thoms)
56. *POLYGONUM HYDROPIPER* Linn.
57. *RANUNCULUS SCCELERATUS* Linn.
58. *RHUS INSIGNIS* Hook. f.
59. *RHUS PUNJABENSIS* J. L. Stew ex Brand.
60. *RHUS SUCCEDANEA* Linn.
61. *RHUS WALLICHII* Hook. f.
62. *RUMEX ACETOSA* Linn.
63. *RUMEX ACETOSELLA* Linn
- *64. *RUTA GRAVEOLENS* Linn., var *angustifolia* Hook. f , (S -Somalata, H.-Gadab, B.-Ermul, Bo.-Satap), cultivated in Indian gardens.
65. *SAPIUM INSIGNE* Trimen.

66. SCHIMA WALLICHII Choisy
67. SEMECARPUS ANACARDIUM Linn f
68. SEMECARPUS TRAVANCORICUS Bedd
69. TRAGIA BICOLOR Miq
- 70 TRAGIA INVOLUCRATA Linn
- 71 URTICA DIOICA Linn
72. URTICA HYPERBOREA Jacq.
- 73 URTICA PARVIFLORA Roxb
- 74 URTICA PILULIFERA Linn.
- 75 WALLICHIA DISTICHA T. Anders
76. XANTHIUM STRUMARIUM Linn

Indian Plants considered to have Insecticidal & Piscicidal Properties.

- 1 ACACIA PENNATA (Linn) Willd
2. ACORUS CALAMUS Linn
- *3 ACORUS GRAMINEUS Soland, Sikkim Himalaya upto 6,000 ft , Khasia hills 4,000-5,000 ft
- 4 ACRONYCHIA PEDUNCULATA (Linn) Miq (Syn — *A laurifolia* Blume).
- 5 ADINA CORDIFOLIA (Roxb) Benth & Hook f.
- 6 AGAVA AMERICANA Linn.
- *7 ALBIZZIA CHINENSIS (Osbeck) Merr (Syn - *A stipulata* Boiv), (B.-Amluki, Bo.-Udala), throughout India, ascending to 4,000 ft. in the Himalaya
- 8 ALBIZZIA PROCERA (Roxb.) Benth.
- 9 ANACARDIUM OCCIDENTALE Linn
- 10 ANAGALLIS ARVENSIS Linn
- 11 ANAMIRTA COCCULUS (Linn.) Wight & Arn.
- 12 ANDRACHNE CORDIFOLIA Muell Arg.
- 13 ANNONA RETICULATA Linn.
14. ANNONA SQUAMOSA Linn
- *15. APAMA TOMENTOSA Engl (Syn - *Bragantia tomentosa* Blume); Assam & Manipur.
- 16 ARENGA OBTUSIFOLIA Mart
17. ARISAEMA SPECIOSUM (Wall.) Mart.
18. ARISAEMA TORTUOSUM (Wall.) Schott.

19. ARISTOLOCHIA BRACTEATA Retz.
20. ARTEMISA ABSINTHIUM Linn.
- 21 ARTEMISIA VULGARIS Linn
- 22 ASCLEPIAS CURASSAVICA Linn
- 23 AZADIRACHTA INDICA A. Juss (Syn -*Melia azadirachta* Linn)
- 24 BALANITES AEGYPTIACA Delile (Syn -*B roxburghii* Planch)
- 25 BAMBUSA BAMBOS Druce (Syn -*B. arundinacea* Willd)
- 26 BARRINGTONIA ACUTANGULA (Linn) Gaertn
- 27 BARRINGTONIA ASIATICA (Linn) Kurz Syn *B speciosa* Forst)
- *28. BARRINGTONIA RACEMOSA (Linn) Roxb , (S -Nipr, H -Ijjul), Western sea coast of India, from Konkan to Travancore, Assam, Sunderbans.
29. BERBERIS ARISTATA DC (possibly some other species of Berberis also)
- 30 BUTEA MONOSPERMA (Lam) Kuntze (Syn -*B fiendosa* Koen ex Roxb.)
- 31 CAESALPINIA NUGA (Linn) Ait.
- 32 CALLICARPA LONGIFOLIA Lam. var. *lanicolaria* C B Clarke
33. CALONYCTION MURICATUM (Linn.) G Don (Syn -*Ipomoea muricata* Jacq)
34. CALOPHYLLUM INOPHYLLUM Linn
- 35 CANNABIS SATIVA Linn
- *36. CARDARIA DRABA Desv (Syn -*Lepidium draba* Linn.), (Afgh -Bijindak) a weed of cultivation in the Punjab & N W. Frontier Pakistan
- 37 CAREYA ARBOREA Roxb
38. CASEARIA GRAVEOLENS Dalz
- *39 CASEARIA TOMENTOSA Roxb., (H & Ro -Chillara), throughout India, ascending to 3,000 ft in Himalayas
- 40 CASSYTHA FILIFORMIS Linn
- 41 CENTRATHERUM ANTHELMINTICUM (Willd) Kuntze (Syn.-*Vernonia anthelmintica* Willd)
- 42 CERBERA MANGHAS Linn. (Syn -*C. odollam* Gaertn.)
43. CHRYSANthemum CINERARIAEFOLIUM Vis.
44. CHRYSANthemum COCCINEUM Willd.
- 45 CIMICIFUGA FOETIDA Linn.
46. CINCHONA CALISAYA Wedd.
47. CINCHONA OFFICINALIS Linn.

48. CINCHONA SUCCIRUBRA Pav. ex Klotzsch.
- 49 CINNAMOMUM CAMPHORA Nees & Eberm.
- 50 CLEISTANTHUS COLLINUS Benth. & Hook f
51. CORYPHA UMBRACULIFERA Linn.
- 52 CROTON OBLONGIFOLIUS Roxb.
- 53 CROTON TIGLIUM Linn
- 54 CUCUMIS SATIVUS Linn
- 55 CURCUMA LONGA Roxb
- *56 CYMBOPOGON NARDUS (Linn) Rendle (Syn - *Androp on nardus* Linn)
(Bo -Kamkher, H -Ganjni), throughout the hotter parts, wild or
cultivated
- 57 CYNANCHUM ARNOTTIANUM Wight
- 58 DALBERGIA STIPULACEA Roxb.
- 59 DELPHINIUM BRUNONIANUM Royle
60. DELPHINIUM CAERULEUM Jacquem ex Cambess
- 61 DELPHINIUM ELATUM Linn.
- 62 DERRIS ELLIPTICA (Roxb) Benth
- 63 DERRIS FERRUGINEA (Roxb.) Benth
- 64 DERRIS SCANDENS (Roxb.) Benth
- *65 DERRIS TRIFOLIATA Lour , var *uliginosa* (Roxb ex Willd) (Syn -
D uliginosa Benth , *Robinia uliginosa* Roxb ex Willd)
- 66 DIOSCOREA HISPIDA Dennst (Syn -*D. daemonia* Roxb)
- 67 DIOSCOREA PRAZERI Prain, Burkill (Syn -*D deltoidea* Wall , var
sikkimensis Prain)
- 68 DIOSPYROS EBENUM Koen
- *69 DIOSPYROS MONTANA Roxb , (S -Tumala, H -Lohori, B -Bangab),
throughout India, from Kanara eastwards in the sub-Himalayan
tract
- *70 DIOSPYROS PANICULATA Dalz , (S -Thinduka, M -Karinthuvari),
S M Country, Travancore, Kanara, Malabar
- 71 DODONAEA VISCOSA (Linn) Jacq
- 72 DOLICHANDRONE FALCATA Seem
- 73 DURANTA REPENS Linn. (Syn.-*D plumieri* Jacq)
74. EDGEWORTHIA GARDNERI Meissn
75. ENTADA PURSAETHA DC (Syn -*E. scandens* Benth)
- *76 EREMOSTACHYS SUPERBA Royle ex Benth., W. Himalaya, Punjab,
Garhwal.

77. EREMOSTACHYS VICARYI Benth
78. EUCALYPTUS GLOBULUS Labill
- 79 EUPATORIUM ODORATUM Linn
- 80 EUPHORBIA ANTIQUORUM Linn
- 81 EUPHORBIA NERIIFOLIA Linn
82. EUPHORBIA ROYLEANA Boiss
83. EUPHORBIA THYMIFOLIA Linn.
84. EUPHORBIA TIRUCALLI Linn
85. EXCOECARIA AGALLOCHA Linn
86. FLUGGEA LEUCOPHYRUS (Koen) Willd
- 87 FLUGGEA VIROSA (Roxb. ex Willd) Baill. (Syn -*F. microcarpa* Blume)
- *88. GARDENIA CAMPANULATA Roxb , (Burm -Hsathanpaya), foot of Sikkim Himalaya, Assam, Sylhet, Chittagong, Bihar.
- 89 GAULTHERIA FRAGRANTISSIMA Wall
- 90 GLORIOSA SUPERBA Linn
91. GNETUM SCANDENS Roxb.
- 92 GYNANDROPSIS GYNANDRA (Linn) Merr. (Syn.-*G pentaphylla* DC.)
- 93 GYNOCARDIA ODORATA R Br
- 94 HARPULLIA CUPANIOIDES Roxb.
- 95 HEDERA HELIX Linn
96. HYDNOCARPUS KURZII (King) Warb (Syn -*Taraktongenos kurzii* King)
97. HYDNOCARPUS LAURIFOLIA (Dennst) Sleumer (Syn -*H wightiana* Blume)
- 98 HYDROCOTYLE JAVANICA Thunb
- 99 JATROPHA CURCAS Linn
- 100 JUGLANS REGIA Linn
- 101 KALANCHOE SPATHULATA (Poir) DC.
- *102 LAGENANDRA TOXICARIA Dalz (Many authors view this species as synonymous with *L ovata* (Linn) Thw of Ceylon), (Bo Rakhalu, M -Maravara, Tsjmul), Konkan to N. Kanara, Travancore, Mysore, Coorg, Cochin
- 103 LASIOSIPHON ERIOCEPHALUS Dcne
- *104. LINOSTOMA DECANDRUM Wall , Sylhet, Chittagong
105. MADHUCA LATIFOLIA (Roxb) Macbride (Syn -*Bassia latifolia* Roxb)

- 106 MADHUCA LONGIFOLIA (Linn.) Macbride (Syn.—*Bassia longifolia* Linn)
107. MAESA INDICA Wall
- 108 MELALEUCA LEUCADENDRON Linn.
- 109 MELODINUM MONOGYNUS Roxb
110. MILLETTIA AURICULATA Baker ex Brand
- *111 MILLETTIA PACHYCARPA Benth , forests of Garo & Khasia Hills, Sikkim & Assam upto 4,000 ft
- *112 MILLETTIA PISCIDIA Wight & Arn , Sikkim and Assam
- 113 MUNDULEA SERICEA (Willd) Greenway (Syn —*M suberosa* Benth.)
- 114 MYRICA NAGI Thunb.
- 115 NICANDRA PHYSALOIDES Gaertn
- 116 NICOTIANA RUSTICA Linn.
- 117 NICOTIANA TABACUM Linn
118. NIGELLA SATIVA Linn
- *119 OCIMUM GRATISSIMUM Linn , (S -Vantulshi, H & B -Ramtulshi); common wild plant in W India cultivated in gardens in Bengal, Nepal, Deccan Peninsula
- 120 OUGEINIA DALBERGIOIDES Benth
- 121 PACHYGONE OVATA (Poir) Miers ex Hook f & Thoms.
122. PEGANUM HARMALA Linn
- 123 PHYLLANTHUS URINARIA Linn
- 124 PICRASMA JAVANICA Blume, var *nepalensis* Badhwar nov comb. (Syn -*P nepalensis* Benn).
125. PIERIS OVALIFOLIA D Don
126. PITHECELLOBIUM BIGEMINUM Mart (Syn —*Pithecolobium bigeminum* Benth).
127. POGOSTEMON HEYNEANUS Benth (Syn -*P patchouli* Hook. f. in Fl. Brit -Ind , non. Pellet)
- 128 POLYGONUM FLACCIDUM Meissn
- 129 POLYGONUM HYDROPIPER Linn
- 130 PONGAMIA PINNATA Linn Merr (Syn.-*P glabra* Vent)
- 131 PYGEUM GARDNERI Hook. f., (Bo -Daka), W Ghats of Bombay & Madras States, hills of Travancore, Malabar, Nilgiris, Pulneys the Deccan.
- 132 RANDIA DUMETORUM Lam. (Split up into three species by Gamble)
133. RANDIA ULIGINOSA DC.

- 134. RAUWOLFIA SERPENTINA Benth ex Kurz
- 135 RHODODENDRON BARBATUM Wall ex G Don , (Nepl.-Guras), Kumaon to Bhutan at 8,000 to 12,000 ft., Sikkim
- 136 RHODODENDRON FALCONERI Hook f , (Nepl -Kurling), East Nepal to Bhutan at 9,000 13,0000 ft
- 137 RICINUS COMMUNIS Linn.
- 138 RUTA GRAVEOLENS Linn. var *angustifolia* Hook f
- 139 SANTALUM ALBUM Linn
- 140 SAPINDUS MUKOROSI Gaertn , (S -Phenila, H , B & Bo -Ritha) cultivated throughout N W India, Bengal & Assam, wild on the Himalayas upto 4,000 ft
- 141 SAPINDUS TRIFOLIATUS Linn
- 142 SAPIUM INDICUM Willd
- 143. SARCOSTEMMA ACIDUM (Roxb) Voigt (Syn -*S brevistigma* Wight & Arn)
- 144 SAUSSUREA LAPPA C. B Clarke
- 145 SCHLEICHERA OLEOSA (Lour) Merr (Syn -*S trijuga* Willd)
- 146 SCLERIA PERGRACILIS (Nees) Kunth widely scattered from Garhwal at 5,000 ft to Assam, Bihar, Chota Nagpur & Deccan
- 147 SOPHORA MOLLIS R Grah , (P -Buna, Arab -Arghavan), plains and low hills of the North-West, Hazara & the Salt Range to Kumaon & Nepal, upto 7,000 ft., & Bushahr & near Dehra Dun
- 148 SPHAERANTHUS INDICUS Linn
- 149 STEPHANIA HERNANDIIFOLIA (Willd) Walp , (S -Vanatik-tika, B -Agnad), W & E. Coast, Dehra Dun, Biharam (Sikkim, Bengal, Assam, Cachar)
- 150. STRYCHNOS COLUBRINA Linn
- 151 STRYCHNOS NUX-VOMICA Linn
- 152. TAXUS BACCATA Linn.
- 153 TEPHROSIA CANDIDA (Roxb) DC
- 154 TEPHROSIA PURPUREA (Linn) Pers
- *155 TEPHROSIA VOGELII Hook f , cultivated by tea planters in Assam
- 156 TERMINALIA BELLERICA (Gaertn) Roxb.
- 157. THEVETIA PERUVIANA (Pers) Merr (Syn -*T nereifolia* Juss ex. Steud)
- *158 TRIGONELLA FOENUM-GRÆCUM Linn , (H B & Bo -Methi), Punjab & Kashmir, cultivated in many parts of India

159. VERBASCUM THAPSUS Linn
- *160 VITEX NEGUNDO Linn , (S. & H -Nirgundhi, B -Nishinda), throughout India in the warmer zone
- 161 WALSURA PISCIDIA Roxb
- 162 WIKSTROEMIA INDICA (Linn) C. A Mey var. *viridiflora* (Meissn) Hook f.
163. ZANTHOXYLUM ALATUM Roxb
- *164 ZANTHOXYLUM HAMILEONIANUM Wall , (Nep -Purpuray timur), throughout Assam, in low level forests, except in the Khasia and Jaintia hills

CHAPTER X

B i b l i o g r a p h y

In the first edition of The Review, a list of the papers published on the subject under the Indian Council of Medical Research (formerly Indian Research Fund Association) was given. As the scope of the present review has been widened to include all the work done on Indian Indigenous Drugs anywhere, the question of including a comprehensive and if possible a complete bibliography for the new edition was discussed and agreed to by the Secretary Indian Council of Medical Research. In view of the fact that no such bibliography has so far been published and also because this subject has been assuming increased importance both in India and abroad, it was decided that a comprehensive bibliography should form a particular feature of the present edition, inspite of the fact that size of volume would be considerably increased.

During the last 3 or 4 decades the subject of Indian Indigenous Drugs has received considerable attention of many of the scientific workers in this country and their papers have been published in various journals in India and elsewhere. The subject has interested many foreign workers also who have made very important contributions in foreign scientific journals. An attempt has been made to get all the references together, particularly of the work done on the plants of Indian origin. This is a special feature of the bibliography which is given in this Chapter.

To facilitate consultation by workers references have been arranged plant-wise (which are given in alphabetical order,) and in sequence of years in which the papers have been published. In order to attract the immediate attention of the workers the un-orthodox method of giving the title of the paper first and names of authors afterwards has been adopted.

Abbreviations of the journals used are also given in a separate list.

The inclusion of all the available bibliography in the present volume has involved considerable labour and careful sifting. It is hoped that it will be useful to those who wish to consult the previous work done when taking up a plant for investigation. It is by no means such an exhaustive bibliography as we would have wished but it is hoped that it may serve as a first compilation of its kind which can be gradually added to.

List of Journals and Books with the Abbreviations.

Agri Jr Ind	.	Agricultural Journal of India
Agri Ledger	..	Agricultural Ledger (Govt of India)
Amer Chem Jr	..	American Chemical Journal
Amer Jr. Pharm	...	American Journal of Pharmacy.
Analyst	...	Analyst.
Ann Bot.	...	Annals of Botany.

Ann. Chem.	...	Annalen der Chemie.
Ann Chim	...	Annales de Chemie
Ann Chim Phys	...	Annales de Chimie et de Physique
Ann Chim Appl	...	Annales de Chimie analytique appliquee
Ann. Inst Past	..	Annales de l' Institut Pasteur
Arch de Pharm.	..	Archiv de Pharmacie
Arch Exper Path Pharm		Archiv fur experimentelle Pathologie and Pharmakologie
Arch Hyg.	...	Archiv fur Hygiene
Arch Pharm	.	Archiv der Pharmazie
Biochem d Pflzen	...	Biochemie der Pflanzen
Biochem Jr	.	Biochemical Journal
Biochem Ztsche		Biochemische Zeitschrift.
B M J	...	British Medical Journal
Buchn Repert Pharm	...	Repertorium der Pharmacie (Buchner)
Bull Acad Sci Allahabad		Bulletin of the Academy of Science of Uttar Pradesh of Agra & Oudh, Allahabad
Bull Bot Gard Kew.	...	Bulletin Botanical Garden Kew
Bull Sc Pharm.	...	Bulletin des Sciences Pharmacologiques
Bull. Soc Chim	...	Bulletin de la Societe Chimique de France
Bull Soc Chim Biol	...	Bulletin de la Societe de Chimie Bio- logique
Bull Tokyo Inst Technical		Bulletin, Tokyo Institute of Techno- logy
C C	...	Chemisches Centralblatt
Chem Abst	..	Chemical Abstract (American)
Chem Weekbl.	...	Chemisches Weekblad
Chem Ztg	...	Chemiker-Zeitung
Chin Jr Physiol	..	Chinese Journal of Physiology
Compt Rend	...	Comptes rendus hebd des Seances de l' Academie des Sciences
Comp Soc Biol	...	Comptes rendus hebd des Seances de la Societe de Biologie.
Curr Sci	...	Current Science
Helv. Chim Act.	...	Helvetica Chimica Acta
Ind Farming	...	Indian Farming
Ind. For Rec.	...	Indian Forest Records (and Bulletin) Dehra Dun
Ind. For.	...	Indian Forester
Ind. Jr. Agric Sci	...	Indian Journal of Agricultural Science

Ind. Jr. Ent.	...	Indian Journal of Entomology.
Ind. Jr. Med. & Phys Sc.		Indian Journal of Medical and Physical Science
I. J. M. R.	...	Indian Journal of Medical Research.
Ind. Jr. Pharm.	...	Indian Journal of Pharmacy
Ind Jr. Vetr Sci Animal Husbandry	...	Indian Journal of Veterinary Science & Animal Husbandry
I. M. G	...	Indian Medical Gazette.
I M J.	...	Indian Medical Journal
J C. S.	...	Journal of the Chemical Society, London.
Jr. Agri Res	...	Journal of Agricultural Research
Jahrb. Prakt Pharm	...	Jahrbuch fur Praktische Pharmazie
Jr. Amer Pharm. Assoc	...	Journal of the American Pharmaceutical Association
Jr Amer Chem Soc		Journal of the American Chemical Society
Jr Anamalai Univ		Journal of Anamalai University
Jr Assoc Agri Chem		Journal of the Association of Official Agricultural Chemists
Jr. Biol Chem	...	Journal of Biological Chemistry
Jr Bombay Nat Hist Soc		Journal of Bombay Natural History Society
Jr Pharm	.	Journal der Pharmacie fur Aerzte
Jr Ind Chem Soc	..	Journal of the Indian Chemical Society
Jr Ind Inst Sci	..	Journal of the Indian Institute of Science, Bangalore
Jr Malaria Inst India	.	Journal of the Malaria Institute of India
Jr Mysore Univ		Journal of the Mysore University
Jr Pharm Chim		Journal de Pharmacie et de Chimie
Jr Pharm Soc Japan		Journal of the Pharmaceutical Society of Japan.
Jr Pharm Exp Therap	.	Journal of Pharmacology and Experimental Therapeutics.
Jr Physiol.	...	Journal of Physiology.
Jr Pract Chem	...	Journal fur Praktisch Chemie.
Jr Russ Phys Chem Ges.		Journal der Russischen Physikalisch-chemischen Gesellschaft.
Ji Sci Indust Res	...	Journal of Scientific & Industrial Research
Jr Soc Chem Ind	..	Journal of the Society of Chemical Industry.

Chemical examination of the seeds of *Abrus precatorius* L.
Constitution of abrine, N Ghatak, *Bull. Acad. Sci United Provinces
Agra, Oudh, Allahabad*, 295, 1934

Chemistry and toxicology of *Abrus precatorius* or seed or Jequirity.,
Carlos L. Carbone Schi. *Scmana Med (Buenos Aircs)* 1947, 275.

Some components of the seeds *Abrus precatorius* Akira Yokoo,
(Tokyo Inst. Technol) *Bull. Tokyo Inst Technol.*, 13 43 (1948)

Chemical examination of the fixed oil from the seeds of *Abrus pre-*
catorius, Moolraj Mandiratta and S Dutta; *Ind Soap Jour* , 1949,
195

ABUTILON AVICENNAE Gaertn

The oil from the seeds *Abutilon avicennae* Gaertn G R. Stepanov.,
Trudin Inst Novogo, Lulyanogo Surya, 123 1933, *Chem Abst*
1933, 4945

INDICUM G Don

Cambir, Chem. Exam of seeds of *Abutilon indicum*, G. Don Indr
Raj & Joshi, Sham Sunder , *J Ind Chem. Soc.* 29 (1952), 451

ACACIA Willd

Nierenstein, M -The catechins of the Cutch-producing Acacias.,
J. Ind Chem Soc 7 (1930), 279

Indian Wattle Bark *Jr. Sci Indl Res* , 1947, 6A, 447.

ACACIA ARABICA Willd

The tannin content of *Acacia arabica* pods, *Anon Bull Imp Inst.*
28, 1 1930

Gum arabic, influence of heat on the chemical composition and phys-
ical properties Moorjani & Narwani, -*Jr. Ind Chem Soc* 25
(1948), 502

CATECHU Willd

Constitution of Catechol M. Nierenstein, *J Am Chem. Soc* 52,
1672, 1930

L Epi-Catechin from *Acacia catechu* Rao, P R and Sheshadri,
T R , *J Sci Indl Res* , 1948, 7B, 59

ACACIA CHUNDRA (Willd) D. C

Acacia chundra or *Acacia Sundra* Chatterjee, D., *Sci & Cult.*,
1949, 14, 290

LEUCOPHLOEA Willd

Chemical examination of flowers of *Acacia leucophloea* Willd.
Mukherjee, S. K. and Murty, V. V. S , *Jr Sci. Indl. Res.*, 1952,
11B, 125.

ACALYPHA Linn.

The correlation between chemical composition of Anthelmintics and
their therapeutic values in connection with the Hookworm Inquiry
in the Madras Presidency, XVIII, Catherlics, *Oleum ricini*, *Oleum*
tiglii, *Aloe*, *Succies acalyphae*, Caines, J. F. and Mhaskar, K. S.,
Ind. Jour. Med. Res., 1923, 11, 103.

A INDICA Linn.

Chemical investigation of the plant *Acalypha indica* Isolation of triacetoneamine, a cyanogenetic glucoside and quebrachitol. Claude Remington and G C S Roet. Onderstepoort, *J Vet. Sci. Animal Ind.* 9, 193 1937 *Chem Abst* 4629, 1938

The presence of Hydrocyanic acid in stock feeds and other plants. *Acalypha indica* L Donw O Steyn, *J S African Vet Med. Assoc* 9, 60, 1938 *Chem Abst* 8617, 1938

ACHILLEA MILLEFOLIUM Linn

Oil of *Achillea millefolium*, L R E Kremens, *J Am Pharm Assoc* 14, 399 1925

The ethereal oil content of fragrant plants of the southern territory. *Achillea millefolium*, O Sobolevskii, *Chem Abst* 3017, 1928.

The non pre-existence of azulene in milfoil Katherine Graham, *J Am Pharm Assoc* 22, 819 1933

Achillea millefolium Linn constituents of the petroleum ether extract of the blossoms R L Macmurray, *Am. J Pharm* 105, 573, 1933

Phytochemical notes The sterols of *Achillea millefolium*. Ole. Gisvold, *J Am Pharm Assoc* 24, 107, 1953

Volatile oil from Western yarrow *Achillea millefolium*, lanulosa R L, Macmurray, *J. Am Pharm Assoc* 25, 304, 1936

Investigation of the antipyretic action of native medicinal plants. *Achillea millefolium* Maksym Naksyn Nikonorow, *Acta Polon, Pharm* 3, 23, 1939 *Chem Abst.* 3766, 1941

ACHRAS SAPOTA Linn

Achras sapota, fatty acids & glycerides of the oil from sapota seeds Vidyarthi, N L & Mallya, M Venkatesh; *J Ind Chem Soc.* 16 (1939), 443.

ACHYRANTHES ASPERA Linn

Chemical examination of the seeds of *Achyranthes aspera* Linn Gopalachari, R & Dhar M L, *J. Sci Indl Res*, 1952, **11B**, 209

ACONITUM Linn

The Indian varieties of Aconite Their chemical composition and biological assay Chopra, R N, Gupta, J C & Ghosh, N. N; *Ind. Jour Med Res*, 1928, 15, 873.

Aconite alkaloids Action of methyl alcoholic sodium hydroxide on atisine, iso-atisine and dihydroatisine-Walter, A Jacobs and Hayman E Craig, *J. Biol. Chem.* 147, 567, 1943

A. ANTHORA Linn

Atisine from *Aconitum heterophyllum* Wall, and anathorine from *Aconitum anthora*, L. Andoregoris, *Comp Rend* 205, 1007, 1937

BALFOURII Stapf

The alkaloids of some Indian Aconites (*A. balfourii*, *A. deinorrhizum* and Chumbi Aconite). Thomas Anderson Henry and Thomas Marvel Sharp, *J Chem Soc* 1105, 1928

CHASMANTHUM Stapf, ex Holmes.

Aconitine from Cashmerian *Aconitum chasmanthum* K H Bayer and Tara Chand Razdan, *Pharm Zentralhalle*, 72, 146, 1931

. DEINORRHIZUM Stapf

The alkaloids of some Indian Aconities (*A. balfourii*, *A. deinorrhizum* and Chumbi Aconite) Thomas Anderson Henry and Thomas Marvel Sharp, *J Chem. Soc* 1105, 1928

HETEROPHYLLUM Wall.

Atisine from *Aconitum heterophyllum* Wall and anthonine from *Aconitum anthora* L Andoregoris, *Comp Rend* 205, 1007, 1937

. NAEPELLUS Linn

New alkaloid from *Aconitum napellus* Henrich Schulz and Gottfried Berger, *Arch. Pharm* 262, 556, 1924

Aconite alkaloids New alkaloids in *Aconitum napellus* Werner Fredenberg and E F Rogers, *J. Am Chem Soc* 59, 2572, 1937

SOONGARICUM

Alkaloids of *Aconitum soongaricum* Alkaloids of Ranunculaceae. S Yunusov, *J Gen Chem (U S S. R)* 18, 515, 1948, *Chem Abst* 7940, 1948

CORUS CALAMUS Linn

Notes on some Indian Essential Oils *Acorus calamus*, Linn, (Calamus Root) Rao B Sanjiva, Sudborough J J, and Watson H E, *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol 8A, Part X, pp 149-151

Essential oil from the rhizomes of *Acorus calamus*, N C Kelkar & B Sanjiva Rao, *J. Ind Inst Sci* 17, 24 1933

Studies in Indian Essential Oils VI—Essential Oil from the rhizomes of *Acorus calamus* Linn Kelker N C and Rao B Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1933, Vol 17A, Part II, pp. 25-31

A. GRAMINEUS Soland.

Constituents of the essential oil *Acorus gramineus*, Soland, Y. Kimura, *J. Pharm Soc Japan* No 431, 380, (1926)

ACTAEA SPICATA Linn

Tinns, *Actaea spicata* Halmut Thaler Mikrokosmos; 22, 94, 1929, *Chem Abst* 2596, 1929

ACTINODAPHNE Nees

Actinodaphne and Litsea fats as raw material for a valuable new detergent, S. V. Puntambekar: *Ind Forester* 60, 707, 1934.

A HOOKERI Meissn.

Actinodaphne hookeri Meissn; Actinodaphine, an alkaloid from. Krishna, S. & Ghosh T. P, *J Ind Chem Soc.* 9 (1932), 429.

Actinodaphne hookeri Meissn. The fat and oil from the seeds of An indigenous source of Lauric acid Puntambekar, S. V & S Krishna, *J Ind Chem Soc* 10 (1933), 395.

ADANSONIA Linn

Some new data on the pharmacology of Adansonia leaves. Daniel Incent, *Chem Abst* 1698, 1951

ADENANTHERA PAVONINA Linn

Oil from the seeds of Adenanthera pavonina A source of lignoceric acid, S M Mubidri, P. Ramaswami Ayyar and H E Weston, *J Ind. Inst Sci* 11 A, 173, 1928

Chemical examination of the leaves of Adenanthera pavonina Linn Patel, C S, Shah C C, & Parikh, H P, *Curr Sci*, 1947, 16, 344

ADHATODA VASICA Nees

A preliminary note on the pharmacology and therapeutics of Adhatoda vasica (Basak) Chopra, R N & Ghosh, Sudhamoy, *Ind. Med. Gaz*, 1925, 60, 354.

Some observations on the Pharmacological Action and Therapeutic properties of Adhatoda vasica (Basak) Chopra, R N, & Ghosh, S, *Ind Jour. Med Res*, 1925, 13, 205

Vasicine-an alkaloid present in Adhatoda vasica Nees Part II Ghish, Tarak Prasad, *J Ind Chem Soc* 4 (1927), 1

Report of the Imperial Agriculture Chemist, Ash of the Adhatoda vasica Plant, B Viswanath, *Sci. Reports Imp Inst Agr. Research, Pusa*, 1934-35, 103, 1936

A pharmacological study of stem and leaf of Adhatoda vasica Nees. Prasad, S & Prabhu, P N, *Ind Jour Pharm*, 1950, 12, 200

ADINA CORDIFOLIA Hk f

Adina cordifolia Hook A yellow colouring matter from the wood of Jagraj Behari Lal & Dutt, Sikkibhushan, *J Ind Chem Soc* 12 (1935) 257

Fish poison from the wood of Adina cordifolia Rao, N P., *J Sci. Indl Res*, 1949, 8B, 95

AEGLE MARMELOS Corr

Preliminary chemical examination of Aegle marmelos. Dikshit, Brij Behari Lal, & Dutt, Sikkibhushan *J Ind Chem Soc* 7(1930), 756

On the active principles of the bark of Aegle marmelos Correa Mookerji, Asima (Miss), *Curr. Sci*, 1943, 12, 209

On the alkaloidal isolation from the matured bark of Aegle marmelos. Chakravarty, K. K., *J Ind Chem Soc* 21 (1944), 401

Constituents of the matured bark of Aegle marmelos Correa. Asima Chatterjee and Sudhangsu Sekhar Mitra; *J. Am. Chem. Soc*, 71, 606, 1949.

Aegle marmelos, essential oil from leaves of Baslas & Deshpande, *J Ind Chem Soc* , 26 (1949), 23

L-d-Phellandrene from the essential oil from leaves of. Baslas & Deshpande; *J Ind Chem. Soc.* 26 (1949) 231

Aegle marmelos, Essential oils from the leaves, twigs & fruits of Baslas & Deshpande. *J. Ind Chem Soc* 23 (1951), 19.

Studies on the active principles isolated from the leaves of *Aegle marmelos* Chatterjee Asima (Nee Mukerjee, & Bose Sukumar, *J Ind Chem. Soc* 29 (1952), 425

AEROTHECIUM LANATUM Wakker

A preliminary note on the isolation and pathogenicity of *Aerothecium lanatum* Wakker and *Helminthosporium tetramera* McKinney on the leaves of *Oryza sativa* Linn Ganguly, A K., Ganguly, D C, *Sci & Cult* , 1941, 6, 424,

AESCULUS Linn

The physiological significance of the glucosides in *Aesculus* and *Salix*, Gerhard Kerston, *Chem Abst* 6772, 1934

A HIPPOCASTANUM Linn

The hydrolysis of ptytic compounds derived from seeds of hemp horse bean horse chestnut and flax, wheat and embryos of rye W, Jarosza, *Chem Abst* 5501, 1934.

Production of alcohol from horse chestnuts, Zabrodskii, golde berg Fuks and Kogan *Chem Abst* 7415, 1934

Plant pigments, the occurrence of L and B carotene in different natural products, horse chestnut P Karyer and W Schlientz Helv, *Chem Acta* 17, 7, 1934

Inhibiting effect of Pilocarpine on intestine treated with extract of horse chestnut. H Busquet Comp, Redn Soc. Biol 118, 234, 1935 *Chem Abst* 3035, 1935.

Chemistry of *Aesculus* saponin and its non sugar components E Bures and K Babor, *Chem Abst* 5454, 1935

Use of *Aesculus hippocastanum* (horse chestnut) as a raw material for distilleries, B Lampe and R deplanque, *Chem Abst.* 4616, 1936.

Electric treatment of horse chestnuts and acorns. Hayao Muraoka; *Chem Abst* 3268, 1936

Saponin content and varying saponin content of certain plants and seeds *Aesculus hippocastanum* L M Roberg and E Marchal Jahrb, Wiss Botan. 84, 710, 1937 *Chem. Abst.* 177, 1938.

Saponins from domestic (German) plants *Aesculus hippocastanum* E Wagner, *Chem Abst* 3032, 1941.

AGAVE AMERICANA Linn

Saponins A new method for preparing the foam producing principle of *Agave americana*, L. J. Balansard and P. Flandrin; *Bull. Soc. Chem Biol.* 27. 618, 1945.

AGERATUM CONYZOIDES Linn.

Travancore Essential oils Essential oil from *Ageratum conyzoides* (Appa grass). Moudgill K L, *Journal of the Indian Chemical Society, Calcutta*, 1925, Vol I

Perfumed oils of French upper Ubangi, *Ageratum conyzoides* L L Jolly, *Chem Abst* 2750, 1637

ALLANTHUS GLANDULOSA Desf

Book of *Ailanthus glandulosa* Desf and its bitter principle ailanthin R Wasichy and S Oerik, *Chem Abst* 255, 1934

Ailanthus wood as paper making material L Vidal and M Aribert, *Paper Trade Jour* 85, 7, 49, 1927

Tests carried out at the French School of paper making, qualities of *Ailanthus* wood (*Ailanthus glandulosa*) L Vidal and M Aribert *Papier*, 30, 183 1927 *Chem Abst* 2062, 1927

Essential oils of the flora of Tadshikistan, *Ailanthus glandulosa*, V Isaev; *Chem Abst* 3180, 1934

Ailanthus glandulosa, Anna M Bernasconi, *Chem Abst* 8473, 1938

A. MALABARICA D. C

Chemical Examination of *Ailanthus malabarica* DC, Rastogi, R P Sharma, V N & Dhar, M L., *J Sci Indl Res*, 1952, 11B, 124

ALANGIUM BEGONIAEFOLIA Roxb

A note on the development on the Embryo sac and Endosperm in *Alangium begoniaefolia*, Roxb (*Marlea begoniaefolia*, Roxb) Mitra, J N & Datta, R M., *Sci & Cult.*, 1947, 12, 451

A. LAMARCKII Thw.

Alangium lamarckii Its chemistry and pharmacological action Chopra, R N & Chowhan J S, *Ind Jour Med Res*, 1934, 21, 507.

Chemical examination of the seeds of *Alangium lamarckii*, A Lakshminarasimhaiah, B L Manjunath and B. S. Nagaraj, *J. Mysore Uni* B 3, 113, 1942

Chemical examination of the seeds of *Alangium lamarckii*, isolation of alangol Prithvi Nath Bhargava and Shakhibhushan, Dutt *Proc. Ind. Acad. Sci.* 16A, 328, 1942

Chemical investigation of *Alangium lamarckii* Thw. Basu, N K., Nair, N S, & Bhattacharya, N N *Ind. Jour Pharm*, 1950, 12, 98

ALLIUM CEPA Linn.

Examination of some Hungarian onions A modified method for crude fiber. Zoltan sandor., *Chem Abst* 1788, 1934.

Carbohydrates of the bulbs of *Allium odorum* and *Allium cepa*, Yoshihiro Kihara. *J Agr. Chem. Soc Japan*, 11, 548, 1935

Food values and contents of Vitamin C of Hungarian, Holland and Spanish onions. Zoltan Sandor, *Chem. Abst.* 1896, 1935.

Essential oil of onion plants *Allium cepa* establishments; *Antonic Chiris, Perfumes France* 15, 219, 1937.

Pharmacological trials of some domestic plants *Allium cepa* H. Kreitmair, *Chem Abst* 3149, 1937

The antimycolic effect of ethereal oils of the leek and onion family of plants on pathogenic fungi P W Schmidt and V Marquardt, *Chem Abst* 2642, 1937.

Vitamin C content of chillies, onion & garlic, both in raw-state and when boiled with water Biswas, H G, & Das, K L., *Ind Jour Med Res*, 1939, 27, 139.

Onion juice and bacterial growth. James E Fuller and Ernest R Higgins, *Food Research* 5, 503, 1940.

The significance of nucleoelastic enzymes in the phenomenon of bacteriology, the working hypothesis and its basis Bacteriolytic agents of onion juice Ph L Spainer, E I Chartkova, Sh Gering, *Chem Abst.* 2552, 1941.

Antibacterial properties of yeasts, *Fusarium* species, onion and garlic C W Carpenter, *Chem Abst*, 1558, 1946

A study of the oils from the seeds of *Luffa aegyptica*, *Benincasa cerifera* and *Allium cepa*, K D Phadnis, A V. Rege, D. C Pishawikar and S V. Shah., *J. Univ Bombay*, 17A, 62, 1948

Carbohydrates of Garlic (*Allium sativum*) and Onion (*A. Cepa* L. Srinivasan, M, Bhatia, I S, Satya Narayana, M N, *Curr Sci* 1953, 7, 203

A MINUS

The inactivation of antibacterial agents and their mechanism of action *Allium sativum* and A minus C. J Cavallito, John Haya Bailey, T H. Haskell, J. R Maccromick and W F Warner, *J Bact* 50, 61, 1945

A ODORUM Linn

Carbohydrates of the bulbs of *Allium odorum* and *Allium cepa* Yoshihiro Kihara, *J Agr Chem Soc, Japan*, 11, 548, 1935

A PORRUM Linn.

The antimycolic effect of ethereal oils of the leek and onion family of plants on pathogenic fungi. P W Schmidt and V. marquardt *Chem. Abst.* 2642, 1937

A. SATIVUM Linn.

Bactericidal properties of acrolein (present in garlic) Richard E. Vollrath, Lucile Walton and Carl C Lindegrem, *Proc. Soc Acad Expt Biol Med* 36, 55, 1937

Vitamin C content of chillies, onion & garlic, both in raw state and when boiled with water, Biswas, H G., & Das, K. L., *Ind. Jour Med. Res.* 1939, 27, 139

Antibacterial principle of *Allium sativum*. its precursor and essential oil of garlic Chester J. Cavallito, John H. Bailey and Johannes S. Buck, *J Am Chem Soc* 67, 1032, 1945

Investigation on Plant antibiotics Pt I Studies on Allicin, the antibacterial principle of *Allium sativum* (Garbi), Raghunandana Rao, R. Srinivasa Rao, S. & Venkataraman, P. R., *J Sci Indl Res*, 1946, 5, 31

Oxidase of garlic, J. Sugihara & W. V. Cruess, *Chem Abst*, 6508, 1946

On the effect of garlic on bacteria, Omer Ozek, *Chem Abst*, 6547, 1946

Pharmacological investigation of *Allium sativum* general action Action on arterial pressure and respiration Giuseppe Santelippo and Giuliano OHaviono, *Chem Absta* 7384, 1946

Allin the pure mother substance of garlic oil A. Stoll and E. Scelebeck, *Experimentia*, 3, 114, 1947

Experimental studies of the Pharmacology of the active principles of *Allium sativum* (garlic), Enrique Umberto de Torres Casana, *Chem Abst* 2172, 1947

Antibiotic principles of *Allium sativum* (Lahsan), Datta, N. L., Krishnamurti A. & Siddiqui, *S J Sci Indl Res*, 1948, **7B**, 42

Allicin the active principle of *Allium sativum*, isolation, physical properties and antibacterial action, Chester J., Cavallito and John, H. Bailey, *J Am Chem. Soc* 66, 1950

A preliminary report on 'Loop Top' of Garlic Mahmud, K. A., *Sci & Cult*, 1952, 71, 296

ALOE Tourn. ex Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency, XVIII Catherlics Oleum ricini, Oleum tiglii, Aloe, Succies acalyphae, Caines, J. F., Mhaskar, K. S., *Ind Jour Med Res*, 1923, 11, 103

Areca nuts and aloes H. Octtel Apoth, *Ztg* 48, 1457, 1933

Chemical examination of Indian Aloes R. N. Chopra and N. N. Ghosh, *Arch Pharm*, 276, 348, 1938

Further studies on Cathartic action in mice, Senna, Aloe Casara and bile salts, Lloyd W. Hazleton and Kathleen D. Talbert, *J Am Pharm Assoc* 33, 170, 1944

Indian Aloes, Mukerji, B., & Karkum, J. N., *Ind Jour Pharm* 1950, 12, 236

A. ABYSSINICA Lam

A chemistry of Ethiopian Aloe and possibility of use of *A. abyssinica* Lan. arck, E. Baccari, *Chem Abst* 4547, 194

A VERA Linn

Indian Medicinal Plants Studies in the specifications of Aloe vera Linn. Sen Gupta, S S , & Gupta, H N , *Ind Jour Pharm* , 1949, 11, 27.

ALSOPHILA GLABRA Hook

Spore germination in Alsophila glabra Hook Kachroo, P , *Sci & Cult* , 1952, 17, 342

Alsophila glabra (B₁) Hook from Sylhet Islam, A S , *Sci & Cult* , 1952, 17, 473

ALSTONIA Br

Pharmacology and toxicity of Alstonia alkaloids, Patricia Koegh and F. H Shaw, *Australian J. Expt Biol Med Sci* 21, 183, 1943

A SCHOLARIS Br

Total alkaloids in Alstonia scholaris The Popular and well known indigenous antimalarial drug Rakshit, J N , *Sci & Cult* 1944, 9, 303, Quinine and Alstonia scholaris (CHHATM) in Malaria Das Gupta, B M , Siddons, L B , & Chakaravarti, H *Ind Med Gaz* , 1943 79, 408

Total alkaloids in Alstonia scholaris Dutt, A T , *Sci & Cult* 1944 9, 556

ALTHAEA OFFICINALIS Linn

Oils from Althaea officinalis and Malva arborea, H Ya Tropp, *Chem Abst* 1597, 1936

A. ROSEA Linn

Damping off in Hollyhocks Srivastava, H C , *Sci & Cult* , 1951, 17, 91

AMARANTHUS GANGETICUS Linn.

Availability of Calcium in Lady's finger (Hibiscus esculentus) cabbage (Brassica oleracea, capitata), Drumstick (Moringa oleifera) and Amaranth Tender (Amaranthus gangeticus) Pt I Availability of Calcium in vegetables determined by experiments on growing rats Basu, K P , & Ghosh, D , *Ind Jour Med Res* , 1943, 31, 37

Examination of the fatty oil from the seeds of Amaranthus gangeticus Chidambaram & Iyer, *J Ind Chem, Soc* 22 (1945), 117.

AMARANTUS BLITUM Linn

Colchicine induced polyploidy in Amaranthus blitum Tandon, S L , & Chinoy, J J , *Sci & Cult* , 1950, 15, 398

AMOMUM XANTHIOIDES Wall

Constituents of Amomum xanthioides Wallich T. Kariyone and Y. Yoshida, *J. Pharm Soc. Japan* 50, 54, 1930.

ANACARDIUM OCCIDENTALE Linn

The food value of the nut of Anacardium occidentale (Hijli Badam). Chatterjee, N. K , *Ind Med. Gaz* , 1930, 65, 12.

Reaction products, Cashew nut-shell oil The Harvil corp. *Chem. Abst* 2584, 1931.

Composition of Rubber and Cashew nut-shell liquid, Mortimer T. Harvey, *Chem Abst* 5720, 1930

The industrial utilization of Cashewnut, Feloz Malberti, *Chem Abst* 1815, 1932.

A new globulin from Cashew nut (*Anacardium occidentale*) Damodaran and Tarakad G Sivaswamy, *Biochem J.* 30, 604, 1936

Cashew Nut-Shell oil as a mosquito larvicide Wats R C, & Bharucha, K, H., *Curr Sci*, 1937-38, 6, 216.

Cashew Apple a rich source of Vitamin C K Mitra, *Science & Culture*, 6, 186, 1940

Phenols from Cashew nut-shell liquid. Mortimer T. Harvey, *Chem. Abst* 1210, 1941

Study of the nuts of acajou (*Anacardium occidentale*) Francica Rosa Bonchristiano, *Chem Abst*, 339, 1941)

Ethers of cashew nut-shell liquid Solomon Caplan, *Chem. Abst* 4779, 1941

Mildew in Cotton, Synthetical experiments in antiseptics for textiles Derivatives of Cashew nut oil R C Gandhi and K. Vankataraman, *Ind Jour Chem Soc & Ind*, 89, 1942

Biological value of the protein of the Cashew nut, Cyro Camargo Nogueira, *Chem Abst*, 3263, 1942

Components of the latex of *Anacardium occidentale* Linn. (Cashew nut) H J Backer and N H Haack, *Chem Abst*, 1394, 1943.

Subsidiary constituents of Cashew Nut shells Pt I, Bahl, O P., Rastogi, Vishwa Nath Sharma & Salimuzzaman Siddiqui; *J Sci Indl Res*, 1946, **8B**, 222

Hypoglycemic action of *Anacardium occidentale* (Cashew) in normal individual, Francisco Arduino and Maria de leudes N G. Soares., *Chem Abst*, 10296, 1951

ANAGALLIS ARVENSIS Linn

Fine suspended poisonous plants, *Anagallis arvensis*. E. Murray Pullar, *Australian, Vet J* 15, 19, 1939

ANAMIRTA COCCULUS W & A

Preparation of purotoxin, *Anamirta cocculus*. E. P Clark, *J. Am. Chem Soc*, 57, 1111, 1935.

ANANAS Tourn ex Linn

A preliminary study of the anthelmintic activity in vivo of fresh pine apple juice Conarado F., Assenjo, *J Am. Pharm Assoc.* 29, 8, 1940

ANCHUSA TINCTORIA Lam.

The constituents of Alkanet root (*Anchusa tinctoria*) Part II Anchusin and its derivatives. Majundar & Chakravary, *J. Ind. Chem Soc* 17 (1940), 272

ANDROGRAPHIS ECHIOIDES Nees

A note on *Andrographis echioides* Nees, Rangaswami, S., Subbarao, V., *Ind Jour Pharm*, 1951, 13, 63

A PANICULATA Nees

Andrographis paniculata Nees, on the bitter principle from. Part I Moktader, A & Sircar, S S Guha, *J Ind Chem Soc* 16 (1939), 333
Studies in the specification of Indian medicinal plants Pt II *Andrographis paniculata* Sen Gupta, S B, Dutta, P P., & Banerjee, S, *Ind Jour Pharm* 1948, 10, 72

Studies in the specification of Medicinal Plants *Andrographis paniculata* Nees Sen Gupta, S B, Banerjee, S, & Chakravarty, D, *Ind Jour Pharm*, 1949, 11, 77

Andrographolide, the active constituent of *Andrographis paniculata* Nees A preliminary communication Chakravarti R N, & Mrs Chakravarti, D, *Ind Med Gaz*, 1951, 86, 96

ANDROPOGON Linn.

The constituents of some Indian Essential Oils Part VIII The Essential Oil from a new species of *Andropogon* occurring in the Etawah District, U P Simonsen J L, *Indian Forest Records*, 1924, Vol X Part VIII

Toxicity of Sorghums, Em Miege, *Chem Abst* 3013, 1933

Hydrocyanic acid content of Sorghum varieties James F Couch Reinhold R Bruise and J W Martin, *Chem Abst* 4629, 1939

A JWARANCUSA Jones

The Essential oil from *Andropogon jwarancusa*. Jones and the Constitution of Piperitone—J L Simonsen, (*Trans Chem Soc* 1921, Vol. 119 p 1644)

The Essential oil from *Andropogon jwarancusa* and the Constitution of piperitone John Lionel Simonsen, *J Chem Soc* 1644, 1921.

The constituents of some Indian Essential Oils Part IV-The Essential Oil from *Andropogon jwarancusa*, Jones Simonsen J. L. and Rau M Gopal, *Indian Forest Records*, 1922, Vol IX Part IV, pp 18-22

The constitution of the terpenes present in the essential oil from *Andropogon jwarancusa*, Jones. John Lionel Simonsen, *Jour. Chem Soc* 2292, 1922

The constitution of the Terpene present in the Essential oil from *Andropogon jwarancusa*, Jones J. L Simonsen, (*Trans Chem Soc* 1922, Vol 121, p 2292)

A. KUNTZEANUS Hack

Studies in Indian Essential Oils Essential oil from the flower heads and stalks of *Andropogon kuntzeanus* Hack Var. *Foveolata*, Hack, Rao B. Sanjiva *Journal of the Indian Institute of Science, Bangalore*, 1932, Vol 15A, Part VII, pp. 75-77

A SQUARROSUS Linn.

The Cus Cus Oil in India-Puran Singh; (*Chemist & Druggist* 1914, Vol. LXXXV) p 51, Aug, 8.

ANETHUM SOA Roxb

Notes on some Indian Essential oils Anethum soa, Roxb (Dill oil). Rao B Sanjiva, Sodborough J J and Watson H E, *Journal of the Indian Institute of Science, Bangalore*, 1925 Vol 8A, Part X, pp 183-184

Anethum soa, green herb and seeds—*Ind Acad Sc Vol XII, Sept*, 1939, p 251)

Chemical investigation of the essential oils derived from Anethum soa Roxb oil from the green and the seeds Braj Kishore Malvia and Sikhishushan Dutt, *Proc Ind Acad Sci* 12A, 251, 1940

ANGELICA ARCHANGELICA Linn

The essential oil in the roots of Angelica archangelica L grown in Dutch soil S P, Dijkstra, *Pharm Weekblad*, 81, 387, 1946

ANISOMELES MALABARICA Br

Chemical constituents of Anisomeles malabarica (N O. Labiatae) Essential oil Pt I., Rao, S B, & Majumdar, D N., *Ind Jour Pharm*, 1945, 7, 123

ANNONA MURICATA Linn.

The alkaloids of Annona muricata Linn Th -M Meyor, *Chem. Abst.* 8206, 1941

A RETICULATA Linn

Alkaloids from Annona reticulata Linnacus Alfred C, Santos Philippine, *J Sci* 43, 561, 1930

A SQUAMOSA Linn

Alkaloids from Annona squamosa leaves N Trimurti, *J. Ind Inst Sci* 7, 232, 1924

Oils and fats from the seeds of Indian Forest, plants Annona squamosa Ram Chandra, R V Ghanekar and P R Ayer, *J Ind Inst Sci* 10A, 1927.

Isolation of Anonine from annono sauamosa Linn, Felciana R Reyes and Alfredo C Santos Philippine, *J Sci* 44, 409, 1931.

Anonine Alfredo C Santos Phillipine, *J Sci* 47, 357, 1932

Seed oil of Formosan Plants Constituents of seed oils leguminosae, Annona squamosa Kinzo Kafulice and Chintu Hata, *J Chem Soc Japan*, 55, 369, 1934

ANOGEISSUS LATIFOLIA Wall

The tanning value of Anogeissus latifolia leaves *Anon Bull Imp Inst* 29, 137, 1931, 425, 1929

ANTHEMIS NOBILIS Linn.

Anthemis nobilis flowers, E Berthlome, *J. Pharm Belg* 15, 595. 1933

Beiglan essential oils Anthemis nobilis, Ernst S. Gremtner, *Am. Perfumer*, 33, 71, 1936

ANTIARIS TOXICARIA Leschen

A preliminary note on the pharmacological action of Antiaris toxicaria Chopra, R N & De, Premankur, *Ind Jour Med Res.*, 1934, 21, 513

Glycosides and aglucons L Antiarin (Antiaris toxicaria) K Dobel, E Schittlar and T Reichstein, *Helv Chem Acta* 31, 688, 1948

APIUM GRAVEOLENS Linn

Oil of Celery (Apium graveolens) Ernest G Guenther, *Amer. Perfumer*, 31, 75, 1935

AQUILARIA AGALLOCHA Roxb

The Nature of "Agaru" formation Bose, S R., *Sci & Cult*, 1938, 4, 89

ARACHIS HYPOGEA Linn

The properties of arachin and conarochin and proportionate occurrence of these proteins in the peanut D Breise Jones and Mullard J Horn, *J Agr Res* 40, 673, 1930, *Chem News*, 141, 38, 1930

Utilization of ground nut meal Basu, U P & Sen Gupta, S K, *J Ind Chem Soc* 21 (1944), 389

ARCHANGELICA OFFICINALIS Hoffm

Addition of nitrosyl chloride to B Phellandrene and the occurrence of the Phellandrene in some essential oils Archangelica officinalis Trustham F West, *J Soc Chem Ind* 58, 122, 1939

ARCTOSTAPHYLOS UVA-URSI Spreng

The constituents of diuretic drugs, the flavoxol glucoside of Folium Uva-Ursi and Folium vaccinii, Harnkitu Nakamura, Tatu Ota and Genitiro Hukuti, *J Pharm Soc Japan*, 55, 800, 1935

Arbutin content of infusion, decoctions and maceration of bearberry leaves (Uva Ursi) L Zechner, *Pharm Monatsh* 17, 47, 1936

Effect of synthetic growth substances on various types of cuttings of Arctostaphylos uva-ursi J A De France, *Proc Am Soc Hort Sci* 36, 800, 1939

Standardisation of the leaves of Arctostaphylos unva-ursi, D S Blinits-ka, *Chem Abst* 7488, 1939

Conditions under which some substances of Uva Ursi can exert an antibacterial action in urine, G Madaus and Fr E Koch, *Chem Abst* 7489, 1939

Composition of the ashes of the drugs Ash of Spanish Uva Ursi leaves, L Rosenthaler and G Beck, *Pharm Acta Helv U*, 186, 136

ARECA CATECHU Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency XIV Pulvis arecae. Caines, J F & Mhasker, K S, *Ind Jour Med Res*, 1921, 9, 206

Analysis of areca nuts, P Bourect; *Bull. Sci Pharmacol*, 98, 1933

The catechol in the fruit of *Areca catechu* Linn Ryo Yamamoto and Tosihiro Muraoka, *Chem Abst* 727, 1933

Areca nuts and aloes H Octtel, *Apoth Ztg* 48, 1457, 1933

Betel nut as a useful teniofuge Hsio lian Lik, *Chinese Med Jour* 50, 1273, 1936.

Steroles, unsaponifiable constituent of *Areca catechu*, L Sho Kuwada and S. Yoshiki, *J Pharm Soc, Japan* 266, 1937

The Pharmacological action of areca alkaloids, Arecoline hydrochloride Chukadonago Folia, *Pharmacol Japan*, 26, 73, 1938

Koleroga disease of Areca nut Uppal, B N, & Desai, M K *Curr Sci* 1939, 8, 122

Principal physical and chemical characteristics of dum palm nut fat as originally present and after hydrolysis, Ivo Ubaldini, *Chem Abst* 419, 1939

Anthelmintics containing areca nut, H Schlegel Arch *Expt Path Pharmacol*, 192, 389, 1939

Improved assay for Areca, *Rept Chem Abst* 2134, 1940

The pharmacological action of areca alkaloid, Arecaidine hydrochloride and arecaidenemethyl iodide, Ches Kadonaga Folia, *Pharmacol Japan*, 28, 57, 1940

Possible use of tannin from the kernels of green betel nuts Luz Baens, *Chem Abst* 6143, 1941

Pharmacology of the alkaloids and alkaloid drugs (areca nuts) Hans Braun, *Chem Abst* 4490, 1941

Nicotine like action of arecolina, U S V Euler and B Dome, *Acta Pharmacol, Toxicol*, 263, 1945

The areca and the betel Andre Mericier, *Chem Abst* 956, 1946

ARGEMONE ALBA Lestib

The alkaloids of Argemone alba, Lestib, P A Foots, *J Am Pharm. Assoc* 21, 246, 1932

A. MEXICANA Linn

Alkaloids of Argemone mexicana Alfredo C Santos and Pacifica Adkilen, *J Am Chem Soc* 54, 2923, 1932

Maxican argemone, Ode Almeda costa, *Chem Abst* 1811, 1934

(Maxican Poppy) Argemone mexicana L. Oswaldo de Almeida Costa, *Chem Abst*. 4901, 1935.

Investigation into the epidemiology of Epidemic Dropsy Pt XIII Application of the biological test to modified Argemone oil and its derivatives Lal, R. B., Das Gupta, Agarwal, S. P.. *Ind. Jour. Med. Res*, 1941, 29, 813.

Investigation into the epidemiology of Epidemic Dropsy Pt XIV
Feeding experiments on human subjects to test the toxicity of some
of the derivatives and modifications of Argemone oil Lal, R B, Das
Gupta, Agarwal, A. C, *Ind Jour Med Res*, 1941, 29, 839

A sensitive test for the detection of Argemone oil Sarkar, S N
Curr Sci, 1941, 10, 405

Chemical test for detection of Argemone oil Mukherji, S P, *Curr
Sci*, 1942, 11, 279

A sensitive chemical test for the detection of argemone oil, the
specificity of the test S N Sarkar, *Ann. Biochem and Exptl Med*
2, 101, 1942

A chemical method for the estimation of alkaloids present in Arge-
mone oil and its application to a mixture of argemone and mustard
oils, Sarkar, S N Rahman, Md Bazlur, *Curr Sci*, 1945, 14, 196

Detection of Argemone oil in Mustard oil, A K Sen, *J. Proc Inst
Chem (India)*, 18, 102, 1946

Argemone oil, Sen, A K *Ind Med Gaz*, 1946, 81, 126

Argemone and Mustard seeds Sanyal, P K, *Ind Med. Gaz.*, 1950,
85, 498

Epidemic Dropsy A new test for Argemone oil Chakravarti,
R N, Chaudhuri, R N., & Chakravarty, N K, *Ind Med Gaz* 1950,
85, 344

ARGYREIA SPECIOSA Sweet

Argyrea speciosa, fatty oil from the seeds of Kelkar, Phalnikar &
Bhide, *J Ind Chem Soc*, 24 (1947), 83, 6

ARISTOLOCHIA INDICA Linn

Essential oil from the root of *Aristolochia indica*, Part I and II
Krishnaswami, Rao Manjunath and Menon, *Journal of the Indian
Chemical Society, Calcutta*, 1935, Vol XII

Aristolochia indica, Chemical examination of the roots Part I
Krishnaswami, P R, Manjunath, B L and Venkata Rao, S, *J Ind
Chem Soc* 12 (1935), 476

Aristolochia indica, Chemical examination of the roots of Part II
The essential oil Krishna Rao, U S, Manjunath, B L & Menon,
K N, *J Ind Chem Soc* 12 (1935), 494

Aristolochia indica Linn Chemical Examination of the root Part III
Isolation of the alkaloid Aristolochine Krishna Swami, P R Manju-
nath, B L, *J Ind. Chem Soc.* 14 (1937), 39

ARISTOLOCHIACEAE

Studies on Indian Medicinal Plants, Pt. I Aristolochiaceae. Chakra-
varty, H L., *Ind. Jour Pharm.*, 1944, 6, 96

ARNICA RUPP. ex Linn

Constituents of Arnica flowers Chemical & Pharmacological study
of arnica preparations, H Thies, *Chem Abst.* 3070, 1945.

A. MONTANA Linn.

Arnica montana (Physiological action of). H S. Simpson, *J Am Inst Homeopathy*, 19, 213, 1926

Flower constituents of Arnica montana, L H Dieterle and K Fay., *Arch Pharm*, 277, 65, 1939

Constituents of the flowers of Arnica montana L.H. Dieterle and K Engelhard, *Arch Pharm*. 278, 225, 1940

Action of Arnica montana on the circulation, A W Forst, *Arch Exptl Pharm*, 201, 242, 60, 1943

Pharmacology of Arnica monatana, Otto Gessner, *Abst.* 2121, 1950.

Action of different constituents of Arnica montana on the isolated frog heart. Elma Barz, *Chem Abst*, 46, 86

ARTABOTRYS Br.

Alkaloids of Artabotrys Artabotrin and a new alkaloid Suaveolin, Alfredo, E Santos and Feliciano R Ryeas, *Chem Abst* 2251, 1933.

A SUAVEOLENS B.

Alkaloidal content of Artabotrys suaveolens Blume Juaquin M Maranon, *Philippine J Sci.* 38, 259, 1929

ARTEMISIA Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum rutae (Ruta graveolens Linn), Butea monosperma Roxb, Melia azadirachta Linn, Punica granatum Linn, Picrasma excelsa Swartz, Vernonia anthelmintica Willd, Cocos nucifera Linn, (coconut), Caines, J F & Mkhaskar, K S., *Ind Jour Med Res*, 1923, 11, 353.

Indian Santonin Chopra, R N & Chandler, A C, *Ind Med Gaz* 1924, 59, 537

Indian species of Artemisia, Chopra, R N & Mukerjee, B, *Ind Med Gaz*, 1931, 66, 666

Indian Artemisias S Krishna and R. S Verma, *Quart Jour. Pharm Pharmacol*, 6, 23, 1933

The cultivation of Artemisia. Verma, R. S and Krishna, S, *Curr. Sci*, 1935-36, 4, 29.

A short report on the economical value of Artemisia growing in the N. W. F. Province Qazilbash, N A, *Curr Sci* 1935-36, 4, 51.

Mode of action of Santonin and Chenopodium oil, H. A. Oelkers, *Arch Exptl Path Pharmacol*, 196, 161, 1940

Synthesis of Santonin. Paranjape, K (Miss), Phalnikar, N. L., Bhide, B. V. & Nargund, K S, *Curr Sci.*, 1943, 12, 150.

A new synthesis of Santonin. (Miss) Mukerji, D., *Sci. & Cultr.* 1947, 13, 258.

Commercial utilization of Indian Santonica for manufacturing santonin. Qazilbash, N A, *Ind Jour. Pharm*, 1950, 12, 175.

A. ABSINTHIUM Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency III, Oleum absinthii. Caines, J F. & Mhaskar, K S, *Ind Jour. Med Res*, 1920, 7, 690

The reason for the curative action of *Artemisia absinthium* (Worm Wood) on rat Scabies G Madaus and Fr E Koch, *Chem. Abst* 5541, 1944

Formation of bitter glucosides in the glandular hairs of certain plants (*Artemisia absinthium*) Jean Politis, *Comp Red* 222, 910, 1946

A KURRAMENSIS

The Kurram santonica Qazilbash, N A, *Curr Sci*, 1943, 12, 233

A MARITIMA Linn

Camphor from Erven Worm Wood, A L Mudighoyan, *Artemisia maritima Chem Abst*, 5820, 1943

A SACRORUM Ledb var minor

Oil of *Artemisia sacrorum* Ledb Var minor Ledh Z. G. Chestova, *Chem Abst* 3940, 1936

A SCOPARIA Waldst. & Kit.

Chemical examination of *Artemisia scoparia*, isolation of an essential oil and lactone Dharmpal Parihar and Shikibhushan Dutt, *Proc Ind Acad Sci*, 25 A, 153, 1947

A VULGARIS Linn

Nutritive value of the leaves of *Artemisia vulgaris*, L Var Indica Maxim R Sakai and T Masaki, *J. Pharm Soc Japan*, 53, 251, 1933

Blumea densiflora and *Artemisia vulgaris*, their insecticidal and larvicidal properties R N Chopra and S M. Ghosh, *J Malaria Inst India*, 3, 495, 1940

ARTOCARPUS HIRSUTA Lamk

Artocarpus hirsuta Lamk Fixed oil from the seeds of Varier *J. Ind Chem Soc* 22 (1945), 275.

A. INTEGRIFOLIA Linn

Artocarpus integrifolia Crystallographic investigation of Artostemonone, the stenone isolated from the Indian summer fruit by means of Goniometer & X-rays Nath, M C & Mukherji, D. L.; *J Ind Chem. Soc* 16 (1939), 229

On the occurrence of a mixed inflorescence in *Artocarpus integrifolia* Linn Guha, M. P., *Sci & Cultr*, 1945, 11, 99

Vivipary in *Artocarpus integrifolia* Linn Venkate, V., *Sci. & Cult.*; 1950, 15, 361

ARUNDO DONAX Linn

Arundo Tourn. *J Sci Indl Res.*, 1946, 6A, 276.

ASARUM EUROPEUM Linn.

Composition of Asarum oil (*Asarum europeum* L) Viktor Bruckner and Tibor Szeki, *J. Prakt. Chem*, 134, 107, 1932

Comparative results of toxicological investigations of the Romanian plant *Asarum europeum* and the Japanese *Sieboldii* Julium Orient., *Chem Abst*, 6247, 1934.

Active principles of *Asarum europeum* L. Leaves M A. Abdulmenev, *Chem Abst*, 7411, 1946.

ASCLEPIADACEAE

The existence of Resinole in Asclepiadaceae Rama Murti & Sheshadri, T R, *Ind Jour Pharm*, 1944, 6, 86

ASPARAGUS Linn

Sapogenins (Sapogenins) from Indian Medicinal Plants I Sapogenins from *Asparagus* Rao, S B, *Ind Journ Pharm*, 1952, 14, 131.

A ADSCENDENS Roxb

Studies on tuber mucilages Pt V Mucilage from the tuber of *Asparagus adscendens* Roxb Rao, P A, Beri, R. M, & Budhiraja, R. P, *J Sci. Indl Res*, 1952, 11B, 127

A RACEMOSUS Willd.

Studies of Tuber Mucilages Pt IV Mucilage from the tubers of *Asparagus racemosus* Rao, P S, Beri, R M & Budhiraja, R P, *J Sci Indl Res*, 1951, 10B, 261

Studies on tuber mucilages Pt VI. Partial hydrolysis of the mucilage from the tubers of *Asparagus racemosus* Willd Rao, P.S & Budhiraja, *J Sci Indl. Res*, 1952, 11B, 209

ASTERACANTHA LONGIFOLIA Nees

Investigation of Indian Medicinal plants, *Asteracantha longifolia* N K Basu, S B Lal and S N Sharma, *Quat Jour Pharm. Pharmacol*, 20, 38, 1947

Investigation of Indian Medicinal plants, *Asteracantha longifolia*, *Trianthema monogyna*, *Boerhaavia diffusa* N K. Basu, S B. Lal and S N Sharma, *Quart Jour Pharm Pharmacol*, 20, 38, 1947

Chemical investigation of *Asteracantha longifolia* Nees Basu, N K & Gode, K, *Ind Jour Pharm*, 1952, 14, 212

ASYSTASIA COROMANDELIANA Nees

A *Cercospora* leaf spot of *Asystasia coromandeliana* Nees. Mahmud, K. A *Sci & Cult*, 1952, 17, 474.

ATALANTIA MONOPHYLLA Corr.

Essential oil from *Atalantia monophylla* M T. Chobe and B. Sanjiv Rao, *Proc. Soc. Biol Chemist. India*, 2, 16, 1937.

Oil of *Atalantia monophylla* M T Chobe and B. S. Rao; (*Proc. Soc. Biol Chem India*, V 2, 1937) Part I

Essential oil from *Atalantia monophylla* Nayak & Guha, *J. Ind Chem Soc* 28 (1951), 713.

ATHYRIUM FILIX-FOEMINA Roth

A study of *Athyrium filix-foemina* S Trupp and Forres, J. Goodrich, *J. Am Pharm Assoc* , 29, 286, 1940

ATRIPLEX HORTENSIS Linn

Salt bushes (*Atriplex hortensis*) and their allies in the United States G L Bidevall and N S Golding, *Chem Abst* , 1120, 1926

ATROPA Linn

Distribution of alkaloids in two species of belladonna. E Est , F. Cabelleron, Josefina Tomas and Y Royo, *Chem Abst* 7600, 1937

Manufacture of Atropine from Belladonna Roots Srivastava, G. P , & Basu, N K , *Ind Jour Pharm* , 1944, 6, 3

Increased alkaloidal contents of induced polyploids of *Datura*, *Atropa* and *Hyoscyamus* *Datura* species Hack Marris Rowson, *Quart Jour Pharm Pharmacol* 18, 175, 1945

Note on the quality of Tincture Belladonna of Indian manufacture Bhattacharya, A., *Ind Jour Pharm* , 1947, 9, 78

A comparative study of *Atropa* species grown in Kashmir Kapoor, L D , Handa, K L & Kartar Singh, *Ind Jour Pharm* , 1951, 13, 249

A ACUMINATA Royle.

Assay of Indian belladonna root W A N Markwell, *Pharm J.* 146, 259, 1941

Seasonal variation in the alkaloidal contents of the leaves of *Atropa acuminata* , I C Chopra, K L Handa and L D Kapoor, *Ind Jour Agri Sci* , 16, 144, 1946

Pharmacognostic and chemical studies of Indian Belladonna, *Atropa acuminata* Herber W Youngken and William E Hassan, *J Am. Pharm Assoc Sci. Ed* , 37, 450, 1948

Indian belladonna, R Chatterjee and J K Lahiri; *J Am Pharm Assoc* , 38, 11, 1949

Cultivation of *Atropa acuminata* Royle ex Lendly Kapoor, I, D , Handa K L & Chopra, I C , *J Sci Indl Res* , 1952, 11A, 534

A BELLADONNA Linn

Some field tests showing the comparative efficiencies of Derris, Pyrethrum and Belladonna powders on different insects, Clyde C., Hamilton and Lousie G. Gemelle, *J Econ Entmol* 27, 446, 1934

Detection of Scopoletin in radix gelsemi and radix belladonna Robert Fischer and Herbert Ehrlich, *Arch. Pharm* , 274, 268, 1936.

The Bulgarian belladonna root. P Kuiper and P. Van der Weiler, *Pharm Week Bland*, 74, 1546, 1937

Analysis of the alkaloidal mixture in *Atropa belladonna* A Kuhn and G Schafer, *Chem Abst*, 4277, 1938

Drug extraction The effect of pressure and vaccum on efficiency of extraction belladonna root Em J Husa and Geo R. Jones, *J Am Pharm, Assoc*, 27, 852, 1938

Belladonna root and Bulgarian treatment Angelo Firrari, *Chem Abst*, 6397, 1938

Variation in the alkaloidal content of *Atropa belladonna* during a vegetative period A Kuhn and G Schafer, *Pharm Exnt Zentral-halk*, 80, 151, 1939

Intubation studies of the human small intestine The effect of atropine and belladonna on the motor activity of the small intestine and colon, Kendall A Elson and J L Drossener, *Am J Digestive Diseases Nutrition*, 6, 589, 1939

Bulgarian treatment of Parkinsons disease, Pharmacological aspects and chemical effects of alkaloids of belladonna root Hermann Vollmer, *Chem Abst* 5539, 1940

The value of Palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves Bernard S Einstein and Frank J Slama, *J Am Pharm Assoc*, 29, 370, 1940

Dynamics of accumulation of alkaloids extractives and inorganic compounds in different growth stages and diferent parts of the Belladonna plant N A Valyashko and A Sova; *Farmatsiya*, 4, 20, 1940

The growth effects of thiamine chloride, ascarbic acid and Phyto-hormones on Belladonna and Ricinus Loxis C Zoft, *J. Am Pharm Assoc* 29, 487, 1940

The determination of the quality of Belladonna N V Kunznestov, *Chem Abst* 856, 1940

The effect of varying temp, upon the total alkaloid content of folia Stromonii and folia Belladonna H Fluck, *Chem Abst* 3878, 1940

Suggestions for the collection, preparing and keeping Belladonna root for Pharmaceutical purposes Arturo Nanniz, *Chem Abst* 2672, 1941

Alkaloids of Bulgarian Belladonna root Harold King and Hancelot L Wave, *J Chem Soc* 331, 1941

Alkaloids of Belladonna root H Kreitmair, *Chem Abst* 9541, 1941
Germination of Belladonna seeds R Melville, *Pharm J.* 146, 116, 1941.

Study of Portugese Belladonna. Manuel Riberiro Cabral, *Chem Abst* 1811, 1942.

Amino neuteric determination of alkaloids in Belladonna, lobelia herb and harmel roots G. Ya Kait, *Chem. Abst* 2996, 1942.

Experimental investigation of the influence of Belladonna, atropine and other substances on the strychnine poisoning of white mice Fr E. Koch and Fr Naumann, *Chem Abst* 6623, 1942.

Some diseases of Belladonna in California and their control Johan T. Middleton, *Plant diseases Rept* 25, 513, 1941 *Chem Abst* 2985, 1942

Analysis of alkaloidal mixture in *Atropa belladonna* L. W. Marki, *Pharm Acta Helv*, 18, 215, 1943

Belladonna cultivation in Eastern Washington, Morris Wolfred; *Am. J. Pharm*, 115, 100, 1943

The influence of the application of artificial fertilizer on the concentration of active ingredients (total alkaloids) in the roots of *Atropa belladonna* E. De Conno, *Chem Abst* 3218, 1943

The differentiation of *Scopolia* and *Belladonna* leaves. D. Markovic, *Chem. Abst.* 10, 1, 9, 1944

Tincture of Belladonna Timoteo. A. Estevz, *Chem Abst*, 2790, 1944

Tetraploidy in *Atropa belladonna*, Gyula Szomolany, *Chem. Abst* 3318, 1944

Synthesis of Hyoscyamine in *Atropa belladonna* and *Datura stramonium*, B. T. Cromwell, *Biochem. Jour*, 37, 717, 1944

Culture study of the drug plant *Atropa belladonna*, W R. Brewer and Alex Laurie, *Chem Abst* 3419, 1944.

A belladonna and Hyoscyamus extract which is stable in the air and contains the alkaloids of the drug in mutually unchanged relative proportion, J Gjesing Andersen, *Chem Abst.* 6497, 1944

Biological assay of galenical preparations of Belladonna Jeanne Levy, *Bull Soc Chem Biol* 27, 431, 1945

Biosynthesis of the Belladonna alkaloids W O. James, *Nature*, 158, 654, 1946

Culture studies on the drug plant *Atropa belladonna*, W R Brewer and Alex Laurie, *Chem Abst* 4480, 1946

Experimental work in breeding medicinal plants. Brief report concerning the examination of properties of *Atropa belladonna* in 1948-49 Alenka Mstnak, *Chem. Abst* 1650, 1950

Influence of Belladonna tincture on the gastric activity of children. E. Pere Delgado and Garcia Morato Castano, *Chem Abst* 4576, 1950.

Belladonna roots and leaves cultivated in this country (as compared with those existing in the market). Felix A. Fares Rail; *Chem. Abst.* 6580, 1950.

Cultivation studies of the Solanaceous drugs Alkaloidal content of mature and unmatre belladonna leaves W R Brewer and L. David Hiner, *J Am Pharm Assoc* 39, 638, 1950

The inhibition phenomena in *Atropa belladonna* Juan Bautista Abad Manvique; *Chem Abst* 3098, 1950

ADIRACHTA INDICA A. JUSS. (MELIA) AZADIRACHTA Linn)

Chemical composition of Nim or Margosa oil *Ind. Jour. Med Res* 1920, 8, 356

Odorous and bitter constituents of neem oil E R Watson, N G Chatterjee and K C Mukerjee, *J Soc Chem Ind* 42, 387, 1923

Odoriferous principle of Neem Oil—*J S C I*, Sept, 1923, Page **387 T**

The correlation between chemical composition of Anthelmintics and their therapuetic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum Rutae (*Ruta graveolens* Linn), *Butea monosperma* Roxb, *Melia azadirachta* Linn, *Punica granatum* Linn, *Picrasma excelsa* Swartz, *Vernonia anthelmintica* Willd, *Cocos nucifera* Linn (coconut) Caines, J F & Mhaskar, K S., *Ind Jour Med Res.*, 1923, 11, 353.

A note on the efficacy of neem batties in the destruction of rat and rat fleas in rat burrows Tiwari, C D & Lal R B, *Ind Med Gaz* 1925, 60, 310

Neem oil, The bitter principle of Neem oil Part I Sen, Rajindra Nath & Banerjee, Ganapari, *J Ind Chem Soc* 8 (1931), 773

Melia azadirachta, The Chemistry of Oleo margosa from Neem oil Part I Insolation of the constituents of the oil—Mohammad Qudrat-i-Khuda, Ghosh, Subbash Kumar and Mukerjee Austosh, *J Ind Chem Soc*, 17, (1940) 189

A Margosa tree without the bitter principle Thirumalachar, M J, *Curr. Sci*, 1941, 10, 413.

A Margosa tree without the bitter principle Jacob, Cherian, K, *Curr. Sci.*, 1941, 10, 335 & 414

Fatty acids of neem oil, C J Dasa Rao and T R Seshadri, *Proc. Ind. Acad. Sci*, **15A**, 161, 1942

Bitter principles of the Neem oil Rangaswami, S., *Curr. Sci*, 1942, 11, 367

Utilization of Nim oil and its bitter constituents (Nimbidin series) in pharmaceutical industry. Salimuzzaman Siddiqui and Chittaranjan Mitra; *Jour Sci Indl Res*, (1945-46), 4, 5.

Chemical examination of the flowers of *Melia azadirachta*. Subramanian, S. Sankara & Rangaswamy, S; *Curr. Sci*, 1947, 16, 182

Chemical examination of Nim blossoms (*Melia azadirachta*, Flora), Chittaranjan Mitra, Narasimha Rao, P., Satyendra Bhattacharji & Salimuzzaman Siddiqui, *J Sci Indl Res*, 1947, **6B**, 19

Denaturation of alcohol by Neem cake distillate Godbole, N N & Pande, G D, *J Sci Indl Res*, 1947, **6B**, 121

A note on the chemical examination of Nim blossoms (*Melia azadirachta*) Mitra, Chittaranjan & Siddiqui, S, *Curr Sci*, 1948, 17, 51

A note on the chemical examination of Nim (*Melia azadirachta*) exudate Satyendra Bhattacharji, Chittaranjan Mitra & Siddiqui, S, *J Sci Indl Res*, 1949, **8B**, 187.

A note of the chemical examination of Nim (*Melia azadirachta*) Trunk bark Satyendra Bhattacharji, Mitra C and Siddiqui S, *J Sci Indl Res* 1949, 813, 188

Structure of the Neem (*Azadirachta indica*) Gum, Mukerjee, S, & Srivastava, H C, *Curr Sci*, 1951, 20, 127

The active constituents of Nim seeds (*Azadirachta indica* Linn) Nimbidin and its composition. Sen, A. C, Sriram Singh & D N Majumdar *Ind Jour Pharm*, 1951, 13, 265

Preliminary study of antibacterial substances from *Melia azadirachta*, Chopra, I C, Gupta, K C, Nazir, B N, *Ind J Med Res*, 1952, 40, 511.

Chemical examination of the root bark of Nim (*Azadirachta indica* Syn *Melia azadirachta*) Siddiqui S, Mitra C, Rao P N, *J S I R*, 1953, **4B**, 152

Chemical examination of the Trunk of Nim (*Azadirachta indica* Syn *Melia azadirachta*) Siddiqui S, Bhattacharji S., Mitra C, *J S I R*, 1953, **4B**, 154

BALANITES AEGYPTIACA Wall

Sapogenins of *Balanites aegyptiaca* George A R, Kon and W T Weller, *J Am Chem Soc* 794, 1939

Some tropical seeds and their contents. *Balanites aegyptiaca*, Franz Berger, *Chem Abst*, 1528, 1939

Steroles from *Balanites aegyptiaca* Russel E Marker, Wagner T.P, Paul R, Ulshafer Dale, P J Goldsmith and Clarence H Ruof, *J Am Chem Soc.*, 65, 1247, 1943

Oil from fruit of *Balanites roxburghii* C B Patel, *Curr. Sci* 12, 58, 1943.

A note on the oil from the fruit of *Balanites roxburghii* Patel, C B *Curr. Sci*, 1943, 12, 58.

Balanites aegyptiaca and its various uses in Tehad territory P Creach, *Chem Abst.* 3777, 1944

Oil from the kernels of Lalob fruit, *Balanites aegyptiaca*, S A. Hussain, F G Golllear and R.T O. Connor; *J. Am Oil Chemists Soc* 26, 730, 1949

A chemical study of *Balanites aegyptiaca* Jean S. Goldfiem, *Chem Abst* 8572, 1949.

Balanites roxburghii, saponin from the seeds of Dhekni & Bhide *J Ind. Chem Soc.* 28 (1951), 588

BALANOPHORA POLYANDRA Griff

On the occurrence of monoecious *Balanophora polyandra* Griff in the Visakapatam District, Madras Presidency Dutt, B.S.M., *Sci. & Cult*, 1951, 17, 131

BALSAMODENDRON MUKUL Hook.

The chemical investigation of gum resin from *Balsamodendron mukul* Hook Ghosh, S & Chopra, R N, *Ind Jour. Med Res* 1942, 309, 331.

BAMBUSA Schreb

Constituents of Bamboo K Azami and S Sengoku, *Chem. Abst* 2248, 1926

Japanese plants, the occurrence of free pentose in Bamboo shoots, Shigeru Komatsu and Yasuo Sasaoka, *Bull Chem Soc. Japan*, 2, 57, 1927

The composition of Japanese Bamboo. Y. Veda, K. Kasama and K. Kimura, *Chem Abst* 3977, 1928

Biochemical study of Bamboo Shigeru Komatsu, *Chem Abst* 4807 1930

Biochemical studies in Bamboo Chemical development in the growth of shoots Chōji Tanaka, *Chem Abst* 983, 1931

Chemical investigation of Bamboo The pentosans of Bamboo Sulezo Oguri, *J Soc Chem Ind Japan*, 34, 233, 1931

Biochemical studies on Bamboo, Wie Sun Tac, *J Chem Soc Japan*, 52, 882, 1931

Chemical study of Bamboo Study of Bamboo cellulose. Sutez Oguri, *J. Soc Chem Ind Japan*, 35, 347, 1932

Lethal properties of aqueous extract of young Bamboo shoots Stewart, A.C, Moorthy, U N, *Ind Med. Gaz.*, 1935, 68, 320.

B ARUNDINACEA Willd

Chemical and pharmacological examination of young sprouts of *Bambusa arundinacea*. N N. Ghosh & R N. Chopra; *Arch. Pharm* 2760, 351, 1938.

BARRINGTONIA ACUTANGULA Gaertn.

Chemical examination of the seeds of *Barringtonia acutangula* Gaertn
J K. Lahiri and S Ghosh , *J Am Pharm Assoc.* 31, 193, 1942

BASSIA LATIFOLIA Roxb

The uses of Mowrah (*Bassia latifolia*) , A S Carlos , *Chem Abst* 1719, 1927

Production of alcohol from spent Mohwra (*Bassia latifolia*) flowers
D N Sahasraduthe and V G Patwardhan , *Poona Agr Coll Mog* 22, 45, 1930

Indian seed fats, Mowha (*Bassia latifolia*) and Tamal (*Garcinia morella*)
fats D R Dhingra, G L Seth and P C Speers , *J Soc Chem Ind* 52, 116, 1933

Some Indian vegetable oils (*Bassia latifolia*), G N Bhattacharya ,
Ind J Phys 10, 403, 1936

The unsaponifiable of Mahua oil from the United Provinces of India,
(*Bassia latifolia*) N N Godbole and P D Srivastava, *Chem Abst* 5193, 1937.

Kinetics of saponification of Mohawa flowers (*Bassia latifolia*) , R N
Gobhill and R N Gobhill and N G Chatterjee , *Ind Soap J* 5, 101, 1938

Fatty acids and glycerides of solid seed fats, seed fat of Madhuca
(*Bassia latifolia*), Mowarahfat T P Hilditch and M B Ichaporia ,
J Soc Chem Ind , 57, 44, 1938

Mohua oil soap and rancidity (*Bassia latifolia*) A N Ghose , *Ind
Soap J* ; 4, 265, 1938

Sapogenins, Bassic acid (*Bassia latifolia*) B Jason Heywood George,
A R Kon and Lancelot L Wane , *J Chem Soc* 1124, 1939

A preliminary note on the preparation of syrup from Mahua flowers
(*Bassia latifolia*) V S Abhyankar and N Narayana , *Poona Agr Coll*
33, 168, 1942

Industrial Utilization of Mahua Chatterji, N G , *J Sci Indl Res* ,
1944-45, 3, 265

Causation of rancidity in Mahwa oil (*Bassia latifolia*) B P Srivastava
and S N. Sethumadhya Rao , *Soap Perfumer & Cosmetics*, 24, 673,
1951.

B. LONGIFOLIA Linn

Chemical examination of the essential oil derived from the fruit of
Bassia longifolia Salgursarn Nigam and Shikhibhushan Dutt ,
Ind Soap J , 11, 131, 1945

BAUHINIA ACUMINATA Linn

Some abnormal flowers of *Bauhinia acuminata* Linn Trivedi, B S , &
Nigam, P N , *Sci & Cult* , 1952, 17, 438

B VARIEGATA Linn

Bauhinia variegata Linn. The fatty oil from the seeds of. Punlambekar, S V & Krishna, S , *J Ind Chem Soc* 17, 1940), 96

Bauhinia variegata leaves as cattle feed Kehar, N D & Goswami, M N., *Sci. & Cult* , 1951, 16, 476.

BELAMCANDA CHINENSIS Leman

Components of commercial Herbageranu (*Belamcanda chinensis*) Toshio Tominaga , *J Pharm Soc Japan* 62, 189, 1924

Glucoside from *Belamcanda chinensis*. L.C Mannich P. Schumann and Wan Ho Lin., *Arch Pharm* 275, 317, 1937

BENINCASA CERIFERA Savi

A study of the oils from the seeds of *Luffa aegyptica*, *Benincasa cerifera* and *Allium cepa* K D Phadnis, A V Rege, D G Pishawikar and S.V Shah , *J Univ Bombay*, **17A** 62, 1948

BERBERIS Linn

Berberine in the treatment of oriental sore Das Gupta, M B & Dikshit, B B., *Ind Med Gaz* , 1928, 64, 67.

Pharmacological action of Berberins Chopra R N , Dikshit B B & Chowhan, J S , *Ind Jour Med Res.*, 1932, 19, 1193

Berberine and Berberine-containing plants in pharmacology and therapeutics Chopra, R N , Dikshit, Chowhan, J S , *Ind Med Gaz* , 1932, 67, 194

Importance of Berberis in German therapy Walther Awe and Gg Schweizer , *Chem Abst* 5997, 1938

Chemical assay of Rasaunt and Hing from the Punjab market. Grewal, K S , Kochhar, B D , *Ind Jour Med Res* , 1940, 28, 463

Alkaloids in the Himalayan Berberis, Histological distribution of. Chatterjee, R., *Sci & Cult* , 1942, 7, 571

Himalaya species of Berberies, chemical studies on Chatterjee, R., *Sci. & Cultr* , 1942, 7, 619

The histological distribution of Alkaloids in the Himalayan Berberis. R Chatterjee , *J Am Pharm Assoc* 32, 1, 1943.

Berberine in Malaria A preliminary note Brahmachari, U. N , *Ind. Med. Gaz.*, 1944, 79, 259.

B AQUIFOLIUM Pursh.

Estimation of alkaloids in preparations of *Berberis vulgaris* and *B. aquifolium*. H. Neugebauer and K. Brunwer, *Chem. Abst.* 3669, 1939.

B. ARISTATA DC

Berberine salts from "Rasot" roots, *Berberis aristata* Ray, J N, Roy, B S., *Sci & Cultr*, 1941, 6, 613

Alkaloidal constituents of the Bark of *B. aristata* (Rassaunt), Chakravarti, K K, Dhar, D C & Siddiqui, S, *J Sci Indl Res*, (1950) **9B**, 161

Alkaloidal constituents of the bark of *Berberis aristata* (Rassaunt). Chakaravarti, K K, *J Sci. Indl Res*, 1950, **9B**, 306

B FLORIBUNDA Wall

Berberis floribunda, isolation of Oxyacanthine, barbamine, berberine, epeberberine, Palmatine, dehydrocorvadaline, Jatrorrhizine & columabmine Chatterjee, *J. Ind Chem Soc*, 28 (1951), 225

B. HIMALAICA Arndt

Plant alkaloids, Part IV *Berberis himalaica* & *B. tinctoria* Chatterjee, R. Guha, M P. & Das Gupta, A K. *J Am Ind Chem Soc*, 29 (1952), 921.

B. INSIGNIS Hook

Umbellatine from *Berberis insignis* Hook R Chatterjee, *J Am. Pharm Assoc* 30, 247, 1941.

B. INSIGNIS Hook. f & B. UMBELLATA Wall

Berberis umbellata. The alkaloid of Part I Isolation and examination of Umbellatine Chatterjee, R *J Ind Chem Soc*, 17 (1940), 289

The alkaloid of *Berberis umbellata* Part II Chatterjee, R. *J, Ind Chem Soc* 19 (1942), 233

The alkaloid of *Berberis umbellata*, Part III. Chatterjee, G R, *J Ind Chem Soc* 19 (1942), 385

General pharmacology of umbellatine, a new alkaloid isolated from *Berberis umbellata* Wall and *Berberis insignis* Hook f and its use in the treatment of oriental sore. Gupta; J C & Kahali, B S, *Ind. Jour. Med Res*, 1944, 32, 53

B VULGARIS Linn

Berberis vulgaris (pharmacological action). H. S Stimpson, *J. Am Inst Homeopathy*, 19, 235, 1925

Berberine in the "Common Berberry" (*Berberis vulgaris*). E R Schultz, *J Am. Pharm Assoc* 15, 33, 1926

Pharmacology of *Berberis vulgaris*, L Henri Leclerc, *Chem Abst* 3731, 1935.

Estimation of alkaloids in preparations of *Berberis vulgaris* and *B. aquifolium*. H. Neugebauer and K. Brunwer, *Chem. Abst* 3669, 1939.

Pharmacological and chemical investigation of Berbery (*Berberis vulgaris*) Z Supek and D Tonnie, *Chem Abst* 9246, 1949

BETA VULGARIS Linn

Secondary products in the extraction of glutamine from beets (*Beta vulgaris*), C Ravenna and R Nuccorni, *Chem. Abst* 5486, 1929

Study of the activity of enzyme in the live leaves of *Beta vulgaris*
A S Okanenko, *Chem Abst* 5688, 1931

A hypoglucemic substance in Beets W Rychlik, *Chem Abst* 6842, 1934

A chemical study of Sugar beet during the first growth year Frank Knowles J E Watkin and F W F Hendry, *J Agr Sci* 24, 368, 1934

The estimation of sugar in the leaf of the manglod (*Beta vulgaris*).
James E. Van der Plank, *Biochem J* 30, 475, 1936

BETULA Tourn

Changes of some of the physical constants of Birch tar on prolonged storage with access to air T G Kovalev and V V Illanionc, *Chem Abst* 7061, 1935

B UTILIS D Don

Sesquiterpenes in Birch (*Betula utilis* D Don) K A Vesterberg and F. Nydahl, *Chem Abst* 4059, 1927

Chemical composition of tree bark Birch bark (*Betula utilis* D Don)
M. Sharkov and Belyaevskii, *Chem Abst* 5368, 1933

Tannin extract from Birch bark (*Betula utilis* D Don) A A. Mukhamedov, *Chem Abst.* 5573, 1933

The pigment from Birch bred K H Bauer and Dierich Ber **66B**, 1053, 1933

Birch Sap and its use (*Betula utilis* D Don) C Voss, *Chem. Abst* 3260, 1934

The paraffin hydrocarbon in the oil from Birch trees (*Betula utilis* D Don) F Petru and J Hadaeck, *Chem Abst* 3770, 1935

BIXA ORELLANA Linn

Pharmacologically utilizable constituents *Bixa orellana* L. Friedrich W Friese, *Pharm. Zentrachale* 76, 4, 1935

Philippine Annato dye as colouring agent (*Bixa orellana*) Santiago Tachico and Augustus P. West, *Philippine J Sc* 61, 429, 1936.

South American drugs. Chemical composition of Boldus, bixol a new alcohol from the oil of *Bixa orellana*, M. Bachstsz and G. Cavellini, *Chem Abst.* 8107, 1937.

Alleged presence of carstene in annato seeds (*Bixa orellana*). Gilberto Guimaraes Villela, *Chem. Abst.* 224, 1943

BLEPHARIS EDULIS Pers.

Blepharis edulis Pers Constituents of the seed of Part I Jagraj Behari Lal, *J Ind Chem Soc* 13 (1936), 109

Blepharis edulis Pers. Constituents of the seeds Part II The comp of the oil. Pendse, G P & Jagraj, Behari Lal, *J Ind. Chem Soc.* 14 (1937), 362

BLUMEA BALSAMIFERA D C

Sombaong *Blumea balsamifera* D C (Chemical investigation), Henri Leclerc, *Chem Abst* 2891, 1941

B DENSIFLORA D C

Blumea densiflora and *Artemisia vulgaris*, their insecticidal and larvicidal properties R N Chopra and S. M. Ghosh, *J Malaria Inst India*, 3, 495, 1940

B. ERIANTHA DC

Blumea eriantha DC Natural flavones Part IV On the constitution of Erianthin, the yellow colouring matter of. By Prafulla Kumar Bose & Dutt, Phanibhushan, *J Ind Chem Soc* 17, (1940), 45.

B. LACERA D. C

Blumea lacera, essential oil from the leaves Baslas & Deshapande, *J Ind Chem Soc* 27 (1950), 25

B. MALCOMII Hook f

The Essential Oil from *Blumea malcomii*—J L Simonsen and M Gopal Rau, (*Trans Chem Soc* 1922, Vol 121, p 876)

BOERHAAVIA DIFFUSA Linn

The pharmacology and therapeutics of *Boerhaavia diffusa* (Punarnava) Chopra, R N, *Ind Med Gaz*, 1923, 58, 203

Chemical examination of Purer-Nava or *Boerhaavia diffusa* Linn. Radha Raman Agarwal and Shikibhushan Dutt, *Proc Acad Sci* 4, 73, 1934

Chemical examination of Punarnava or *Boerhaavia diffusa* Linn Isolation of an alkaloid punarnavine Radha Raman Agarwal, Shikibhushan Dutt, *Proc Acad Sci* 5, 240, 1935

A comparative study of *Boerhaavia diffusa* Linn and the white and red flowered varieties of *Trianthema portulacastrum* Linn *Ind Jour. Med Res.*, 1940. 28, 475

BOMBAX MALABARICUM D C.

Bombax malabaricum T P Ghosh; *Ind Forester*, 61, 93, 1935.

Bombax malabaricum Oil from the seeds of Rao, Rao & Venkateswarlu, *J Ind Chem. Soc.* 20 (1943), 403

BORASSUS FLABELLIFER Linn

Investigation on the Palm juice Analysis of the fresh and fermented juice (Borassus flabellifer) V S Basrur and M Qureshi, *J Osmania Univ* 7, 19, 1939

Edibles from Borassus flabellifer (Palmyra palm) with special reference to Nira or sweet toddy K Mitra and H C Mitra, *Ind Jr Agr. Sci* 10, 824, 1940

Biological value of proteins of Palmyra fruits—Borassus flabellifer Linn S N Sarkar and A C Bose, *Ann Biochem Exptl Med.* 5, 59, 1945.

BOSWELLIA SERRATA Roxb

The Constituents of some Essential oils Part VII—The Essential oil from the gum-oleo-resin of Boswellia serrata, Roxb Simonsen J L, *Indian Forest Records*, 1923, Vol IX, Part XI

Constituents of the essential oil from the gum-oleo-resin of Boswellia serrata Roxb O Roberts, *Chem Abst* 566, 1924

Chemical constitution of gum from Boswellia serrata, M A Malandkar, *J. Indl Inst Sci* 8A, 240, 1925

Preparation of turpentine from Boswellia serrata (Roxb), gumoleo-resin. R S Pearson and Puran Singh, (*Indian Forest Records* Vol VI, Part VI)

BOTRYCHIUM VIRGINIANUM Sew.

Phyllody in Botrychium virginianum Sew Govindu, H C. & Narayana, H S., *Sci & Cult*, 1949, 15, 199.

BRAGANTIA WALLICHII Br

Chemical examination Manjunath, B L. & Shankara Rao, M S. *J Ind Chem Soc* 15 (1938) 646.

BRASSICA CAMPESTRIS Linn

The nitrogenous compound in oil cake of Brassica campestris Var. Chinensis Kiyohisa Yoshimura and Shiro Fujise, *J. Chem. Soc. Japan*, 45, 427, 1924

B JUNCEA Hook f & T

A simple method for the removal of the toxic alkaloids from mustard oil adulterated with Argemone Roy, A C., *Curr Sci*, 1950, 19, 91

Argemone and Mustard seeds Sanyal, P K., *Ind Med Gaz.* 1950, 85, 498.

Purification of mustard oil contaminated with Argemone Oil. Om Parkash, Atma Ram & Brahma Prakash, *Curr. Sci.* 1951, 20, 16.

B. OLERACEA Linn

Availability of Calcium in Lady's finger (*Hibiscus esculentus*), Cabbage (*Brassica oleracea* Var *capitata*), Drumstick (*Moringa oleifera*) and Amarnath Tender (*Amaranthus gangeticus*) Pt I. Availability of Calcium in vegetables determined by experiments on growing rats. Basu, K P & Ghosh D., *Ind Jour Med Res.*, 1943, 31, 29.

Availability of calcium in Lady's finger (*Hibiscus esculentus*), Cabbage (*Brassica oleracea* Var *capitata*), Drumstick (*Moringa oleifera*) & Amarnath Tender (*Amaranthus gangeticus*) Pt. II Availability of calcium in vegetables determined by metabolism experiments on a human adult Basu, K P Ghosh, D., *Ind. Jour. Med Res* , 1943. 31, 37.

Colchicine induced polyploidy in *Brassica oleracea* Var *Botrytis* L Tandon, S L , *Sci & Cult* 483, 16, 1951

BROSIMUM GALACTODENDRON D Don ex Sweet

Cow tree Chatterjee, D , *Sci & Cult* 1950, 16, 116

Introduction of 'Cow Tree' in India, Biswas, K , *Sci & Cult.*, 1950, 16, 195.

BRUCEA SUMATRANA Roxb

Composition of the seeds of *Brucea sumatrana*. Heikei Uno, *J. Pharm. Soc. Japan* 63, 579, 1943

BRYONIA ALBA Linn.

Bryonia alba, a typical animal proving T G Mitchell, *J, Am Inst. Homeopathy*, 19, 105, 1926

BRYOPHYLLUM CALYCINUM Salisb

Some common indigenous remedies, *Picrohiza kurroa*, *Erythrina indica*, *Sansiviera zeylanica*, *Pongamia glabra*, *Hygrophylla spinosa*, *Bryophyllum*, *colycisa Rheum emodi*, *Solanum indicum* R N Chopra, & S Ghosh, *Ind Med Record* 55, 77, 1935

Organic acids of the leaves of *Bryophyllum calycinum*, Geo W Pucher, *J Biol Chem* 145, 511, 1942.

BURSERA Linn

Mysore Linaloe oil *J Sci Indl Res* , 1944-45, 3, 516

BUTEA Roxb,

The treatment of intestinal worms with the Indigenous drugs. *Butea*, *Embelia* and *Kamala* Mukerji, A.K., Bhaduri, N V , *Ind Med. Gaz* , 1947, 82, 66.

B. MONOSPERMA (Lam.) Kuntze. (B FRONDOSA Roxb)

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency. Santonin, *Oleum rutae* (*Ruta graveo-*

lens Linn), Butea monosperma Roxb , Melia azadirachta Linn , Punica granatum Linn , Picrasma excelsa Swartz , Vernonia anthelmintica Wild , Cocos nucifera Linn. (coconut), Cames, J. F. & Mhaskar, K S , *Ind Jour Med Res* , 1923, 11, 353.

The chemical examination of the oil from the seeds Butea frondosa Roxb Tummin Katti, M C & Manjunath, B L , *J Ind Chem Soc* 6 (1929), 839

Butea frondosa flowers, isolation of a crystalline glucoside of Butin Jagaraj, Behari Lal & Dutt, Sikhishushan, *J Ind Chem Soc* 12 (1935), 262

Butea frondosa Roxb On supposed occurrence of acids with uneven number of C atoms in vegetable oils and fats Part I. The acid fraction of Mean M W 354 from the seeds Krishna Rao, U. S & Manju Nath, B L , *J Ind Chem Soc* 12 (1935), 611

Constitution of Burtin Isolated from Butea frondosa flowers. Behari, L J , *Curr Sci* , 1936-37, 5, 426

Butea frondosa Studies on the enzymes of the seeds, Part I Proteolytic enzymes Chatterjee, N. R , Ghosh, S & Chopra, R N , *J Ind Chem Soc.* 15 (1938), 101

Butea frondosa Studies on the enzymes Part II Lipolytic enzymes. Chatterjee, N R , Ghosh, S & Chopra, R N , *J Ind Chem Soc* 15 (1938), 107

Occurrence of free buteen and butin in the flowers of Butea frondosa. P Bhaskara Ramamurti and T R Sheshadri, *Proc Ind Acad. Sci.* 12A, 477, 1940

Preliminary report on the presence of an estrogenic substance in the storage root of Butea frondosa Supvanta, *Chem Abst* 2929, 1940

The Phlobatannins of Kino and buteagums G V Krishnamurti and T R Sheshadri, *Proc Ind Acad Sci* 22A, 134, 1945.

Chemical investigation of the seeds of Butea frondosa, Examination of the fixed oil Dharmpal Parihar and Sikhishushan Dutt, *Ind Soap Jour* 12, 26, 1946.

B SUPERBA Roxb

Butea superba Roxb Khasem Pangson Vongse *Chem Abst* 3088, 1938.

Chemical composition of flowers of Butea superba Subba Rao, V., Sheshadri, T R , *J Sci. Indl. Res* , 1949, 8B, 178.

CAESALPINIA BONDUCELLA Fleming

Some constituents of Caesalpinia bonducella nut. Part II Bonducella Nut oil Godbole, S. N Paranjpe, D. R. & Shrikhande, J. G ; *J. Ind Chem Soc* 6 (1929), 295.

Caesalpinia bonducella Flem. Chemical examination of seeds. Part I. Tummin Katti, M. C.; *J. Ind. Chem. Soc.* 7 (1930), 207.

Search for an antimalarial drug in the indigenous materia medica
Pt. II. *Caesalpinia bonducella* Fleming Mukherji, B, Ghosh B K &
Siddons, L B, *Ind Med Gaz.* 1943, 78, 285

C DIGYNA Rottl

Investigation on the isolation and constitution of the tannin from
Indian Teripods. Biswas, H G, *Sci & Cult*, 1943, 9, 90

Gallic acid from Teripods. Biswas, H G, *J. Sci Indl Res*,
1948, **7B**, 27

CALENDULA OFFICINALIS Linn

Saponin Series, Gualc saponin and saponin from *Calendula officinalis*
Alfred Winterstein and Gertraud Stein, *Chem Abst* 5170, 1951

CALLITRIS RHOMBOIDEA R Br

Notes on some Indian Essential Oils *Callitris rhomboidea* R Br
(Syn *Frenela rhombidea*. Endl) Rao B Sanjiva, Sudborough J J
and Watson H E, *Journal of the Indian Institute of Science, Banga-*
lore, 1925, Vol **8A**, Part X, pp 144-145

CALOPHYLLUM INOPHYLLUM Linn

The proximate composition of *Palomaria* seed, oil and resin
(*Calophyllum inophyllum*) F. A. Soliven, *Chem Abst* 3118,
192, 1924

CALOPLACA ELEGANS

Indian Lichens *Caloplaca elegans* Neelakantan, S & Seshadri, T R
J Soc Indl Res, 1952, **11B**, 126

CALOTROPIS Br

African arrow poisons Heart poisons in *Calotropis latex*, Gerhard
Hesse Franz Reicheneder and Hans Eysenboch, *Chem Abst*
1742, 1939

African arrow poisons Crystalline *Calotropis* resin Gerhard
Hesse, Hans Eilbracht and Franz Reicheneder, *Chem Abst*
4387, 1941

The Digitalis like principles of *Calotropis* compound with other
cardioactive substances K K, Chem C I Bliss and E Brown
Robbins *J Pharmacol.* 74, 223, 1942.

C GIGANTEA Br.

Calosterol, a sterol present in the milky juice of *Calotropis gigantea*
Kalipoda Basu and Madhab Chander Nath; *Biochem. J* 28, 1561,
1934.

Calotropis gigantea. On the proteinase in the milky juice, its puri-
fication and activation by ascorbic acid and glutathione. Basu,

Kalipada & Madhab Chandra Nath, *J Ind Chem Soc.* 13 (1936), 34
Chemical composition of *Calotropis gigantea* wax and resin, components of the latex P Bhaskara Ramamurti and T R Seshadri, *Proc Ind Acad Sci.* **18A**, 145, 1943

Chemical composition of *Calotropis gigantea* The resins of the root bark K J Balkrishna, P Bhaskara, Ramamurti and T R Sheshadri, *Proc Ind Acad Sci* **22A**, 138, 1945

Chemical composition of *Calotropis gigantea* flowers A comparison of the various parts of the plant P Bhaskara, Ramamurti and T R. Sheshadri, *Proc Acad Sci* **22A**, 304, 1945

Madar Juice, N Pitchandi (*Calotropis gigantea*) *J Proc Inst of Chemists*, 20, 34, 1948

A preliminary study of the toxicity of *Calotropis gigantea* Rathnasabapathy, V, Rao K & Krishnaswamy, A, *Ind Jour Med. Res*, 1949, 37, 483.

C PROCERA Br

African arrow poison calotropin (*Calotropis procera*) Gerhard Hesse and Franz Reicheneder, *Chem Abst* 1031, 1937

CALYOPTERIS FLORIBUNDA Lamk

The anthelmintic constituent of the leaves of *Calycopteris floribunda* Araya Puram, Ratnagiriswaran, Kumar B Sehra and Krishanna Swami Vankataraman, *Bioch J* 28, 1964, 1934

The constituents of calycopterin The yellow colouring matter of the leaves of *Calycopteris floribunda* Shah, R C Virkar, V V & Venkataraman, K, *J Ind Chem Soc* 19 (1942), 135

Anthelmintics—Anthelmintic activity of *Calycopteris floribunda* M L Khorana, D K Motiwala and K Venkataraman, *Proc Ind Acad Sci* **27A**, 121, 1948

CANANGA ODORATA Hk f & T

Cultivation of *Cananga odorata* and experimental production of oil of Ylang-Ylang in Java. W Bobloff, *Chem Abst* 985 1927.

C PATCHOULI

Analytical characteristics of Dutch East Indian oils of *Cananga patchouli* and Vetiver D R Koolhas and P A Rowan, *Perfum France*, 15, 245, 1937

Cananga patchouli and Vetiver oil, D R Koolhas and P A Rowan, *Perfumery Essential Oil Record* 29, 53, 1938

CANARIUM COMMUNE Linn

The fatty oil of the seeds from *Canarium commune*. Alph. Steger and J Van Loon; *Chem Abst* 3937, 1940

C STRICTUM Roxb

Travancore Essential Oils Black Dammar Moudgil K L and Aiyar K Sitarama, *Department of Industries, Trivandrum*, 1923, *Bulletin* No XVII, pp 11-17

CANAVALLIA ENSIFORMIS D C

A toxic principle in the seeds of *Canavalia ensiformis* A Orru and A Fractoni, *Chem Abst* 6123, 1946

CANNABIS Tournef

The hydrolysis of ptytic compounds derived from seeds of hemp, horse bean, horse chestnut, flax, wheat and embryos of rye W Jarosza, *Chem Abst* 5501, 1934

Resin content and Physiological activity of hemp grown in the region of Paris Rene Paris and Marce Lousie Merac, *Chem Abst* 9377, 1949

C SATIVA Linn (C INDICA Lamk)

Chemical examination of India hemp and its preparations R Weitz and Dardanne, *Bull Sci Pharmacol* 31, 321, 1924

The valuation of Charas Chatterjee, D N & Roy, N B , *Ind Med Gaz* , 1929, 64, 373

Cannabis indica, resin, Part II, Robert Sidney Cahn *J Chem Soc* 630, 1931

Cannabidiol and Cannabiol constituents of *Cannabis indica* resin. A Jacob and A R. Todd, *Nature*, 145, 350, 1940

The use of Hemp Drugs in India for Euphoric purposes Chopra, R N , *Sci & Cultr* , 1940, 5, 761

Cannabis sativa in relation to mental diseases and crimes in India Chopra, R N., Chopra, G S , & Chopra, I C , *Ind Jour Med Res* , 1942, 30, 155

Observations on the physiologically active fraction of Indian Hemp, *Cannabis sativa* Linn , Bose, B C & Mukerji, B , *Ind Jour. Med Res* , 1945, 33, 265

CAPPARIS SEPIARIA Linn

Vivipary in *Capparis sepiaria* Linn Mahabale, T S., *Sci & Cultr* , 1939, 5, 372

C SPINOSA Lam

Analysis of *Capparis spinosa* seed oil D Zabramnyl, A Ochakovskil and N. Petrova, *Chem. Abst* 6822, 1941.

CAPSELLA BURSA-PASTORIS Moench

Medicinal action of Capsella bursa-pastoreis, and also of its parasites, Cystopus candidus and Peronospora parasitica W Harste, *Arch Pharm*, 266, 133, 1928

Alcoholic extract of Capsella bursa-pastoris Attilio Bartola, *Chem Abst* 6063, 1932

C ANNUUM L

Pharmacodynamic action of the active principles of chillies (Capsicum annuum) Jose Lille and Eliseo Ramirez, *Chem Abst* 4836, 1935

Pungency in chillies (Capsicum annuum) a Mendelian character Deshpande, R B, *Curr Sci*, 1935-36, 4, 418

The nitrogen complex of Indian foodstuffs Condiments Pt II Chillies (Capsicum annuum) Coriander seeds (Coriandrum sativum) Narasimhamurthy, G, *Ind Jour Med Res*, 1968, 25, 863

(Capsicum) chillies, Vit C contents of Indian foodstuff Rothenheim, C A, Shaik Mohammad, H S & Cowlagi, S S, *J Ind Chem Soc* 15 (1938), 15

A case of variegation in Capsicum annuum L, Deshpande, R B, *Curr Sci* 1939, 8, 313

Vit C content of chillies, onion & garlic, both in raw state and when boiled with water Biswas, H G & Das, K L, *Ind Jour Med Res*, 1939, 27, 139

CARICA PAPAYA Linn

Protolytic enzymes of Carica papaya, N C Nag and H N Banerjee, *Trans Bose Research Inst Calcutta*, 1, 1930-31

The colouring matter of Carica papaya Ryo Yamamoto and Scityu, *Chem Abst* 3480, 1933

Carica xanthin, Kryptoxanthin monopalmate (Carica papaya), P Karrer, W Schlentz, *Helv Chem Acta*, 55, 1934

Carica xanthin of the Papaya and Citrus poonensis Hort Ryo Yamamoto and Akimasa Kato, *J Agr Chem Soc Japan*, 10, 754, 1934.

Carica xanthin a colouring matter in the fruit pulp of Carica papaya and Citrus poonensis, Ryo Yamamoto and Yosiakin Kato, *Chem Abst* 806, 1935

An alkaloidal principle isolated from Carica papaya seeds Tryambak B. Panse and Anand S Pranjpe, *Rascyanam* 1, 215, 1941

Some chemical constituents of papayas and their relation to flavour. S J. Lynch and W.M Fifield; *Chem Abst* 1135, 1941,

Carica papaya Linn , Intro-carpellary Fructification in Salam, M A , *Sci. & Cultr* 1942, 8, 143

The biological value of the proteins of papaya (*Carica papaya*) and Lady's fingers *Espil Med* 2, 71, 1942

Carpasemine isolated from *Carica papaya* seeds T B Panse and A S Pranype, *Proc Ind Acad Sci* **18A**, 140, 1943

Puetro Rican fatty oils Characteristic composition of expressed papaya (*Carica papaya*) seed oil C F Asenjo and J A Hoyco, *Oil and Soap*, 20, 217, 1943

Internal proliferation in *Carica papaya* L Sattar Khan, *Sci & Cultr* , 1946, 12, 194.

A case of vivipary in *Carica papaya* Linn Sen, P , *Sci & Cultr.* 1946, 12, 153

Papain Activation of Ray, *J Ind Chem Soc* 23 (1946), 313.

Carica papaya L , study of the fruit, C A Lolano, Salcedo, *Chem Abst* 2507, 1947

Damping off of Papaya (*Carica papaya* L) seedlings Babu Singh & Paharia, K. D., *Sci. & Cult* 1952, 17, 477

Safflower oil The component glycerides of Vidyarthi, *J Ind Chem Soc* 20 (1943), 45.

CARTHAMUS TINCTORIUS Linn

Australian research on the use of Safflower oil C, Macmillin, *Chem Abst* 6546, 1948.

A rapid method for determining the oil content of Safflower and sunflower seeds W Keith Kennedy and John Urau, *Chem Abst* 5209, 1949

Utilization of the Tobacco seed and Safflower seed oil in varnish and paints. Pt I Sharma, P G , Budhiraja. N C & Aggarwal, J S , *J Sci Indl Res* , 1951, **10B**, 33

CARUM CARVI Linn

Shiyah Zirah (black caraway), H Stanley, *Redgrove Pharm J* 127, 496, 1931.

The essential oil of Japanese orandahakka (Crapemint), Carvone Torice Nagasawa, *J Chem Soc Idn. Japan*, 41, 252, 1938.

C COPTICUM Benth

The manufacture of Thymol from Ajowan Lahhani I V , Sudborough J J and Watson H E., *Journal of the Indian Institute of Science, Bangalore*, 1921, Vol 4, Part V pp 59-84

Ajowan Oil—R R Sobte and P Singh; (*Perfum. Record*, 14, 1923 399)

The Manufacture of Thymol Inugauti N N , Bhate S R and Habib Uasan, *Department of Industries and Commerce, Hyderabad*, 1924, *Bulletin No. 8*

Production of Thymol from Ajowan !seeds Verghese James Gulali K C & Joshi, M L , *Curr Sci* , 1949, 18, 17

CARYOPHYLLUS AROMATICUS Linn

Clove industry in India (Article), *Sci & Cultr* 1938, 4, 221

CASSIA ABSUS Linn

Oil of Cassia absus Zafaruddin Ahmed, *Chem Abst* 8376, 1935

Cassia absus A new formula for Chaksine The alkaloid of C. absus and some expts On its constituents Kapur, Hansraj, Gaind, Kidar Nath, Narang, Kartar Singh, Raj, Jiranendra Nath, *J Ind Chem Soc* 17 (1940), 28

Studies in the alkaloids of Cassia absus Linn Pt I Vishwa Nath Puri, Vishwa Nath Sharma & Salimuzzaman Siddiqui, *J Sci Indl Res* , 1945-46, 4, 701

Chaksine Aggarwal, M L , Ray, J N , Sen, D C , *Sci & Cultr* , 1946, 12 201,

Pharmacological investigation of Chaksine, an alkaloid from Cassia absus Linn Pradhan, S N , Varadan, K S , Roy C , De N N., *J S I R* 1953, **8B**, 358

A note on the new formula for Chaksine, Siddiqui, S & Ahmad, Z, *J Ind Chem Soc* 18 (1941), 589

C ANGUSTIFOLIA Vahl

The effective constituent of Senna leaves. W Straub and H Gebhardt, *Arch Exptl Path Pharmacol* 181, 399, 1936.

A new crystalline component of Senna leaves (Cassia angustifolia), P Bhaskara, Ram Murti and S. Rangaswami, *J. Pharm* 2, 203, 1940

Further studies on Cathartic action in mice, Senna, Aloe, Cascara and bile salts, Lloyd W Hazleton and Kathleen D Talbert, *J Am. Pharm Assoc* 33, 170, 1944

Factors effecting the cathartic activity of Senna in mice Lloyd W Hazleton and Kathleen D. Talbert, *J Am Pharm Assoc* 34, 260, 1945

Crystalline active Senna glucosides, Sandoz Ltd, *Chem Abst* 587, 1945

Evaluation of the purgative activity of Senna extract, composition of bioassay and chemical assay of Senna H. O J Collier, E C. Filher, S. K. Paris and D. M. Bellis Wuart, *J. Pharm. Pharmacol* 21, 252, 1948,

Crystallised glucosides from Senna leaves and pods Sandoz Ltd, *Chem Abst* 2376, 1949

Biological assay of vegetable purgatives, Senna leaf, fruit and their preparations, T C Lou, *J Pharm Pharmacol* 1, 673, 1949.

The repeated administration of Tinnevely and Alexandria senna to mice Maribelle Woods and I W Crote, *J Am Pharm Assoc* 40, 198, 1951

C. FISTULA Linn

Abnormal flowers of Cassia fistula Linn Raghavan, T S , & Venkatasubban, K R , *Curr Sci* 1934, 35, 3, 256

The use of Cassia fistula in the treatment of Black-water fever Venkatachalam K & Ratnagiriswarn, *Ind Med Gaz.* 1941, 76, 211

Pharmacological investigation of Cassia fistula L and some Brazilian species of Cassia Richard Wasiky, *Chem Abst* 1562, 1943

The fruit-pulp of Cassia fistula Linn Karim, M A & Guha-Sircar, S S *Sci & Cultr* 1946, 12, 194

A study of Cassia fistula Pulp Miss Modi, F K. & Khorana M L , *Ind Jour Pharm* , 1952, 14, 61

C OCCIDENTALIS Linn

Poisoning with Cassia occidentalis is due to a toxic albumin Raymond Moussa, *Chem Abst* 3320, 1925

The fatty oil of Cassia occidentalis Linn seed A. Steger J Van Loon, *Chem Abst* 2208, 1934

C TOMENTOSA Linn

Anthracene derivatives from the genus Cassia Cassia tomentosa and C corymbosa E Maurin; *Chem Abst* 1289, 1927

C. TORA Linn

Constituents of Folium Ketsumu (Cassia tora) Genichire Fukuchi and Kanco Imai, *J. Pharm Soc. Japan*, 70, 568.

Cassia tora Linn. Chemical examination of the seeds of. Part I Subba Jois, H & Manjunath. B L., *J Ind. Chem Soc.* 7 (1930) 521.

CASSYTHA Linn.

Cassytha L on Viscum Dutt. B.M.S., *Sci. & Cultr.*, 1951, 16, 373.

C. FILIFORMIS Linn.

On Cassytha filiformis Linn. Srivastava, J , *Curr. Sci* 1945, 14, 242

On the range of Hosts of Cassytha filiformis Linn. Nayar, B.K & Nayar, P N , *Sci & Cultr.*, 1952, 17, 383.

CASUARINA EQUISETIFOLIA Forst.

Sex in *Casuarina equisetifolia* Forst. Chandrasekharan, S N., Parachasarathy, S V & Sundararaj, D D., *Sci & Cult*, 1949, 15, 158.

CATHA EDULIS Forsk.

Central stimulating action of the alkaloid Cathine (*Catha edulis*) *Arch. Exptl Path and Pharmacol* 198, 100, 1941.

CEDRELA TOONA Roxb

Essential oil from the wood of *Cedrela toona*, Roxb. P. Parameswara Pillai and B Sanjiva Rao, *J Soc. Chem Indl* 50, 220, 1931.

Cedrela toona Roxb Isolation of a lactone, an essential oil and a colouring matter from the wood of. Parihar & Dutt, *J Ind. Chem Soc* 27 (1950), 77.

CEDRUS DEODARA Loud.

The constituents of some Indian Essential Oils Part III. The Essential Oil from *Cedrus deodara*, London Simonsen J L, and Rau M Gopal, *Indian Forest Records* 1922, Vol IX, Part IV, pp. 13-17

CELASTRUS PANICULATUS Willd.

Seed and fruit coat fats of *Celastrus paniculatus*, B.G Gunde and T P Hilditch, *J Chem. Soc.* 1980, 1938

A note on the chemical examination of *Celastrus paniculatus*, Warsi. Sharifuddin Ahmad *Curr Sci*, 1940, 9, 134

Chemical investigation of *Celastrus paniculatus* Willd. N.K Basu, P R Pabrai, *J Am Pharm Assoc* 35, 272, 1946

A note on the chemical investigation of the fruit of *Celastrus paniculatus* Willd (N O Celastraceae)

Shah, M M, Phalankar, N L, Bhid, B V, *Curr Sci* 1947, 16, 57

CENCHRUS PAUCIFLORUS Benth

New Plant immigrant, *Cenchrus pauciflorus* Benth (Gramineae, A). Sen, S K, *Sci & Cultr*, 1942, 459, 7.

CENTIPEDA ORBICULARIS Lour.

Centipeda orbicularis. Vyas, B N & Sinha, H K, *Ind Med. Gaz.*, 1930, 65, 75

Chemical examination of *Centipeda orbicularis* Lour Rastogi, R P Sharma, V N. & Dhar, M L, *J S I R*, 1953, 7B, 332.

CEPHAELIS IPECACUANHA Rich

Review of the Pathogenesis of Ipecac and its alkaloids, Megavack, T H, *J. Am Inst of Homeopathy*, 21, 300, 1928

Indian Ipecacuanha. Chopra, R.N. & Mukerjee, B., *Ind. Med. Gaz*, 1932, 67, 88.

Treatment of chronic intestinal amoebiasis with Gavano, a derivative of Ipecacuanha. Chopra, R N , Sen, S. & Sen, B, *Ind. Med Gaz* , 1934, 69, 130,

Action of emetine on the activity of the adrenal & thyroid glands. Chopra, R.N , Gupta, J.C & Roy, A C , *Ind Jour Med. Res.*, 1935, 22, 771.

Radix Ipecacuanha and its alkaloids as expectorant and emetics Pfisers A , *Pharm Ztg* 81, 503, 1936.

The emetic action of Ipecacuanha, Mukhnacheva A I , *Chem. Abst* 6449, 1936.

Biological studies on tincture Ipecac Astrue A , Girovex J and S Barrau, *J Pharm Chem* 1, 185, 1940

Assay of Indian Ipecacuanha Guha, R C , & Mukerji, B , *Sci. & Cultr.*, 1945, 11, 204.

Pharmacognostic studies on 'Ipecacunha'. Bal, S N., & Datta, S C *Sci & Cult.*, 1945, 10, 448

Pharmacognostic studies on Indian Ipecacuanha Bal, S N , & Datta S C , *Sci & Cult* , 1946, 12, 201.

Pharmacognostic studies on Indian Ipecacuanha Bal, S N & Datta, S C *Ind Jour. Pharm* , 1946, 8, 76.

Ipecacuanha (*Cephaelis ipecacuanha*) of Brazil Arnaldo Augusto Addor, *Chem Abst* , 1812, 1947

Observations on the growth of *Cephaelis ipecacuanha* (Bort) A Rich in Calcutta. Mitra, G C & Chakravarti, D , *Sci & Cult* 1948, 13, 504.

On the cultivation of Ipecacuanha in India Biswas, K P & Sampathkumaran, M A , *Sci & Cult* 1948, 14, 160.

A process for the preparation of emetine HCl from Indian Ipecacuanha Shah, S K , *J Sci Ind Res* , 1951, 10B, 76

CEPHALANDRA INDICA Naud

Cephalandra indica (Telakucha) in Diabetes Chopra, R.N , & Bose, J P , *Ind Med Gaz.*, 1925, 13, 11

Observations on the antidiabetic preparations of *Cephalandra indica* (Telakucha) Chopra, R N , & Bose, J P , *Ind Med. Gaz* 1925, 60, 201.

CERBERA ODOLLAM Gaertn

Oil and fats from the seeds of Indian forest plants, oil from the seeds of *Cerbera odollam*. Ram Chandra, Ghanekar R V. and Ayyar, P.R. *J Ind Inst Sci* 10A, 20, 1927

A preliminary study of pharmacological action of the glucoside of *Cerbera odollam* Chopra, R N., Bose, B.C., Gupta, J.C. & Chopra, I.C., *Ind. Jour Med Res.*, 1942, 30, 107.

CHAMAENERIUM ANGUSTIFOLIUM Schur.

Comparative plant chemistry *Chamaenerium angustifolium* Konstantia Puringer, *Montash* 44, 247, 1924

CHEIRANTHUS CHEIRI Linn.

Heart active glucosides of *Cheiranthus cheiri* Jaretzky R and Wilke M, *Arch Pharm* 270, 81, 1932.

CHENOPODIUM Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency. Oleum Chenopodii Caines, J.F & Mhaskar, K S, *Ind Jour Med Res*, 1920, 7, 570.

Indian Chenopodium Chopra, R N., & Mukerjee, B, *Ind Med Gaz* 1932, 67, 5

A comparative record of anthelmintic treatment with tetrachloroethylene and oil of Chenopodium Manson, D, *Ind Med Gaz*, 1934, 69, 500

Pharmacological study of Chenopodium oil M Aissi Manicini and Leonardo Donatelli, *Chem Abst* 4451, 1935.

Pharmacological investigation on Chenopodium oil. Leonardo Donatelli, *Chem Abst* 4217, 1936.

The comparative value of oil of Chenopodium and tetrachloroethylene as anthelmintic for use in mass treatment. Hare, K P., Dutta, S C, *Ind Med Gaz*, 1939, 74, 198

A short note on the mass anthelmintic treatment by oil of Chenopodium and tetrachloroethylene Majumdar, D C, *Ind Med Gaz*, 1940, 75, 652

Pharmacology of chenopodium oil Oelkers H A, *Arch Exptl Path. Pharmacol* 195, 315, 1940

Effect of heat and light on the Ascaridol content in oil of Chenopodium Mukerji, A R, & Sen Gupta, K K, *Ind Med Gaz.*, 1945, 80, 347

C ALBUM Linn.

A phytochemical study of the fruits of *Chenopodium album* Tonuta J and Kaufmann K.L, *Pharm Arch* 12, 27, 1941.

C AMBROSIODES Hance

Saponin and sapogenin of *Chenopodium ambrosiodes* L. O. Dafert, Bauer, F Baur, M Capesius V and Grifinger S, *Chem Abst* 6937, 1934.

Studies on saponins of *Chenopodium ambrosiodes*, Grifinger L S, *Chem. Abst* 4839, 1937.

Investigation of the oil from *Chenopodium ambrosiodes* L. Var anthelminticum Gray. cultivated in the gardens of medicinal plants of the Stepan Botany Univ. in Wilno. Tadensz Burchacinshi; *Chem. Abst* 8916, 1939.

A preliminary investigation of the yield and composition of the oil distilled from *Chenopodium ambrosioides* L; Var *anthelminticum* Gray. R.E Shapter, *J. Coun Sci Ind Res.* (Australia) 14, 201, 1941.

Culture tests on *Chenopodium ambrosioides* L. Var *anthelminticum*. The value and determination of the volatile oil. Schmonthuin, A & Schmied, E, *Chem. Abst.* 216, 1944

CHLOROXYLON SWIETENIA Dc

Alkaloid isolated from *Chloroxylon swietenia*, constitution of. Mookerjee & Basu, *J. Ind Chem Soc.* 23 (1946), 1.

Chloroxylon swietenia. Alkaloid from the bark of. Mookerjee & Bose, *J. Ind Chem Soc* 23 (1946)

CHRYSANthemum CINERARIIFOLIUM Vis.

Insecticidal principle in *Chrysanthemum cinerariifolium*. Insecticidal principle produced on dry distillation of *Chrysanthemum cinerariifolium* Ryo Yamamoto and Mizuho Sumi H.; *Chem Soc Japan*, 44, 1070, 1923

Pyrethrum and Derris insecticides as arsenical substitute Roy Huston, *Chem Abst* 6234, 1934

Some field tests showing the comparative efficiencies of Derris, Pyrethrum and Belladonna powders on different insects Clyde C. Hamilton and Lousie G Gemelle, *J Econ Entmol.* 27, 446, 1934.

Determination of Pyrethrum I in commercial insecticides containing Pyrethrum and pyrethrum extract D A. Holaday, *Chem Abst* 3078, 1938

Pyrethrum cultivation in Kashmir Fotidar, M R, *Curr Sci*, 1940, 9, 360

Treatment of Scabies and Pediculosis with Pyrethrum Roy, D N, Ghosh, S.N & Chopra, R N, *Ind Med Gaz*, 1941, 76, 333.

Pyrethrum cultivation in Kashmir (*Med & Public Health Sec*), *Sci Cultr.*, 1941, 7, 217

Further work on Pyrethrum in the treatment of Pediculosis. Roy, D N. & Ghosh, S M, *Ind Med Gaz.*, 1942, 77, 480.

A water emulsion of pyrethrum extract for spray killing adult mosquitoes Russel, P P, Knipe, F W, *Ind. Med. Gaz.*, 1942, 77, 477.

Pyrethrin content of Indian Pyrethrum Puntambekar, S.V., *Curr. Sci.*, 1943, 12, 232

Pyrethrum in Kashmir Chopra, I.C., Dhar M.L, Handa, K.L. Habibullah, M. & Asa Nand, *Curr. Sci*, 1945, 14, 104.

Carotenoid pigments of Pyrethrum flowers Subbaratnan, A.V., & Parameshwaram Pillay, P, *J Sci Indl Res.*, 1947, 6B, 100.

Pharmacognostic studies on homegrown Pyrethrum flowers Mitra, G C.; *Ind Jour. Pharm*, 1950, 12, 100

Cultivation of Pyrethrum in Kashmir, Chopra R.N., Kapoor L.D., Handa K.L. and Chopra I C *Ind. Farming* Vol. VIII 78, 1947

CICER ARIETINUM Linn.

γ Galactan contained in the seeds of Cicer arietinum and its molecular constitution N Castoro, *Chem Abst.* 2642, 1925.

The food value of Cicer arietinum. V Zagami, *Chem Abst.* 3764, 1934

CICHORIUM INTYBUS Linn

Cichorium Intybus Linn Chemical examination of the seeds of. Constituents of the oil from the seeds Misra, Ramnath & Dutt Sikhishushan, *J Ind Chem Soc* 14 (1937), 141

CICUTA VIROSA Linn

The toxin in Cicuta virosa E. Svagr. *Chem Abst* 407, 1924.

A case of the poisoning of six boys by the root nodules of Cicuta virosa V M. Karasek, *Chem. Abst* 7418, 1940.

CIMICIFUGA RACEMOSA Nutt

Constitution of the rhizomes of Cimicifuga racemosa Fern Mercier and J Balansaid, *Comp. Rend Soc Biol* 118, 166, 1935

CINCHONA Linn.

The relative value in malaria of the Cinchona alkaloids Quinine, Cinchonine, Quinidine, Cinchonidine & Quindoidine and the two derivatives Hydro-quinine and Ethyl Hydro-Cuprine MacGilchrist, A C, *Ind Jour. Med Res*, 1915, 3, 1.

On the behaviour of Paramecium caudatum towards the Cinchona alkaloids Acton, H W, *Ind Jour. Med Res*, 1921, 9, 339

Comparative action of the principle alkaloids on the dog heart (quinine, quinidine, Cinchonine, (Cinchonidine) A Elerc, C Pezzi and G Perro Gahaud, *Chem Abst* 293, 1924

Pharmacological notes on the principal alkaloids of Cinchona. L. Barthe, *Chem Abst* 700, 1925

A comparative study of the action of Cinchona alkaloids on the uterus Chopra, R N, David, J C & Dikshit, B B., *Ind Jour. Med. Res*, 1928, 15, 571

The cultivation of Cinchona in India A K Y, *Curr Sci.*, 1939, 8, 435

Cinchona plant. Bal, S.N, *Sci & Cultr*, 1939, 4, 617

Bengal Cinchona Plantation (Med & Public Hsalth Sec), *Sci. & Cultr.*, 1941, 7, 300

Cinchona cultivation in India—Sen, S.C., *Curr Sci*, 1941, 10, 223

Quinine and *Alstonia scholaris* (CHHATM) in Malaria. Das Gupta, B.M., Siddons, L.B. & Chakaravarti, H., *Ind Med. Gaz.*, 1943, 79, 408.

Cinchona, pharmacognostic studies on Datta S C, & Bal S N, *Ind. Jour Pharm*, 1946, 8, 85

Pharmacognostic studies on Indian Cinchona Datta, S.C., & Bal, S N, *Sci & Cult*, 1946, 12, 246

Influence of temperature on growth and alkaloidal content of Cinchona seedlings Harold F Winters, Arnaud, J Loustalot and Norman F Childens, *Chem Abst* 2126, 1947

Waste Cinchona materials for production of Quinine Mukerjee, S, *Sci & Cultr* 1949, 13, 372

Cinchona alkaloids of newly cultivated plants of Assam. Mukerjee, S, *Ind Jour Pharm*, 1949, 10, 107

Quinine from Cinchona bark extractions Pt I & II Bhunvara, N.B, & Khorana, M L, *Ind Jour Pharm*, 1949 11 Pt I, 148 and Pt II 152

Structure of Quinamine and Cinchonamine A note on the Str of Quinamine and Cinchonamine, two minor alkds of Cinchona Tyabji, A, *Ind Jour Pharm*, 1950, 12, 243

Cinchona finds a new home Chatterjee, R, *Sci & Cultr*, 1950, 15, 339.

C FEBRIFUGA

Cinchona febrifuga in the treatment of malaria Gore, R N, *Ind Med Gaz.*, 1938, 73, 608

Soluble salts of "Cinchona febrifuga" as a cheap and effective anti-malarial drug Chatterjee, R P, *Sci. & Cultr* 1952, 17, 524

C LEDGERIANA Moens ex Trimen

Vegetative propogation of Cinchona ledgeriana from gootes (Marcotte) and cuttings by treatment with auxins Chakutra, A G, *Sci & Cult*, 1944, 9, 409

CINNAMOMUM Bl.

Cinnamon leaf oil Krishna, S, Kamath, H R., Kudva, K T & Kudva, K G, *J Sci Indl Res*, 1945-46, 4, 464

C CAMPHORA Nees.

Note on the possibilities of camphor cultivation from Cinnamomum camphora in Northern India. Howard S H, Robertson W.A. and Simonsen J L, *Indian Forest Records*, 1923 Vol IX, Part VII, pp 34.

Notes on some Indian Essential Oils Cinnamomum camphora, Nees & Eberm (Syn Camphora officinarum, Baugh, Laurus Camphora, Linn) Rao B Sanjiva, Sudborough J J and Watson H E., *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. 8A, Part X, pp 160-174

Camphor seed oil. Hao Chen, *Ind Res. (China)* 6, 92, 1937.

Constituents of Camphor oil, Acids and Phenols in the brown camphor oil, Kashichi Ono and Monoru Imoto, *J. Chem Soc Japan*, 58, 531, 1937

C JAPONICUM Siebold.

Fat of the seeds *Cinnamomum japonicum*, as a substitute for Cocoa butter. Talsoo Karyone and H Iwao, *J Pharm Sco Japan*, 58, 238, 1938.

C ZEYLANICUM Nees.

Essential oil from the leaves of *Cinnamomum zeylanicum* V P Shintue and B. Sanjiv Rao, *J. Ind Inst Sci* 15A, 84, 1932

Studies in India Essential Oils X-Essential Oil from the leaves of *Cinnamomum zeylanicum*, Breyn Rao B Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1932, Vol 15A, Part VII, pp 84-87.

Cinamon-bark oil of the Seychelles (*Cinnamomum zeylanicum*) W Holdsworth Haines, *Perfumery Essential Oil Records*, 27. 6, 1936.

CISSAMPELOS PAREIRA Linn

Chemical Examination of the roots of *Cissampelos pareira* Linn Bhattacharji, S, Sharma, V N & Dhar, M L, *J Sci Indl Res*, 1952, 11B, 81

A preliminary note on the pharmacological action of the total alkaloids isolated from *Cissampelos pareira* Linn Roy, P K., Dutta, A T, Roy, G K, Mukerjee, B, *Ind J Med Res*, 1952, 40, 95.

CITRULLUS COLOCYNTHIS Schrad

A preliminary note on the chemical examination of the roots of *Citrullus colocynthis* Scharader. Agarwal & Dutta, S, *Curr Sci*, 1934-35, 3, 250

Citrullus colocynthis Examination of the oil from the seeds of Alimchandani, Badami & Katli, *J Ind Chem. Soc* 26 (1949) 515 & 519

C VULGARIS Schrad

Citrullin a new amino acid in press juice of the Watermelon, *Citrullus vulgaris* Schard. Mitsunori Wada, *Chem Abst* 5335, 1930

Proteins and other nitrogenous constituents of Watermelon seeds (*Citrullus vulgaris*) P S Krishnan and T K Krishnaswamy *Bioch J* 33, 1284, 1939

Characteristics and composition of Watermelon seed oil Arthur J. Nolte and Harry W. Von Loeseke; *J Am Chem Soc* 61, 869, 1939

Extracting the therapeutic principle from various parts of Watermelon fruit or plant. Mathew Evertz; *Chem Abst.* 5257, 1941.

The diuretic action of Watermelons I T Isawa, Y Takahasi and T. Togo, *Chem Abst* 521, 1941

Biological value of proteins in Watermelon and Pumpkin seeds Tsao E King, *Chinese J. Physio* 16, 31, 1941

Cucurbit-seed glabulins, use as substitute for edestin in experimental diets, (*Citrullus vulgaris*) Hubull R B . Vockery H B and Nolan, L S *J Nutrition*. 25, 99, 1943

Citrullus vulgaris Component fatty acids of the oil from the seeds of Dhingra & Biswas, *J Ind Chem Soc* 22 (1945), 119

Melon seed oil Safet Riz Alpar and Suat Esin; *Chem Abst* 9700, 1950

CITRUS Linn

Preservation of citrus fruit juices Jiwan Singh Pruthi & Girdhari Lal, *J Sci Indl Res* , 1951, 10B, 36

Nagpur Oranges—(*Proc Nat Acad Sc* Vol I, 1939, pp 175-180)

C ACIDA Roxb

Xanthyletin Isolation of. from Citrus acida bark Mookerjee, *J Ind Chem Soc* 23 (1946), 41

Citrus acida Roxb Natural coumarins isolated from the leaves of Khastagir, *J Ind Chem Soc* 24 (1947), 421-34.

C AURANTIUM Linn

Preparation of Citric Acid from Citrus aurantium or Karana Khatta Dhingra, D R., Gupta, G.N., Nigam, V.N , & Raghvendra Singh, *Ind Jour Pharm*. 1950, 12, 351

C BIGARADIA Loisel.

Notes on some Indian Essential Oils Citrus bigaradia, Risso Rao, B Sanjiva, Sudborough, J J and Watson, H E , *Journal of the Indian Institute of Science, Bangalore*, 1925 Vol 8A, Part X, pp 176-179

C. DECUMANA Linn.

Citrus decumana On the bitter principles of Miss Asima Mukerjee, *J Ind Chem Soc* 17 (1940), 593

Ascorbic acid of Shaddock (*Citrus decumana*) availability of by human subjects. De & Barai, *J. Ind. Chem Soc*. 25 (1948), 389

C. LIMON Burm f

Survey of the possibilities of manufacturing Citric Acid from Lemons in Bihar and Bengal. Ghatak, N. & Karimullah, J., *Sci Indl Res* , 1944-45, 3, 110.

A study of species of Citrus limonis. Thomas Oomen, P , *Ind. Jour Pharm* , 1951, 13, 120.

C POONENSIS Hort ex Tanaka

Carica xanthin of the papaya and Citrus poonensis Hort Ryo, Yamamoto and Akimasa Kato, *J Agr Chem Soc, Japan*, 10, 754, 1934

Carica xanthin, a colouring matter in the fruit pulp of Carica papaya and Citrus poonensis Ryo Yamamoto and Yosiakin Kato *Chem Abst* 806, 1935

CLAVICEPS PURPUREA (Fries) Tulasne

The therapeutic activity of liquid preparations of Ergot of the Calcutta market Chopra, R N & De, Premenkur, *Ind Med Gaz*, 1928, 63, 519

Ergot in India Padwick, G Watts *Curr Sci*, 1941, 10, 488

Can the deterioration of Ergot extracts be prevented in the tropics Bose, I B & Dey, N K, *Ind Med Gaz*, 1942, 77, 286

Germination of Ergot Bose, A B, *Curr Sci*, 1942, 11, 439

Ergot in Indian Kulkarni G S, *Curr Sci* 1942, 11, 246

Indian medicinal importance of Ergot and need for its cultivation Mukerji, B, Bose, A B, *Sci & Cultr*, 1942-43, 8, 267, 348

Claviceps purpurea (Fr) Tul and a new species from India Watts Padwick, G & Azmatullah, *Curr Sci*, 1943, 12, 257

Ergot on Sugarcane in Mysore Thirumalachar, M J, *Curr Sci* 1943, 12, 330

Assay of Indian Ergot Mukerji, B. & Dey, N K., *Curr Sci* 1943, 12, 58

A method for the assay of individual Ergot sclerotium Mukerji, B & De, N K, *Curr Sci* 1944, 13, 128

Ergot on Cynodon dactylon Pers Thirumalachar, M J, *Curr Sci*, 1944, 13, 288

Ergot and sphacelial stages on some wild grasses in Mysore Thirumalachar, M J, *Curr Sci*, 1945, 14, 22.

Ergot sclerotia on Sorghum vulgare Pers Ramakrishnan, T S, *Curr Sci*, 1948, 17, 218

Ergot on bamboo Ramakrishnan, T S & Ramakrishnan, K, *Curr. Sci*, 1949, 18, 344.

Ergot on two grasses from South India Ramakrishnan, T S & Sundaram, N V, *Sci & Cultr*, 1950, 16, 214

CLAUSENA WILLDENOWII Wight & Arn

Oil of Clausena willdenowii—B S Rao and K S Subramanian, *Proc. Indian Acad Sci.* 1A, 1934, 189.

Oil of Clausena willdenowii—B S. Rao and K S. Subramanian, *Proc. Indian Acad Sci. Sect. A*, 3-1936.

Chemical investigation on some Minor Forest Products. The Essential Oil from Karibe, *Clausena willdenowii*, leaves and twigs Rao K. Narain, Varadhan C. and Janniah S. L., *Department of Industries, Bangalore*

Constitution of L. Clausenan (*Clausena willdenowii* W. & A.). Narsimha Rao, P. L., *J. Sci. Indl. Res.*, 1949, **7B**, 11

CLEOME VISCOSA Linn

Cleome viscosa Linn. Chemical examination of the seeds of Part I. The constituents Gupta, Mahadeo Prasad & Dutt, Sikhishushan, *J. Ind. Chem. Soc.* 15 (1938), 532

CLERODENDRON INFORTUNATUM Gaertn

Constitution of clerodin, the active bitter principle of *Clerodendron infortunatum* H. N. Banerjee, *Trans. Bose Research Inst., Calcutta*, 12, 75, 1936-37

Clerodendron infortunatum, chemical examination of. Part I. M. B. Rane & Kondiah K., *J. Ind. Chem. Soc.* 14 (1937), 46.

Clerodendron infortunatum Active principle Clerodin of Chaudhry & Dutt, *J. Ind. Chem. Soc.* 28 (1951), 295

COCCULUS D.C

Sinomenium and Cocculus alkaloids Constitution of Cepharanthine Heisaburo Kondo and Ichiro Keimatu, *Ber.* **71B**, 2553, 1938

C. LAURIFOLIUS D.C

Alkaloid of Sinomenium and Cocculus Alkaloid of *Cocculus laurifolius* H. Mondo and T. Kondo, *J. Pharm. Soc. Japan* 524, 876, 1925

C. SARMENTOSUS Diels

The alkaloids from Sinomenium and Cocculus The alkaloids of *Cocculus sarmentosus* Diels Constitution of menisavine Heisaburo Kondo and Masao Tomita, *J. Pharm. Soc. Japan*, 55, 637, 1935

COCHLEARIA ARMORACIA Linn

The content of *Lepidium latifolium* and *Cochlearia armoracia*, in essential oils S. Rivos Goday and M. Gomez, serranillos, *Chem. Abst.* 4432, 1945

COCOS NUCIFERA Linn.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency. Santonion, *Oleum rutae* (*Ruta graveolens* Linn.), *Butea monosperma* Roxb., *Melia azadirachta* Linn., *Punica granatum* Linn., *Picrasma excelsa* Swartz, *Vernonia anthelmintica* Willd., *Cocos nucifera* Linn. (coconut). Caines, J. F. & Mhaskar, K. S., *Ind. Jour. Med. Res.*, 1923, 11, 353.

Chemistry of the products of *Cocos nucifera*. Juan P.C. Chandersona, *Bioch J* 24, 1493, 1930

Antirachitic effect of Coconut oil and its mode of action on bone Calcification Nimal C Dutt, *Ann Biochem Exptl Med.* 8, 69, 1948.

COFFEA Linn

The chemistry of Coffee bean The unsaponifiable matter of Coffee bean oil Preparation and properties Kahweol, R O. Bengis and R J Anderson, *J Bioch Chem* 97, 99, 1932

Coffee black bean Venkalarayan, S V, *Curr Sci*, 1938, 7, 236.

Chemistry of Coffee Elucidation of the constitution of Cafestrol, Carl H Slotta and Klas Neissar, *Ber* 71B, 2342, 1938

Chemistry of Coffee A new method for the determination of trigonoline Karl Heinrich Slotta and Klaus Neisser, *Ber* 17B, 1987, 1938

Chemistry of Coffee, recent analytical findings Karl H. Slotta and Klaus Neisser, *Chem Abst* 2604, 1939

A coffee substitute *Cassia occidentalis* that is toxic before roasting. P Bruri, *J Pharm Chem* 20, 321, 1942

Coffee oil K H Bauer and R Beu, *Chem Abst* 5607, 1943.

Determination of Coffee in coffee mixtures and substitute by the Tillmanns—Hollatz method Jean Deshuses *Chem Abst* 3033, 1944.

C ROBUSTA Linn

Importance of reserve food material in successful establishment of shoot cutting in *Coffea robusta* L Rallabharaman, T V & Gopalkrishnan, K S, *Curr Sci* 1947, 16, 385

COLCHICUM Linn

Colchicine Bose, A., *Sci & Cultr* 1940, 5, 511

COMBRETUM PILOSUM Roxb

Combretum pilosum Roxb as an anthelmintic for *Ascaris lumbricoides* Ramsay, G C, *Ind Med Gaz*, 1922, 57, 374.

COMMELINA ASIATICA Linn

Chemical Examination of the oil from the seeds of *Commelina asiatica* Linn Aggarwal, J S & Miss Padmini Soni, *J Sci. Indl. Res* 1949, 8B, 49

CONVOLVULUS HAMADAE (Vved) Petrov.

Alkaloids from *Convolvulus hamadae* G V Lazurevskii; *Chem Abst.* 4029, 1941.

C PLURICANTES Chois.

On an abnormal flower of *Convolvulus pluricantes* Chois. Rao, A R ; *Curr Sci.*, 1935-36, 4, 441.

COPTIS TEETA Wall

The constituents of Coptis teeta Wall Chatterjee, A , *Sci & Cultr* , 1950, 15, 330

Alkaloid from Coptis teeta Wall Saw, P L Seshadri, T R , *J Sci Indl. Res* , 1952, **11B**, 308

Plant alkaloids Part II Coptis teeta Wall Chatterji, R Guha, M.P. & Chatterjee, A , *J Ind Chem Soc* 29 (1952), 97.

CORCHORUS Linn

Corchortoxin a cardiac agent from jute seeds P Karrer and P Banerjee, *Helv Chem Acta* 32, 2385, 1949

C OLITORIUS Linn

Corchorus olitorius Composition of the oil from Chakravarti & Sen, *J Ind Chem Soc* 20 (1951), 790

CORIANDRUM SATIVUM Linn

Notes on some Indian Essential oils Coriandrum sativum Linn Rao B Sanjiva, Sudborough J J and Watson H E , *Journal of the Indian Institute of Science, Bangalore*, 1925. Vol **8A**, Part X, pp 179-182

Report on the examination of Indian and foreign coriander (Coriandrum sativum) B. Vishwanath and C V Ramaswamy Ayyar, *Ag Livestock India*, 4, 583, 1934

The fatty and the essential oils from Coriandrum sativum grown in the Northern Caucasus I I Vanin and A A Chernoyarova, *Chem Abst* 3833, 1934

The nitrogen complex of Indian foodstuffs Condiments Pt II Chillies (Capsicum annum) Coriander seeds (Coriandrum sativum) Narasimhamurthy, G , *Ind Jour Med Res* , 1938, 25, 863

Production of faboil from the Coriander seed waste at the Aleksevskii plant F Tansue Slenko, *Chem Abst* 5649, 1938

Production of decaldehyde from Coriander P P Shorygin and V P Osipova, *Chem Abst* 3781, 1942

Decaldehyde from the oil of blooming Coriander Ya Bryusova Ryu Shagolova and N Novikova, *Chem Abst* 3781, 1942

Production of citral from Coriander oil V I Isagulyants and E K Smolyani *Chem Abst* 3782 1942

COSCINIUM FENESTRATUM Colebr

A note on the alkaloids of Coscinium fenestratum (Colebr) Varier, N S & Pillai, P P , *Curr Sci* 1943, 12, 228

The berberine content of Coscinium fenestratum (Colebr.) Child, R., Narhanael, W.R.N , *Curr Sci* , 1943, 12, 255

CRATAEGUS OXYCANTHA Linn.

Constituents of Crataegus oxycantha H Dieterle and O Dorner; *Arch Pharm.* 275, 428, 1937

Investigation of *Crataegus oxyantha* Thomas E Pugh, *Chem. Abst* 3462, 1939

A pharmacological study of *Crataegus* tincture I I Sivertsev, *Chem Abst* 6463, 1939

Pharmacological action of *Crataegus oxyantha* J D P Graham, *Quart Jour Pharm Pharmacol* 13, 49, 1940

CRATAEVA NURVALA Ham

Chemical constituents of the root bark of *Crataeva nurvala* Ham. Bhandari, P R , Dhar, M L , Shafma, V N , *J Sci Indl Res* , 2951, 10B, 105

CRINUM ASIATICUM Linn

Alkaloids of *Crinum asiaticum* L. Var *Japonicum*, Bak Kinzi Tanaka, *J Pharm Soc , Japan* 57, 652, 1937

CROCUS SATIVUS

Saffron, colouring matter Paul Karner and Harry Salomon, *Chem. Acta* 11, 711, 1928

Picrocrocin the bitter principle of Safran H E W Lutz, *Chem Abst.* 110, 1931

On the possibility of cultivation of Safran (*Crocus sativus*) in the Hyderabad state and its importance Inam-Ul-Haq & Sayeeduddin, M , *Cur Sci.*, I, (1932-33), 394

Picrocrocin, the terpene glucoside of Safran and the biogenesis of the Carotenoid Carboxylic acids Richard Kuhn and Alfred Winterstein, *Chem Abst* 161, 1934

Kashmir Saffron with methods of testing its purity K L Budhiraj, *J Ind Chemical Soc Ind and News Ed* 5, 135 1942

CROTALARIA JUNCEA Linn

A note on the occurrence of Tri-cotyledonary seedlings in *Crotalaria juncea* Linn Purkayastha, K.K *Curr Sci* , 1941, 10, 30

C MEDICAGINEA Lam

A note on breaking dormancy in *Crotalaria medicaginea*, Lamb. Satyanarayana Rao, N *Sci & Cultr* , 1947, 12, 503

C SPECTABILIS Roth

Isolation and some properties of an alkaloid from *Crotalaria spectabilis* Roth. W M Neal, L L Rusoff and C F Ahmann, *J Am. Chem Soc* 57, 2560, 1935

Crotalaria spectabilis poisoning in Louisiana livestock P O. Piercy and L L Rusoff, *J Am Vet Med. Assoc* 108, 69, 1946

Livestock poisoning by *Crotalaria spectabilis*. P.L Piercy and L.L. Rusoff, *Chem Abst* 6681, 1946.

CROTON TIGLIUM Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency XVIII Catharatics, Oleum Ricini, Oleum Tiglii, Aloe, Succies acalyphae Caines, J F., Mhaskar, K S, *Ind Jour Med Res.*, 1923, 11, 103

Poison of Croton oil Isolation of Croton resin, thin oil and Phorbol from Croton oil by alcoholysis Bonifaz Flachentrager and George Wigner, *Helv Chem. Acta* 25, 569, 1942

CRYPTOSTEGIA GRANDIFLORA R Br

Extraction of rubber from *Cryptostegia grandiflora* Bhatnagar, S S, Karimullah & Uma Shankar, *J Sci Indl Res*, 1944-45, 3, 441

Extraction of rubber from *Cryptostegia grandiflora* Bhatnagar S S, Karimullah & Uma Shankar, *J Sci. Indl Res*, 1944-45, 3, 263

Cryptostegia grandiflora R Br A war time source of vegetable rubber, Pt VIII Examination of seeds and the fatty oil. Siddiqui, R H & Warsi, S A, *Ind Jour Pharm* 1945, 7, 75

Cryptostegia grandiflora R Br Siddiqui, R H & Warsi, S A.; *Ind Jour Pharm*, 1945, 7, 117

Rubber from *Cryptostegia grandiflora* Bhatnagar, S S, Karimullah & Uma Shankar, *J Sci Indl Res*, 1945-46, 4, 654

CUBEBA OFFICINALIS Rafin

Notes on some Indian Essential Oils *Cubeba officinalis* Misc (Syn *Piper cubeba*, Linn) Rao B Sanjiva, Sudborough J J and Watson H E *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol 8A, Part X, pp 159-160

CUCUMIS SATIVUS Linn

A proteolytic enzyme in cucumber (*Cucumis sativus*) Chopra, R N & Roy, A C, *Ind Jour. Med Res*, 1933, 21, 17

Chemical examination of the seeds of *Cucumis sativus* Linn Miss Padmini Soni, Gupta, S C & Aggarwal J S, *J Sci Indl Res* 1949, 8B, 210

C UTILISSIMUS Roxb

Chemical examination of the seeds of *Cucumis utilissimus* Roxb Bhasin, M M, Gupta, A S & Aggarwal, J S, *J Sci Indl Res*, 1950 9B, 230

CUCURBITA PEPO D.C

Examination of the pumpkin oil from Pumpkin seeds The constitution of obec acid. J.L. Rubsomer and G A. Nesty, *J Am. Chem. Soc* 56, 1784, 1934.

Biological value of proteins in Watermelon and Pumpkin seeds. Tsao E King, *Chinese J Physio.* 16, 31, 1941.

The anthelmintic power of Pumpkin seeds Giuseppe Sanfilippo,
Chem Abst 1346, 193, 32.

CUCURBITACEAE

Nomenclature changes of some common plants of the Cucurbitaceae
Chakravarty, H L, *Sci & Cult*, 1949, 15, 31

CUMINUM CYMINUM Linn

Notes on some Indian Essential oils Cuminum cyminum, Linn
Rao B Sanjiva, Sudborough J J and Watson H E, *Journal of the Indian Institute of Science, Bangalore*, 1923, Vol 8A, Part X, pp. 182-183

CUPRESSUS TORULOSA Don

The constituents of some Indian Essential Oils Part XI—The Essential Oil from the leaves of Cupressus torulosa Don Simonsen J L, *Indian Forest Records*, 1922, Vol X, Part I

CURCUMA Linn

Active constituents of Curcuma E Franquela, *Chem Abst* 3833, 1934

C AMADA Roxb

Curcuma amada—S Dutt and J N Tayal, *Soap Journ.* Feb 1941, p 200

Chemical examination of the essential oil derived from the rhizomes of Curcuma amada Roxb S Dutta and Jagat Narayan Tayal, *Ind Soap J* 7, 200, 1941

C AROMATICA Salisb

Constituents of some Indian essential oils Curcuma aromatica, Puthan Madhathel, Bhaskara Panicker, B Sanjiva Rao and J L Simonson, *J Ind Inst Sci* 9A, 133, 1926

Constituents of some Indian Essential Oils Part XX - Essential Oil from the Rhizomes of Curcuma aromatica Salisb Rao B Sanjiva, Shintre V P and Simonsen, J L, *Journal of the Indian Institute of Science, Bangalore*, 1926, Vol 9A, Part VI, pp 140-144

C CAESIA Roxb

Curcuma caesia—*Proc Nat Acad Sc* Vol V, 1940, pp 64-68

C DOMESTICA Valetton

Constituents of the rhizomes of Curcuma domestica H Dieterle and Ph Kaiser, *Arch Pharm* 217, 337

C LONGA Linn

Studies in Indian Essential oils V-Essential oil from the Rhizomes of Curcuma longa, Linn Kelker N C and Rao B Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1933, Vol 17A, Part II, pp 7-24.

Essential oil from the rhizomes of *Curcuma longa* L N C Kelkar & B Sanjiva Rao, *J Ind Inst Sci* 17, 24, 1933

Indian essential oils Essential oil from the rhizomes of *Curcuma longa* N C Kelkar and B Sanjiva Rao, *J Ind Inst. Sci* 17A, 7 1933

Volatile plant constituents—Tumerone, the aromatic principle of Turmeric oil H Hupe Georges Clar Alexander St Pfau and Pl Plattner, *Helv Chim Acta* 17, 372 1934

Turmeric and vegetable oil as repellents against Anopheline mosquitoes Philip, M I, Ramkrishna, V & Rao, V V, *Ind Med Gaz* 1945, 80, 343

Pharmacological action of essential oil of *Curcuma longa*, Chopra, R N, Gupta, J C & Chopra, G S, *Ind Jour Med Res* 1941, 29, 769

C ZEDOARIA Rose

Note on some Indian Essential Oils *Curcuma Zedoaria*, Roscoe Rao B Sanjiva, Sudborough J J and Watson H E, *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol 8A, Part X, pp 153-155

The constituents of some Indian Essential Oils Part XXIV—The Essential Oil from the rhizomes of *Curcuma Zedoaria*, Roscoe Rao B Sanjiva, Shintre V P and Simonsen J L, *Journal of the Indian Institute of Science, Bangalore*, 1928, Vol IIA, Part XV—I, pp 195-199

CUSCUTA REFLEXA Roxb

Chemical examination of *Cuscuta reflexa* Roxb Radha Raman Agarwal and Sikhibhushan Dutt, *Ind J Chem Soc* 1935, 12, 384

Cuscuta reflexa Roxb Chemical examination of. Part II The constitution of cuscatalin Agarwal, Radha Raman & Dutt, Sikhibhushan *J Ind Chem Soc* 12, (1935), 586

Cuscuta reflexa Chemical examination of Part III The constitution of the oil from the seeds Agarwal Radha Raman & Dutt, Sikhibhushan, *J Ind Chem Soc* 13 (1936), 264

Cuscuta reflexa Roxb Chemical examination of Part IV Isolation of a new yellow flavone, colouring matter from the seeds. Agarwal, Radha Raman, *J Ind Chem Soc* 13 (1936), 531

Parasitism of *Cuscuta reflexa* Roxb & *Loranthus longiflorus* Desr Sheriar, K C, *Sci & Cult*, 1951, 17, 218

On *Cuscuta reflexa* Roxb in the Khasi Hills Kachroo, P, *Sci & Cult*, 1951, 16, 431

CYCAS REVOLUTA Thunb

Constituent of seed oil of *Cycas revoluta* Seuch Ueno, Sumio Matsuda and Taka Kimura, *Chem Abst* 5618, 1950. 42 7 99

CYDONIA VULGARIS Pers.

The fatty oil of Quince seed, *Cydonia vulgaris* A Steger and J Van Loon, *Chem Abst* 2207, 1934

CYMBOPOGON Hack

Cymbopogon Grasses—B S Rao and J J Sudborough, *Journal of the Indian Institute of Science, Bangalore*, Vol 8A, Part II, pp 9 to 27, 1925

Oil of a new Cymbopogon species *Bull Imp Inst* 27, 1929, 459

Studies in Indian Essential oils Essential oil from the flower heads of Cymbopogon, Kotnis M S and Rao B. Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1935, Vol. 18A, Part XVIII, pp 129-133

Cultivation of Cymbopogon grass in Trans-Gangetic tracts, J N Rakshit, *Rakshit Gardens Bulletin No 1, Ghazipur, U.P.* 1936.

Development of Essential Oil Industry Ocimum lemon, O basilicum, O canum, O sanctum, Lemon grass, Motia, Palmarosa grass Jitendra Nath Rakshit, *Sci & Cult*, 1939, 5, 108

C CAESIUS Stapf

Travancore Essential Oils Inch grass Cymbopogon caesius Stapf. Moudgill K L and Aiyar K R Krishna, *Department of Industries, Trivandrum*, 1923, *Bulletin No XVII*, pp 18-25

Travancore Essential Oils Part VI—The Essential oil from Cymbopogon caesius, Stapf Inch grass Moudgill K L., *Journal of the Indian Chemical Society, Calcutta*, 1925, Vol II

Kachi grass oil (Cymbopogon caesius). R B Rao and J J Sudborough, *J Ind Inst Sci* 8A, 9, 1925

Essential oil from Cymbopogon caesius Stapf Moudgill, *J Ind Chem Soc* 2 (1925), 23

C CITRATUS Stapf

Distillation tests on lemon grass, Cymbopogon citratus R Wilbaux; *Chem Abst* 3636, 1937

Essential Oils Lemon Grass Oil. *Department of Industries, Trivandrum, Bulletin No IV.*

C CLANDESTINUS Stapf

Oil of Cymbopogon clandestinus *Bulletin Imp. Inst.* 27, 1929 438.

C COLORATUS Stapf

The constituents of some Indian Essential Oils Part XVII—Essential Oil from the flower heads of Cymbopogon coloratus, Stapf Pillai P Parameswaran, Rao B Sanjiva and Simonsen J L.; *Journal of the Indian Institute of Science, Bangalore*, 1928, Vol 11A, Part XV—1, pp 181-186

C. MARTINI Stapf

Note on the economic uses of Rosha Grass (*Cymbopogon martini*, Stapf)—R.S Pearson, *Indian Forest Records*, 1916, Vol. V Part VII.

Some observations on the Essential oil content of Rosha Grass (*Cymbopogon martini* var. *Motia*) Girdhari Lall, *The Indian Journal of Agricultural Science*, Vol V, Part III, 1935

CYNOMARATHRUM NUTTALLII A Gray

Volatile oil and resin of *Cynomarathrum nuttallii* A Gray (or *Pencedanum graveolens*) E K Nelson, *J Am Chem Soc* 55, 3400, 1933.

CYPERUS ESCULENTUS Linn

Oil value of *Cyperus esculentus* tubers J Pieraerts, *Chem Abst* 1917, 1924

C ROTUNDUS Linn

Essential oil of *Cyperus rotundus* B S Rao, P.B Panicker and J J. Sudborough; *J Ind Inst Sci* 8A, 39, 1925

Essential oil of root of *Cyperus rotundus* L. of Japan Y Kumura and M Ohtani; *J Pharm Soc Japan*, 48, 971, 1928

Oil of *Cyperus rotundus*—B J Hedge and B S. Rao, (*Journ Soc Chem Ind* 54, 1935, T 388)

Some of the constituents of the tuber of Coqui (*Cyprus rotundus* L) Preliminary examination of the tuber and composition of the fatty oil. Conrado F Asenjo, *J Am Pharm Assoc* 30, 216, 1941

C ROTUNDUS & C SCARIOSUS Br

Chemical investigation of *Cyperus rotundus* and *Cyperus scariosus* Basu N K , *Sci & Cultr* , 1944, 10, 131

Oil of Cyperiol (*Cyprus scariosus*) Gupta, G N & Ganesh Chandra, *Curr. Sci* , 1951, 20, 273

CYSTOPUS CANDIDUS

Medicinal action of *Capsella bursa-pastoris* and also of its parasites, *Cystopus candidus* and *Peranospora parasitica*. W Harste, *Arch Pharm* 206, 133, 1938

DAEMIA EXTENSA (Linn) R Br

Pharmacological action of an active constituent isolated from *Daemia extensa* Linn. (Syn *Pergularia extensa*) Pt I. Gupta, J.C., Roy, P K & Dutta, A , *Ind. Jour Med Res* , 1946, 34, 181

Daemia extensa R Br. Datta, A. & Ghosh, S., *Ind Jour. Pharm* 1947, 9, 58.

Chemical examination of *Daemia extensa* R. Br. Ashutosh Dutta and Sudhamoy Ghosh, *J Am. Pharm. Assoc.* 36, 250, 1947.

Pharmacological action of an active constituent isolated from *Daemia extensa* Linn (Syn. *Pergularia extensa*). Pt. II. *Ind. Jour Med Res* , 1950, 38, 75.

DATURA Linn

The alkaloid content of some *Datura* species L. S Dyson and J. A. Coetzee, *S African J. Sci* 40, 162, 1943

Cultivation studies of the Solanaceous drugs Post harvest alkaloidal movement in *Belladonna* and *Datura* species. W R. Brewer and L. David Hiner, *J Am Pharm Assoc* 37, 541, 1949

D ALBA Nees

Alkaloids of *Datura alba* Nees S Osada, *J Pharm. Soc. Japan* 504, 89, 1924

Isolation and separation of main constituents of *Datura alba* Nees Cheoa Sakul Pradisth and A C Santos, *Chem Abst* 5593, 1939

Alkaloids of *Datura alba* Nees Cheoa Sakul Pradisth and Alfrendo E Santos, *Chem Abst* 767, 1940.

Isolation and separation of the main constituents of *Datura alba* Nees Cheoa Sakul Pradisth and Alfrendo E Santos, *Chem. Abst.* 3441, 1940

Fertilizer effects on the growth and alkaloidal content of *Datura alba* Nees Prasad, S , *Ind Jour Pharm.* 1944, 6, 13.

A mosaic disease of *Datura alba* Nees , Capoor, S P. A & Varma, P M , *Curr Sci* , 1949, 17, 151

D FASTUOSA Linn

Datura metel L & *D fastuosa* L Narayanswami, V *Sci & Cult* , 1948, 14, 38

Datura metel L & *Datura fastuosa* L Chatterjee, D Narayanswami, V , *Sci & Cult* , 1948, 14, 206

D INOXIA Mill.

Datura inoxia a potential commercial source of Scopolamine Goe H Garlach, *Econ Botany*, 2, 436, 1948

D METEL Linn

Datura metel L. & *D fastuosa* L. Narayanswami, V , *Sci & Cult.* 1948, 14, 38

Datura metel L & *Datura fastuosa* L Chatterjee, D., Narayanswami V., *Sci & Cultr* , 1948, 14, 206

Datura metel Chatterjee, D., *Sci & Cult* , 1949, 15, 77.

D STRAMONIUM Linn (D. TATULA Linn.)

A study of *Datura stramonium*. Frank H. Eby, Frederick. M. Scholl and David H. Phillips, *J. Am. Pharm. Assoc.* 27, 474, 1938.

The value of Palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves. Bernard S Feinstein and Frank J Slama, *J Am Pharm Assoc.* 29, 370, 1940

The effect of varying temp upon the total alkaloid content of folia stramonii and folia belladonna H Fluck, *Chem Abst.* 3878, 1940.

Synthesis of Hyoscyamine in Atropa belladonna and Datura stramonium B T. Cromwell, *Biochem Jour* 37, 717, 1944

Comparative analysis of normal and tetraploid Datura stramonium and Datura tatula. Ovrville H. Miller and Louis Fischer; *J Am Pharm Assoc.* 35, 23, 1946

DAUCUS CAROTA Linn

Essential oils of Carrot (Daucus carota) G Igolen, *Chem. Abst* 7281, 1936.

Oil of Carrot. Ernest S Guenther, *Am. Perfumer*, 5, 71, 1936.

The carotenoid pigments and the Vitamin 'A' activity of Indian carrot. Sadana, J C & Bashir Ahmad, *Ind Jour. Med. Res.* 1947, 35, 81.

DECALEPIS HAMILTONII W & A

Oil of Decalepis hamiltonii M. G Srinivasa Rao and M. Sesha Iyenger; *Perfumery Records* 14, 1923, 300

Chemical components of roots of Decalepis hamiltonii (Makali veru) Comparison with Hemidesmus indicus (Indian Sarsaparilla). P. Bhaskara Ramamurti and T R. Sheshadri, *Proc. Ind. Acad Sci* 13A, 399, 1941.

DELPHINIUM AJACIS Linn.

Alkaloids of Delphinium ajacis. Mathew V Hunter, *Chemist & Druggist*, 139, 304, 1943

D. ELATUM Linn

Alkaloids of the seeds of Delphinium elatum L Jhon. A. Goodson; *J. Chem. Soc* 139, 1943.

DERRIS Lour.

Rotenone the active constituent of Derris root. Shankichi Takiee; *Ber.* 61B, 1003, 1928

Constituents of Derris root Tubaic acid T. Karizone, K Kondo and Makdie; *J. Pharm Soc. Japan.* 48, 674, 1928.

Recent progress in the chemistry of Derris. R.C Roak, J C Con, *Entomol.* 22, 378, 1929

Rotenone the active principle of Derris root Sankichi Takel, Shakivo Mihajima and Minoru Ono, *Chem Abst* 5565, 1932

Pyrethrum and Derris insecticides as arsenical substitute. Roy. Huston; *Chem Abst.* 6234, 1934.

Some field tests showing the comparative efficiencies of Derris, Pyrethrum and Belladonna powders on different insects. Clyde C. Hamilton and Lousie G Gemelle, *J Econ. Entmol.* 27, 446, 1934.

Rotenone the non-crystalline constituent of Derris root. H L Hallerna, F B. Laforge, *J. Am Chem Soc* 56, 2415, 1934

D. ELLIPTICA Benth

Rotenone the physiologically active constituent of Derris elliptica. A. Buten, *Chem. Abst* 3660, 1928

Toxicological studies of Derris elliptica and its constituent Rotenone. H B Haag, *J. Pharmacol* 43, 193, 1931

Occurrence of Derris elliptica in India Krishna, S. & Ghosh, T.P., *Curr Sci*, 1935-36, 4, 857

D FERRUGINEA Benth.

Denis ferruginea from Assam Krishnan, S & Ghose, T P., *Curr. Sci*, 1938, 7, 22.

D SCANDENS Benth

Chemical Examination of the seeds of Derris scandens. Rao, C. J. Dassa, Subramanian, S. S *Curr Sci.*, 1947, 16, 346.

DICHANTHIUM ANNULATUM Stapf

Bulk emasculation technique as applied to Dichanthium annulatum Stapf Oke, J G., *Sci & Cult*, 1950, 16, 30

DICHROA FEBRIFUGA Lour

Chang Shan—a Chinese antimalarial herb, Dichroa febrifuga C. S. Jang, F. Y Fu, C Y Wang, K C. Haung, G Lu and T C Chou, *Science*, 103, 50, 1946.

Alkaloids with high antimalarial activity from Dichroa febrifuga J B Koepfli, J. F Mead and John A Brockman, *J. Am. Chem. Soc* 69, 1837, 1947

Alkaloids of Dichroa febrifuga. Frederick, A, Kuehl. Jr Claude F. Spencer and Karl Folkers, *J Am Chem. Soc* 70, 2091, 1948

Pharmacology of Chan Shang (Dichro febrifuga), a Chinese anti-malarial herb. C. S Jang and F. Y. Fu and K C Haang and C. Y Wang, *Nature*, 161, 400, 1948

Alkaloids of Dichroa febrifuga. Isolation and degradation studies. J B Koepfli J F Mead and John A Brockman, *J Am Chem Soc* 71, 1048, 1949

DICTAMUS FRAXINELLA Pers •

Oil from the leaves and flowers of Dictamus fraxinella Pers B N. Rutovskii and I V Vingradova, *Chem Abst* 3774, 1926

DIDYMOCARPUS PEDICELLATA Br

Didymocarpus pedicellata. The constituents of Part I Isolation of a new series of colouring matters. Siddiqui, Salimuzzaman; *J Ind Chem. Soc* 14 (1937), 703

Didymocarpus pedicellata. The constituents of. Part II. Comparative studies in the constitution of Pedicine, isopedicin, pedicinin & Pedicellin Sharma, Vishwanath & Siddiqui, Salimuzzaman, *J Ind Chem Soc* 16 (1939)

Constitution of *Didymocarpus pedicellata* Pt V. Studies in the constitution of Pedicinin Kailash Chander Saloaja, Vishwa Nath Sharma & Salimuzzaman, Siddiqui, *J Sci Ind Res*, 1947, **6B**, 57.

DIGITALIS (Tourn) Linn

Physiological standardization of Digitalis with special reference to the method suitable for use in India Tate, G, *Ind Med Gaz*, 1921, 56, 505

Variation in the potency of Digitalis preparations Chopra R N, Bose, S C & De P., *Ind. Med. Gaz*, 1925, 60, 93

Clinical studies on Digitalis in Bengal, Bose, S C., *Ind Med Gaz*, 1925, 60, 147

Variations in the potency of Digitalis preparations in tropics Chopra, R. N, De, Premankur, *Ind. Med. Gaz*, 1926, 61, 57

Observations on the potency of Indian Digitalis Chopra, R N, De, Premankur, *Ind Med Gaz*, 1929, 64, 312

Biological assay of Digitalis preparations in the tropics, Pt III Chopra, R N & Chowhan, J S., *Ind Jour. Med. Res.*, 1933, 20, 1189

The value of palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves. Bernard S Einstein & Frank J Slame, *J Am Pharm Assoc* 29, 370, 1940

Combined Digitalis and Rauwolfia poisoning in a human subject De, M M & Tarapaido Chatterjee, *Ind Med. Gaz*, 1941, 76, 724

Digitalis, their chemical evaluation Notes & News, *Ind Jour. Pharm*, 1946, 8, 50

D LANATA Ehrh

A study on Digitalis lanata Ehrh S C Grown in India Bose, A N & Bose, S, *Ind Jour Pharm*, 1952, 14, 189

DIOSCOREA HISPIDA Dennst

Starch from the tubers of Dioscorea hispida Dennst Rao, P S & Beri, R M, *Sci & Cult*, 1952, 17, 482

DIOSPYROS EMBRYOPTERIS Pers.

Pectin in fruits of the Indian Persimmon (D embryopteris Pers) Biswas, H G., *Sci & Cultr*, 1944, 9, 501.

DIPTEROCARPUS INDICUS Bedd

On the essential oil from the Oleo-resin of Dipterocarpus indicus Krishna G S, Rao, Sukh Dev & Guha, P C, *Sci & Cult.*, 1950, 16, 74

Studies in sesquiterpenes Part VIII Sesquiterpene of the essential oil from *Dipterocarpus indicus*, Bedd Krishna Rao, G S , Sukh Dev & Guha, P C , *J Ind Chem Soc* 29 (1952) 589

DODONAEA VISCOSA Linn

Chemical examination of *Dodonaea viscosa* Linn T P Ghosh, *Indian Forester*, 59, 78, 1933

Chemical examination of the seeds *Dodonaea viscosa*, isolation of dodonin and a fixed oil Dharmpal Parihar and Shikhibhushan Dutt, *Pro Ind Acad. Sci* 26A, 56, 1947

DOLICHOS LABLAB Linn

Oil of *Dolichos lablab* B S Rao and Co-workers , *Perfum Record*, 28, 1937, 413

DROSERA INDICA Linn

A new variety of *Drosera indica* Linn from Kolhapur (S M C) Parandekar, S A , Diwan M G , *Curr Sci* , 1943, 12, 304

DRYOPTERIS

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Enquiry in the Madras Presidency XI Extractum Filicis Liquidum Cains, J F & Mhaskar, K S , *Ind Jour Med Res* , 1921 9, 191

Constituents of *Filexmas*, Part I, Aspidinol. Alexander Robertson and William F Sandroock , *J Chem Soc* 819, 1933

Male Ferns of Kashmir Handa, K L , Kapoor, L D & Chopra I C , *Curr Sci* 1947, 16, 56

A short note on Male-ferns Handa, K L , Kapoor L D & Het Singh, *Ind. Jour. Pharm* 1952, 14, 109

EICHHORNIA Kunth.

Studies in the lignacellulose group Part I An investigation into the constituents of Water Hyacinth (*Eichhornia crassipes*). Hemendra Kumar Sen, Pal, Patitpavan Ghosh, and Sindhu Bhushan., *J. Ind Chem Soc* 6 (1929), 673

A Pamphlet on Water-Hyacinth (Science & Industry Sec) *Sci & Cult* , 1942, 7, 602

The problem of Water-Hyacinth in Bengal Bose, P K., *Sci & Cultr* , 1945, 11, 167

Eradication of Water-Hyacinth by Methoyone on a field scale trial. Mitra, G.P., *Sci & Cult.*, 1948, 14, 143

ELAEIS GUINEENSIS Jacq

Red Palm oil in the treatment of Keratomalacia. Aykroyd. W.R. & Wright, R.E , *Ind. Jour Med Res* , 1937, 25, 7.

ELETTARIA CARDAMOMUM Maton.

Notes on some Indian Essential Oils "Elettaria cardamomum", Maton. (Var a-minor) Rao, B Sanjiva, Sudborough, J J and Watson, H E., *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. 8A, Part X, pp 155-158

Mysore cardamon oil Krishnan, P P & Guha, P C , *Curr Sci* , 1950, 19, 158

ELEUCINE CORACANA Gaertn

The nutritive value of Indian vegetable foodstuffs, Pt V. Nutritive value of Ragi (Eleucine coracana) Niyogi S P , Narayana, N. & Desai, B G , *Ind Med Res* , 1934, 22, 373

The relative value of the proteins of certain food-stuffs in nutrition Pt. V Supplementary values of the proteins of Eleucine coracana (Ragi) and of certain pulses and stem milk powder studied by the nitrogen balance and certain growth methods Swaminathan, M , *Ind Jour Med Res* , 1938, 26, 107

Eleucine coracana Gaertn A new raw material for the malting Industry Sastri, B N Ragi , *Curr Sci* , 1939, 8, 34

EMBELIA Burm.

The treatment of intestinal worms with the indigenous drugs Butea, Embelia and Kamala Mukerji, A K , Bhaduri, N V , *Ind Med Gaz* , 1947, 82, 66

E. RIBES Burm

Constitution of Embelia ribes Part I Kaul, Banerjee, Amarish Chandra Roy, Dutt, & Sikhishushan , *J Ind Chem Soc.* 6 (1928), 577

Embelia ribes Constitution of the active principles of. Part II Kaul, Ramjee, Amarish Chandra Roy, Dutt, & Sikhishushan, *J Ind. Chem Soc* 8 (1931) 231

EMBLICA OFFICINALIS Gaertn

Chemical Examination of the seeds of Emblica officinalis Pt I , The fatty oil and its component fatty acids Dhar, D.C., Dhar, M L. & Srivastava, D L., *J. Sci Indl Res.*, 1951, 10B, 88.

ENHYDRA FLUCTUANS Lour.

Chemical Investigation of Enhydra fluctuans Lour Part I. Chakravarty, R.N. & Dutt, A., *J. Ind Chem Soc* 29 (1952) 374.

ENICOSTEMMA LITTORALE Blume.

Enicostemma littorale Blume in malaria. Rai, B.B., *Ind. Med. Gaz.*, 1946, 81, 506

ENTADA PURSAETHA D.C

A note on the chemistry and pharmacological action of *Entada pursaetha* DC (*E. scandens* Benth) Chopra, R.N, Gupta, J.C, Chopra, G.S & Ghosh, B K., *Ind. Jour Med Res.*, 1940, 28, 469

EPHEDRA Linn

The comparative action of Ephedrine and Pseudoephedrine from Indian varieties of *Ephedra* on heart Chopra, R N, Dikshit, B B & Pillai, K V., *Ind. Jour. Med. Res.*, 1928, 16, 780.

Ephedrine from Indian varieties of *Ephedra*. Chopra, R.N, Ghosh, S., & Dutt, A. T, *Ind Jour. Med. Res*, 1928, 15, 889

A preliminary note on the action and uses of Pseudoephedrine Chopra, R N, Dikshit, B. B. & Pillai, K V, *Ind Med. Gaz*, 1929, 64, 1

Seasonal variations in the alkaloidal contents of Indian *Ephedra* Chopra, R N & Dutta, A T, *Ind Jour. Med Res.*, 1930, 17, 647

Indian *Ephedra*, their Chemistry and Pharmacology Chopra, R N., Krishna, S & Ghosh, T. P., *Ind Jour Med Res*, 1931, 19, 177.

Ephedra and ephedrine in U S S R. *Chem Abst* 6765, 1940.

Indian *Ephedra* research creates demand (Medicine & Public Health), *Sci & Cultr*, 1940, 6, 174.

Ephedra concentrate and the extraction of ephedrine (Science & Industry Sec.), *Sci & Cultr*, 1943, 9, 195.

Indian *Ephedras*, N A Qazilbash, *Quart J. Pharm. Pharmacol* 21, 475, 1948.

Ephedra, Pt. I Estimation of total alkld solvent extraction. Datal, V D & Khorana, M L, *Ind Jour. Pharm.*, 1950, 12, 165.

Pharmacognosy of Indian *Ephedra*. Gupta, B. & Bal, S. N, *J Sci. Indl Res.*, 1952, 11B, 253.

E NEBRODENSIS Tin

Medicinal flora of Sardinia. *Ephedra vulgaris* Rich. and *Ephedra nebrodensis* Tin. Aldo La Floresta, *Arch Pharmacol. Sper.* 69, 67, 1940.

Pharmacological activity of ephedrine hydrochloride prepared from *Ephedra nebrodensis* from the Oliena mountains. Sante Gabatto, *Chem. Abst* 6209, 1946.

E. VULGARIS RICH.

Medicinal flora of Sardinia, *Ephedra vulgaris* Rich. and *Ephedra nebrodensis* Tin. Aldo La Florista, *Arch. Pharmacol. Sper.* 69, 67, 1940.

The Ephedras of Sardinia Chemical investigation of *Ephedra vulgaris* Rich. *Chem. Abst.* 8204, 1941

ERIGERIN CANADENSIS Linn

The essential oil of *Erigerin canadensis* L T K. Geponekov; *J. Applied Chem.* (U.S.S.R.) 6, 111, 1933.

ERYTHRINA INDICA Lam.

Chemical examination of *Erythrina indica* (White variety) Rao, J Virabhadra, *Curr. Sci* , 1945, 14, 198

Some common indigenous remedies *Picrorhiza kurrooa*, *Erythrina indica*, *Sansevieria zeylanica*, *Pongamia glabra*, *Hygrophila spinosa*, *Bryophyllum calycinum*, *Rheum emodi*, *Solanum indicum* R. N. Chopra & S Ghosh; *Ind Med Record* 55, 77, 1935

ERYTHROXYLUM MONOGYNUM Roxb

The Essential Oil from the wood of *Erythroxylum monogynum* Roxb Rao B Sanjiva, Shintre V P and Simonsen J L , *Journal of the Indian Institute of Science, Bangalore*, 1926, Vol. 9A, Part VI, pp. 145-148.

Devadani - root oil B. S Rao and Co-workers, *Perfum. Record*, 28, 1937, 412.

Devadani wood oil S G. Sastry, *Journal of the Mysore Forest Association*, Vol 5, No 4.

EUCALYPTUS Lher

Note on the Eucalyptus Oil industry in the Nilgiris Puran Singh, *Indian Forest Records*; 1917, Vol. V Part VIII

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency VII *Oleum Eucalypti* Cains, J F & Mhaskar, K S , *Ind Jour Med. Res* , 1920, 7, 372

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency IX The Eucalyptus CHCL₃ mixture S C Cains, J F. & Mhaskar, K. S., *Ind Jour Med Res.*, 1920, 8, 384.

E. CITRIODARA Hook.

Leaves of *Eucalyptus citriodara* yielded 1.3 per cent oil. Noemi Carcia Arrillaga *Chem. Abst* 5615, 1942.

E. GLOBULUS Labill.

The essence of *Eucalyptus globulus* Martias Gonzales Atilio Lambardo and M A Mesa De Romero, *Chem. Abst* 4666, 1942.

Essential oil of leaves of *Eucalyptus globulus* from groves in La Plata. Leonor Pelandia Ponce, *Chem. Abst* 5955, 1942.

Essential oil of *Eucalyptus globulus*. B. V. Ramaswami, P. L. Narasimha Rao and P. C. Guha; *J Ind. Inst. Sci.* **28A**, 57, 1946.

Microscopical studies of the leaves of *Eucalyptus globulus* Labil. and three other species grown in India Amiya Datta & Datta, D, *J. Sci Indl. Res.*, 1952, **11B**, 181

EUGENIA JAMBOLANA Lam.

Eugenia jambolana Subba Rao, N. V & Seshadri, T R, *Curr Sci* 1939, 8, 120.

Jaman (Black Berry) seed as a cattle feed Kehar, N D. & Sahai, K, *Sci & Cult*, 1948, 14, 205.

EULOPHIA R Br

On the occurrence of velamen in *Eulophia* R Br Lakshminarayana, S & Venkateswarlu, V, *Sci & Cult*, 1950, 15, 327.

EUPHORBIA Linn.

Triterpenic compounds isolated from *Euphorbia* latices. Gopalachari, R. & Siddiqui, S, *J. Sci Indl Res*, 1949, **8B**, 234

E. GENICULATA Ort.

On the introduction of *Euphorbia geniculata* Ort. in India. Van Steenis, C G G J, *Curr Sci.*, 1937-38, 6, 119.

Fresh locality record for *Euphorbia geniculata* Ort in South India Mayuranathan, P V, *Curr. Sci*, 1937-38, 6, 119.

E. HELIOSCOPIA Linn

Seeds of *Euphorbia helioscopia* L Paul Gillot, *Bull Sci Pharmacol.* 33, 193, 1926

E PEPLIS Linn

Chemical investigation of Egyptian plant *Euphorbia peplis*, Abd El Aziz Sharaf, *British Vet J* 105, 128, 1949

E PILULIFERA Linn

Pharmacognostic study of *Euphorbia pilulifera* Linn. Bose, A B., *Sci. & Cult.*, 1944, 9, 301.

Pharmacology of *Euphorbia pilulifera* Llyod W. Hazelton and Rebecca C Hellerman; *J Am. Pharm. Assoc* 37, 491, 1948.

The isolation of L-inositol from *Euphorbia pilulifera*. Floyd P. Hallet and Lloyd M. Parks *J. Am. Pharm. Assoc.* 40, 474, 1951.

E THYMIFOLIA Linn.

Flavonol glucosides from *Euphorbia thymifolia* L. Makoto Nagase; *Chem. Abst.* 3625, 1942.

E. TIRUCALLI Linn.

Chemical examination of dried latex from *Euphorbia tirucalli*. Dutta, N. L. & Karimullah, *J. Sci Indl Res.*, 3, 213, 1944-45.

Chemical Examination of the dried latex from *Euphorbia tirucalli* Karimullah & Gopalachari, R, *J Sci Indl Res.*, 1949, **8B**, 89.

Chemical Examination of the latex from *Euphorbia tirucalli* Gopalachari, R & Siddiqui, S, *J Sci. Indl Res.*, 1949, **8B**, 129.

Studies in the const. of Euphoron Pt I (*Euphorbia tirucalli*). Gopalachari, R & Siddiqui S, *J. Sci Indl. Res.*, 1949, **8B**, 140.

FAGOPYRUM CYMOSUM Meissn

Phytochemical component of *Fagopyrum cymosum*. Kaneo Imai and Kiyoshi Furuya, *J. Pharm Soc Japan*, 71, 266, 1951

FERONIA ELEPHANTUM Correa.

Feronia elephantum Essential oil from the leaves of Bhatia & Deshpande, *J. Ind Chem Soc* 26(1949), 342.

Investigation on the structure of Ketha (*Feronia elephantum*, family Rutaceae) Gum Mathur, G P., Mukerjee, S, *J. Sci Indl Res*, 1952, **11B**, 344

FERULA Linn.

Chemical composition of *Ferula* spp O M. Efimenko, *J Applied Chem.* (U. S. S R.) 12, 202, 1939.

Chemical assay of Rasaunt and Hing from the Punjab market Grewal, K. S., Kochhar, B. D, *Ind. Jour. Med. Res.* 1940, 28, 463.

F. FOETIDA Regel.

Studies in the specification of Indian Medicinal Plants. Pt I Asafoetida, *Ferula foetida*. Sen Gupta, S B & Das, B. K, *Ind Jour Pharm.*, 1948, 10, 36

F. JAECHKEANA Vatke.

Chemical investigation of the fruits of *Ferula jaeckkeana* Vatke V. P Bersutskii, *Chem Abst* 4522, 1940

FICUS ANTHELMINTICA Martius.

Note on *Ficus indica* Linn. and closely allied American species. *Ficus laurifolia* Hort. et Lam. and *Ficus anthelmintica* Martius. *Curr. Sci.*, (1933-34), 2, 51

F. BENGALENSIS Linn.

Fasciated aerial roots of Banyan & (*Ficus bengalensis* Linn.). Nadkarni, M D., Airan, J. W, Shah, S V., *Curr. Sci.*, 1944, 13, 233.

Investigation on "Ficosterol", a sterol occurring in *Ficus bengalensis*. Nath, M.C. & Debnath, C R.; *Sci & Cultr.*, 1947, 12, 599.

F. CARICA Linn.

The constituents of the leaves of *Ficus carica*. Kunio Okahara, *Bull. Chem Soc. Japan*, 11, 389, 1936

Some pharmacological and biological effects of the latex of *Ficus carica* L. S. B. Ullman, L. Helberstaedter and J. Leibowitz, *Chem. Abst* 3071, 1945, 530

Preparation of a highly active vegetable Rennet from *Ficus carica* Linn Krishnamurthy, C. R., Jaganathan, V. & Subrahmanyam V., *J. Sci. Indl. Res.*, 1945-46, 4, 720

The proteolytic enzyme of the latex of *Ficus carica* Linn. Krishnamurti, C. R. & Subrahmanyam, V., *Sci. & Culture*, 1947, 13, 204.

F. INDICA Linn & F. laurifolia Hort et Lam

Note on *Ficus indica* Linn and closely allied American species—*Ficus laurifolia* Hort et Lam and *Ficus anthelmintica* Martius. *Curr. Sci.*, (1933-34), 2, 51

F. RELIGIOSA Linn

Fascination in an anomalous foliar organ of *Ficus religiosa* Linn De Sarkar, K. & Datta, R. M., *Sci. & Culture*, 1952, 17, 381

FOENICULUM PANMORIUM D. C.

Notes on some Indian Essential oils *Foeniculum panmorium* D. C. (Fennel) Rao B. Sanjiva, Sudborough, J. J. and Watson, H. E., *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. 8A, Part X, pp. 184-186.

FRITILLARIA ROYLEI Hook

Some minor alkaloids of *Fritillaria roylei* T. Q. Chou, *J. Pharm. Assoc.* 36, 215, 1947

FUMARIA OFFICINALIS Linn

Chemical examination of some Indian medicinal plants, *Tinospora cordifolia*, *Solanum xanthocarpum* and *Fumaria officinalis* Pense G. P. & Dutt, S., *Ind. Jour. Med. Res.* 1932, 20, 663

The alkaloids of Fumariaceae plants *Fumaria officinalis* Richard H. F. Manske, *Can. J. Research*, 16B, 438, 1938

Action of Fumarine (*Fumaria officinalis*) Wilhelm Bolm. *Arch. Exptl. Path. Pharmacol.* 195, 304, 1940

FUSARIUM.

Antibacterial properties of Yeasts—*Fusarium* species, onion and garlic. C. Carpenter, *Chem. Abst.* 1558, 1946

GARCINIA INDICA Choisy.

Garcinia indica. Fatty acids and glycerides of the fat from the seeds of Vidyarthi, N. L. & Desa Rao, C. J.; *J. Ind. Chem. Soc.* 16 (1939), 437.

G. MANGOSTANA Linn

Resin of *Garcinia mangostana*. L. Otto. Dragendorff; *Ann.* 482, 280, 1930.

Resin of *Garcinia mangostana*. Otto Dragendorff, *Ann* 487, 62, 1931

G. MORELLA Desr

Indian seed fats. Mowha (*Bassia Latifolia*) and Tamal (*Garcinia morella*) fats. D R. Dhingra, G.L Seth & P. C Speers, *J Soc Chem Ind* 52, 116, 1933.

On 'Morellin' the antibacterial principle of the seeds of *Garcinia morella*. Rao. R., Raghunandana & Natarajan, S, *Curr Sci*, 1950, 19, 59

Antibiotic principles of *Garcinia morella* Pt. I Preparation and antibacterial activity of Morellin, Morellin-t, Morellin M, Morellin L & Iso morellin Narsimba Rao, P L & Verma, S C. L, *J Sci Indl Res*, 1951, **10B**, 184

Antibiotic principles of *Garcinia morella* Part II. Chemistry of Morellin Narasimba Rao, P L & Verma, S C L, *J Sci Indl Res* 1952, **11B**, 206

GAULTHERIA FRAGRANTISSIMA Wall

A pharmacological study of *Gaultheria fragrantissima* S Prasad, *Ind. J. Pharm* 3, 62, 1941.

Detection of Scopoletin in radix gelsemi and radix belladonnanae Robert Fischer and Herbert Ehrlich, *Arch Pharm.* 274, 268

G PROCUMBENS.

Manufacture of Wintergreen oil in India—Puran Singh, *Indian Forest Records*, 1917, Vol V Part VIII

GENTIANA KURROO Royle

Comparative pharmacognosy of *Gentiana kurroo* Royle. & *Picrorrhiza kurroa* Royle ex Benth Bal S. N & Datta, S. C., *Ind Jour. Pharm.* 1945, 7, 109.

GERANIUM MACRORRHIZUM Linn

Oil from *Geranium macrorrhizum*. Schimel and Co, *Chem Zentr* 1519, 1927.

GLORIOSA SUPERBA Linn.

A note on the distribution of *Gloriosa superba* Linn. Sayeeduddin, M & Salam, M Abdusa, *Curr Sci.*, 1934-35, 3, 443.

Alkaloidal constituents of *Gloriosa superba* Linn. Sulbaratnan A V, *J. Sci. Indl Res.*, 1952, **11B**, 446.

GLYCOSMIS PENTAPHYLLA Correa.

Alkaloids from *Glycosmis pentaphylla* Correa, Chatterjee, A. & Majumdar, S. G., *Sci. & Cult.*, 1952, 17, 306.

Glycosin, the new alkaloid of *Glycosmis pentaphylla* Correa. Asima Chatterjee & Majumdar, S. G , *Sci & Cult.* 1953, 10, 506.

GLYCYRRHIZA GLABRA Linn.

The flavanone glucoside of *Glycyrrhiza glabra* Var. *glandulifera*. Regel et Herder, Junzo Shinada and Seuchi Ueda, *Ber* **67B**, 434, 1934

A study of vehicles for medicines The *Glycyrrhiza* vehicles. Bernard Fantus, H A Dyniewicz and J. M Dyniewicz, *J. Am Pharm Assoc* 23,9 15, 1934

Sweet constituents of liquorice root Tatu Kariyone and O Nonaka, *J Pharm Soc Japan*, 57, 166, 1937.

GOSSYPIUM HERBACEUM Linn.

Constitution of Herbacetin and Herbacetin Neelkantam, K & Seshadri, T R , *Curr Sci.*, 1936-37, 5, 476

GYMNEA AURANTIAECUM Wall

Chemical Examination of Meda (*Gymnema aurantiacum*) Tubers Mukherjee, S & Srivastava, H C , *J Sci Indl Res* , 1951, **10B**, 324.

G SYLVESTR A Br

Gymnema sylvestra in Diabetes Mellitus Chopra, R N., Bose, J P , Chatterjee, N R., *Ind Jour Med. Res* , 1928, 16, 115

HAGENIA ABYSSINICA J. F Gmel

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency XII Kouso Cains, J F & Mhasakar, K S , *Ind Jour Med Res* , 1921, 9, 198

HARDWICKIA PENNATA Roxb

The oleo-resin from *Hardwickia pinnata* K S Iyer and J J Sudborough, *Journal of the Indian Institute of Science, Bangalore*, Vol. 2, Part III, pp. 29 to 35, 1918

Hardwickia pinnata Essential oil from the oleo-resin of Dev & Guha, *J. Ind Chem Soc.* 25 (1948), 495

Humulene, presence of, in oleo resin from *Hardwickia pinnata* Dev & Guha, *J. Ind Chem Soc.* 25 (1948), 500

HELIANTHUS ANNUUS Linn.

A rapid method for determining the oil content of Safflower and sunflower seeds. W Keith Kennedy and John Urau, *Chem. Abst.* 5209, 1949

HELMINTHOSPORIUM TETRAMERA Mckinney.

A preliminary note on the isolation and pathogenicity of *Aerothecium lanatum* Wakker and *Helminthosporium tetramera* Mckinney on the leaves of *Oryza sativa* Linn. Ganguly, A.K , Ganguly, D.D. *Sci & Cultr* , 1941, 6, 424.

HEMIDESMUS INDICUS Br.

Chemical examination of the roots of *Hemidesmus indicus* A T Dutt, S Ghosh and R N Chopra, *Arch. Pharm.* 276, 33, 1938

Preparation of the essential oil from the root of *Hemidesmus indicus* (Sarsaparilla) B. Sanjiva Rao, K.S Subhramanian and N C Kelkar, *Proc Soc Biol Chemists India*, 3, 35, 1938

Ether soluble constituents of Sarsaparella root. James C E Simpson and Norman E Williams, *J Chem Soc* 2040, 1939

HERACLEUM NEPALENSE D. Don

Crystalline components of the seeds of *Heracleum nepalense* Chandranath Bhar *Sci & Cult*, 1947, 12, 504

Heracleum nepalense Crystalline components of the seeds of. Bhar. *J Ind. Chem Soc* 25(1948), 139

HERPESTIS MONNIERIA H B.K

The Chemical investigation of the leaves of *Herpestis monniera* Basu, N K & Walia, J S, *Ind Jour Pharm*, 1944, 6, 84

Correct name of *Herpestis monniera* H B K Chatterjee, D., *Sci & Cult*, 1948, 14, 156

HIBISCUS CANABINUS Linn

The colouring matter of Deccan Hemp (*Hibiscus cannabinus*) flowers. Cannaluscitrin and Cannabiscetin. Neelakantam, K, Seshadri, T R, *Curr Sci*, 1937-38, 6, 504.

H ESCULENTUS Linn

The biological value of protein of papaya (*Carica papaya*) and Lady's fingers *Exptl Med* 2, 71, 1942

Availability of Calcium in Lady's finger (*Hibiscus esculentus*), cabbage (*Brassica oleracea* var. capitata), Drumstick (*Moringa oleifera*) and Amarnath Tender (*Amaranthus gangeticus*) Pt. I. Availability of Calcium in vegetables determined by experiments on growing rats. Basu, K.P & Ghosh, D., *Ind Jour. Med Res.*, 1943, 31, 29

Availability of calcium in Lady's finger (*Hibiscus esculentus*), cabbage (*Brassica oleracea* var capitata), Drumstick (*Moringa oleifera*) & Amaranth Tender (*Amaranthus gangeticus*). Pt. II Availability of calcium in vegetables determined by metabolism experiments on a human adult Basu, K P & Ghosh, D., *Ind Jour Med Res*, 1943, 31, 37.

Colouring matter of the flowers of *Hibiscus esculentus* Seshadri, T R. & Viswanathan, N., *Curr. Sci.* 1947, 16, 343.

H SABDARIFFA Linn.

Sida sabdariffa (*Hibiscus sabdariffa*) Henri Leeleve; *Chem. Abst.* 3891, 1939.

The chemical, botanical and pharmacological characters of the Karkade (Resella) *Hibiscus sabdariffa*. G. Reabourg and R.H. Moneeu, *J Pharm Chem.* 292, 1940.

Isolation of Hibiscitrin from the flowers of *Hibiscus sabdariffa*. Constitution of Hibiscetin P Surya Prakash Rao and T R Seshadri, *Proc. Indian. Acad Sci* **15A**, 148, 1942.

Constitution of Hibiscetin. Suryaprakasa Rao, *Curr Sci* 1942, 11, 360.

H. SYRIACUS Linn

A note on the Pistillody in *Hibiscus syriacus* Linn Rao, L. Narayana, *Curr. Sci*, 1936-37, 5, 301

H VITIFOLIUS Linn.

Colouring matter of the flowers of *Hibiscus vitifolius* K. Visweswara Rao and T R. Seshadri, *Proc Ind Acad Sci* **24A**, 352, 1946.

HOLARRHENA ANTIDYSENTERICA Wall

Conessine Darab Dinsha Kanga, Panch Nanda Ramaswamy Iyer & John Lionel Simonsen, *J. Chem. Soc* 2123, 1926

Oil from the seeds of *Holarrhena antidysenterica* Ramachandra R V, Ghanikar and P R Ayyar, *J. Ind Inst Sci* **10A**, 20, 1927.

Observations on the pharmacological action of conessine, the alkaloid of *Holarrhena antidysenterica* Chopra, R N, Gupta, J C., David, J C & Ghosh, S, *Ind Med Gaz*, 1927, 62, 132

The use of a standardized preparation of the total alkaloids of Kurchi Bark in Amoebic dysentery Majumdar, A R, *Ind. Med Gaz*, 1930, 65, 80

New alkaloids from *Holarrhena antidysenterica* seeds Robert D Haworth, *J Chem Soc* 631, 1933

Pharmacological action of Kurchicine (an alkaloid of *Holarrhena antidysenterica*) Chopra, R N & Chowhan, J S, *Ind Jour Med Res*, 1933, 21, 277

The treatment of chronic intestinal amoebiasis with the alkaloid of *Holarrhena antidysenterica* (Kurchi) Acton, H W, Chopra R N, *Ind Med Gaz.* 1933, 68, 6.

A study of the preparations of an efficient extract of Kurchi *Holarrhena antidysenterica* Ghosh, A B, *Ind Med Gaz*, 1933, 68, 12

Some new bases from *Holarrhena antidysenterica* Alfred Bertho, Gustav Von Schuckmann and Walter Sitionberger, *Ber.* **66B**, 786, 1933

The risoles of the latex of *Holarrhena antidysenterica*. J.C. Choudhary and D H Peacock, *J Chem Soc.* 1129, 1935

Kurchi alkaloids. Preparation of conessine and its secondary alkaloids. Alfred Bertho, *Arch. Pharm.* 277, 237, 1939.

The Alkaloids of Kurchee Bark (*Holarrhena antidysenterica*) Part I. Ghosh, Sudhamony & Ghosh, Nagendra Nath, *J. Ind Chem Soc.* 5 (1928), 477.

Holarrhena antidysenterica The alkaloids of. Part I Three new alkaloids from the bark of Indian *Holarrhena* and new method of isolation and further purification of conessine Siddiqui, S & Pillai, *J Ind. Chem. Soc* 9(1932), 553

Holarrhena antidysenterica. The alkaloids of Part II. Two further new alkaloids from the bark and the seeds of Indian *Holarrhena* and their constitutional relationship to conessine. Salimuzzman Siddiqui, *J Ind Chem Soc* 11, 1934, 283

The preliminary pharmacological studies of Kurchi bark Madhev Lal Schroff, S.N Bal and Mohan Lal Dhir, *Ind J. Pharm* 2, 195, 1940

Pharmacological tests on extract and alkaloids of *Holarrhena antidysenterica*. Alfred Bertho, *Arch Exptl Path Pharmacol* 203, 34, 1944

Studies in the Conessine series, Pt IV Action of sulphuric acid on conessine and its conversion to a new isomer Neo-conessine Salimuzzaman Siddiqui & Shyam Kishan Vasisht, *J Sci. Indl Res.*, 1944-45, 3, 559.

Pharmacognostic studies on Kurchi Bark. Datta, S C & Bal, S N., *Ind. Jour Pharm*, 1945, 113, 7

On the assay of alkaloid in Kurchi Bismuth iodide. Mukerjee, S & Dutta, B C., *Sci & Cultr*, 1945, 10, 506

Studies in the conessine series, Pt 7 Salimuzzaman Siddiqui & Vishwa Nath Sharma, *J. Sci Indl Res.* 1945-46, 4, 435.

Studies in the Conessine series, Pt 8 Salimuzzaman Siddiqui & Shyam Kishan Vasisht, *J. Sci Indl Res.*, 1945-46, 4, 440

Chemistry of Kurchi Seeds Pt I Isolation of a crystalline Glycoalkaloid, Irani, (Miss) R.J., *Curr Sci*, 1946, 15, 106.

Chemistry of Kurchi Seeds Pt. II Isolation of the Bromide of a Linoleo-Linolenin from the fatty oil Irani, (Miss) R.J, *Curr Sci.*, 1946, 15, 161.

Chemistry of Kurchi Seeds Pt. III A new & simple method of analysis by Bromoglycerides. Irani, (Miss) R J, *Curr. Sci.* 1946, 15, 191

Chemistry of Kurchi Seeds, Pt, IV Isolation of Galactose from the Picric acid hydrolysis of glyco alkaloid Irani, (Miss) R.J, *Curr. Sci.*, 1946, 15, 229.

Assay of extract of Kurchi Liq Basu, N.K. & Mithal, B.M., *Ind. Jour Pharm.*, 1947, 9, 118.

Assay of Kurchi Bark [Basu, N K & Mithal, B M., *Ind. Jour. Pharm.* 1948, 10, 72.

A method for the assay of Kurchi Bismuth Iodide. Rao, G.K, *Ind. Jour. Pharm.*, 1948, 10, 100.

The decomp of Kurchi alkaloids. Sen Gupta, S.B., Dutta, H.B.; *Sci. & Cult*, 1949, 15, 32.

Holarrhena antidysenterica Wall., estimation of total alkaloids in. Datta, A, Ghosh, B K., *Ind Jour Pharm* 1949, 11, 74

Kurchi bases, labile nature of Basu, N.K. & Battacharya, N N., *Ind Jour Pharm*, 1949, 11, 157.

Studies in the conessine series. Pt. 10 Action of fuming Nitric acid (HNO₃) on conessine Vishwa Nath Sharma & Siddiqui, S., *J Sci. Indl Res*, 1950, 9B, 84.

The seasonal variation of alkaloids in different parts of Holarrhena antidysenterica Wall. and the alkaloidal content in the bark at different ages Datta, A T Ghosh, B K & Gupta, J C, *Ind Jour. Med Res.*, 1950, 38, 467.

Thermolabile nature of Kurchi Bases Thomas Oommen, P., *Ind. Jour Pharm*, 1951, 13, 64.

H. FEBRIFUGA Klotsch

Chemical Examination of Holarrhena febrifuga Klotztsch, Pt I. Salimuzzaman Siddiqui, Shambhu Charan Misra & Viswa Nath Sharma, *J Sci Indl. Res*, 1944-45, 3, 555

HOLIGARNA ARNOTTIANA Hook. f.

Chemical Examination of the constituent of Holigarna arnottiana Hook f. Pt. I latex of Semecarpus travancorica Bed. Pt. II Nair, A.V, Poti, D M & Pillay, P P, *J. Sci Indl Res*, 1952, 11B, 294, 298.

Holigarna lactone—A new aldehydelactone from the seed kernels of Holigarna arnottiana Hook f Nair G V, Poti A N., Pillay P P, *J S I R.* 1953, 3B, 119

HYDNOCARPUS Gaertn

Report of a chemical investigation of Chaulmoogra oil in connection with leprosy treatment Sudhamoy, Ghosh, *Ind Jour Med Res*, 1917, 4, 691.

Chemotherapeutic experiments with Chaulmoogras and allied preparations O.H Schohl, *Phillipine J Sci* 23, 533, 1923

Chaulmoogra oil in the treatment of Trachoma. Gubbay, B B & Row, T M., *Ind Med Gaz.* 1929, 64, 563.

A preliminary note on the action and uses of 'Alepol'. Dikshit, B B & Row, T.M, *Ind. Med. Gaz* 1931, 66, 317

Alepole in Leprosy Dikshit, B B., *Ind Med Gaz*, 1932, 67, 7.

Alepole in the treatment of Leprosy Bhandari, A. D., *Ind Med Gaz*, 1932, 67, 244.

Chemical composition of Congo Chaulmoogra seeds and fat L. Adraims, *Chem. Abst.* 6754. 1946.

Examination of the purity of Chaulmoogras aethylicus and oleum chaulmoogra. J.T. Kruithof; *Chem. Abst.* 2859, 1947.

H. ANTHELMINTHICA Pierre

Hydnocarpus anthelmintica seeds from Ceylon Anon; *Chem. Abst.* 3391, 1930.

Analysis of Chaulmoogra oils Hydnocarpus anthelmintica oil and Taraktogenos kurzii (Chaulmoogra oil). Howard Cole and Humberts to T. Cardoso, *J Am Chem Soc* 61, 3442, 1939

H. WIGHTIANA Blume

Efficacy of ethyl chaulmograta ethyl hvdnocarpate and ethylesters of the total fatty acids of Hydnocarpus wightiana oil Bonifacio De Vera and Casimiro B Lara, *Chem Abst* 1424, 1930.

The constituents of Hydnocarpus wightiana oil Howard Cole. *Chem. Abst.* 1466, 1930

Reduction of irritation by iodised ethyl esters of Hydnocarpus wightiana oil, Howard. Cole, *Chem. Abst* 1466, 1930.

Chemistry of antileprosy oils (Hydnocarpus wightiana) Buu Hoo Paul Cagniant and Joseph Janicaud, *Compt. Rend* 212, 577.

Hydnocarpus wightiana A note on the keeping properties of oil from, and its derivatives Basu, U P & Majumdar, A., *J Ind Chem Soc* 17(1940), 280

HYDROCOTYLE ASIATICA Linn

Chemical examination of the constituents of Hydrocotyle asiatica. M A. Wali and M C Tuminkatti, *Proc Ind Acad Sci*, 5A, 109, 1937

A preliminary note on the fission of Vascular Cylinder in some of the roots of Hydrocotyle asiatica Linn. Chakraverti, D N, *Curr Sci.*, 1940, 9, 230

Identification of Brahmi Nityendra Nath Sircar, *Sci & Cultr.* 1941, 7, 120.

Investigation of Indian medicinal plants, Hydrocotyle asiatica, Vitex negundo and Monniera cunifolia. Basu, N K., Lamsal, P P and Singh, G B., *Quart. J Pharmacol* 20, 135, 1947

HYGROPHILA SPINOSA T Anders

Hygrophila spinosa. Chemical examination of the roots of Ghatak, Narendra Nath & Dutt, Sikhishushan *J Ind Chem Soc* 8 (1931), 23

Some common indigenous remedies, Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum Chopra, R. N. and Ghosh, S, *Ind Med Record* 55, 77, 1935.

An investigation of oil from seed of Hygrophila spinosa Godbole, N. N. Gunde, B D. & Srivastava, P. D, *Oil and Soap*, 18, 206, 1941.

Chemical investigation of Hygrophilla spinosa. Lagawankar, J. D, Phalankar, N.L. and Bhide, B V, *J Univ Bcmabay.* 13, 15, 1945.

HYOSCYAMUS Linn.

The value of Palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves Bernard S Feinstein and Frank J. Slama, *J. Am Pharm Assoc* 29, 370, 1940.

A belladonna and Hyoscyamus extract which is stable in the air and contains the alkaloids of the drug in mutually unchanged relative proportion, J Gjesing Andersen, *Chem Abst* 6497, 1944.

Increased alkaloidal contents of induced polyploids of Datura Atropa and Hyoscyamus Part I Datura species Jack Morris Rowson, *Quart J Pharm Pharmacol* 18, 175, 1945

HYOSCYAMUS NIGER Linn

Indian Henbane Handa, K L., Kapoor, L D & Chopra, I C, *Curr. Sci*, 1947, 16, 315

Cultivation of Hyoscyamus niger Linn Chopra, I C, Kapoor L.D, Handa, K L. *J S I R* 1953, 5A, 238

HYPERICUM PERFORATUM Linn

Hypernin, a glucoside of Hypericum perforatum Zofia Jerzmanowska, *Chem Abst* 7299, 1939

HYPTIS SUAVEOLENS Poit

Essential oil from Hyptis suaveolens Nayak, K G & Guha, P C, *J. Ind Chem Soc* 29 (1952), 183

ILEX VOMITORIA (Soland) Hort

The pharmacological action of Cassine Croceum De and Mundulea suberosa Benth N Sapeika, *S African J Med Sci* 10, 51, 1945

ILICIUM PARVIFLORUM Michx.

Note on the volatile oil of Illicium parviflorum Michx Foote, P A., *J. Am Pharm. Assoc* 27, 573, 1938

I. RELIGIOSUM Sieb & Zucc

A study of Illicium religiosum Chen, K K, *J Am Pharm. Assoc* 15, 861, 1926.

INDIGOFERA ENNEAPHYLLA Linn.

Chemical investigation of Indigofera enneaphylla Linn. and the isolation of its active principle Satyendra Nath Chatterji and Shikhibhushan Dutt, *Proc Natl Inst India* 3, 374, 1937

I. LINIFOLIA Retz

Chemical examination of Indigofera linifolia Retz The isolation of its active principles. Mahado Prasad Gupta and Shikhibhushan Dutt, *Proc Natl Acad Sci. India*, 8, 49, 1938

INULA HELENIUM Hk. f & T.

Examination of plants for insecticidal principles, Elecampane. F. Tutin, *Chem. Abst.* 4337, 1933

Bitter principles of Elecampane roots Karl, Fr., W Hansen; *J. Prakt. Chem.* 136, 176, 1933.

Pharmacological studies of Korean Inula helenium L. Tinsyo Go, *Japan J Med Sci Pharmacol.* 12, 110, 1938

I. ROYLEANA D.C

Inula royleana D C. Its Chemistry, and pharmacological action Chopra, I C., Kohli J D & Handa, K L, *Ind Jour Med Res*, 1945, 33, 139

IPOMOEAE Linn.

The seed structure of Ipomoea, a criticism. Maheshwari, P., *Sci. & Cultr*, 1944, 9, 557.

I. HEDERACEAE Jacq

Studies in the specification of Indian Medicinal Plants Pt. III Kaladana Sengupta, S B, Gupta, H N, *Ind Jour Pharm*, 1948, 10, 106.

I. MURICATA Jacq.

Ipomoea muricata. Fatty oil from the seeds of Kelkar, Phalnikar & Bhide, *J Ind Chem Soc* 24 (1947), 87-90

Ipomoea muricata Chemical examination of the seeds of Misra & Tewari *J Ind Chem Soc* 28 (1951), 221

Chemical examination of seeds of Ipomoea muricata Part II Misra, A L & Tewari, J. D, *J Ind Chem Soc* 29 (1952), 63

Chemical examination of seeds of Ipomoea muricata Part III. Misra, A L. & Tewari, J D, *J Ind Chem Soc* 29 (1952), 430

Chemical examination of Ipomoea muricata seeds, Part IV. Misra A L., Tewari J D, *J I C S* 1953, 6, 391

I PESCAPRAE Roth

A phytochemical study of Ipomoea pes-caprae Gustav E. Cwalina and Glenn L. Jenkins, *J. Am Pharm Assoc* 27, 585, 1938.

Study of the leaves of Ipomoea pescoprae B V Chrestense and J A. Reese, *J Am Pharm. Assoc.* 27, 195 1938.

I. PULCHELIA Roth.

Damping of Ipomoea pulchelia Roth, due to Pythium aphanidermatum (Eds) Fitz Mohmud. K. A & Jain, A C.

I. REPTANS Poir.

The carotenoids and some lipoids of Ipomoea reptans Poir. M. Ishi, *Chem. Abst* 3376, 1935

JASMINUM Linn.

Jasmine Oil—J N. Rakshit, *Perfum Record*, 26, 1937, 241.

Jasmin Flowers—Sadgopal; *S.P.C* July, 1939, p. 589.

Development of essential oil industry (Ocimum lemon, O. basilicum, O. canum, O. sanctum, lemon grass, Motia, Palmarosa grass). Jitandra Nath Rikshit, *Sci & Cultr* 1939, 5, 108.

Chromosome numbers in some ornamental Jasmines. Mridula Dutt, *Sci & Cult.*, 1952, 17, 527.

JATEORRHIZA PALMATA Miers

Bitter principles of Columba root, Palmarin F Wesseky, K Schonol and W Iseman, *Monatsh.* 68, 21, 1936.

Bitter principles of Columba root, Methylation of Columbin. F. Wessely and K Jentzsch, *Monatsh* 70, 30, 1937

Evaluation of Colombo root and its preparations H Neugebourer and K Brunner, *Arch. Pharm* 276, 199, 1938.

JATROPHA CURCAS Linn.

Seeds of Jatropha curcas from the Belgian Congo Adriaens, *Chem Abst.* 5440, 1936.

Chemical investigation of the seeds of Jatropha curcas Linn Desai, C M & Vyas, M T, *Curr Sci*, 1949, 18, 49

J GLANDULIFERA Roxb

Jatropha glandulifera Roxb Examination of the seeds of. Alimchandani, Badami Katti, *J Ind Chem Soc* 26 (1949), 523.

JUGLANS REGIA Linn

The Vit C contents of Tender Walnut (Juglans regia) Ratnanathan, S., *Ind Jour Med Res*, 1942, 30, 513

JUNIPERUS Linn.

Oil of Juniper berries Ernst S Guenther, *Am Perfumer* 33, 64, 1936

Analysis of Juniper berries. P Casparis and W Freund, *Pharm. Acta Helv* 13, 307, 1938

New hydrocarbon from juniper oil P Casparis and W Freund, *Pharm Acta Helv* 14, 1, 1939.

Assay of Juniper for volatile oil content *Chem Abst* 4523, 1940

J COMMUNIS Linn

Constituents of some Indian essential oils, essential oil from the seeds of Zanthoxylum ovalsifolium & Juniperus communis J. L. Simonsen, *Ind Forest Records* 11, 1, 1924

The constituents of some Indian Essential Oils Part XV—The Essential Oil from the seeds of Juniperus communis. Simonsen J L, *Indian Forest Records*, 1925, Vol. XI Part I, pp 6-9

KAEMPFERIA GALANGA Linn.

Constituents of some Indian essential oils Essential oil from the rhizomes of Kaempferia galanga. Puthan Madhathil, Bhaskara Panicker, B. Sanjiv Rao and J. L. Simonsen, *J. Ind Inst of Sci* 9A, 133, 1926.

LACTUCA SCARIOLA Linn.

Kahu (Lactuca scariola) Seed Oil Component fatty acids of. Dhingra & Pershad, *J Ind Chem. Soc* 22 (1945), 127

LAGENARIA VULGARIS Ser

Wilt disease of Lagenaria vulgaris Ser Ashok Kumar Kar, *Sci & Cult*, 1946, 12, 290

An abnormal flower of Lagenaria vulgaris Ser Ahmed Shamsul Islam, *Sci & Cult*, 1951, 16, 427

LANSIUM ANAMALAYANUM Bedd

Chemical Investigation of some minor Forest Products The Essential oil of Chigatmani (Lansium anamalayanum) Rao K A Narain, Varadhan C and Janniah S L., *Department of Industries, Bangalore*

Studies in sesquiterpenes Part X Sesquiterpenes of the essential oil of Lansium annamalayanum Bedd A Somasekar Rao, Dutta, K B, Subdev & Guha, P C, *J Ind Chem Soc* 29(1951), 604

L. DOMESTICUM Correa

Chemical study of the seeds of Lansium domesticum Correa Kassem Pangsrivongse, *Chemical Abst* 3088, 1938

LANTANA CAMARA Linn

Oil from the flowers and leaves of Lantana camara—D. D Kanga, *Journal of the Indian Institute of Science, Bangalore* Vol. I, Pt IX, pp 93 to 95, 1914

Travancore Essential Oils Lantana camara Moudgill, K. L. and Vrindachalam, P N, *Department of Industries, Trivandrum*, 1923, Bulletin No XVII, pp 7-10.

Travancore essential oils from the leaves of Lantana camara. Moudgil, K L, *Perfumery Essent Oil Record* 16, 9, 1925.

Volatile oil of Lantana camara L. Kinzo Kafuku, Tassaku Ikeda and Yasuji Feijita, *J Chem Soc Japan*, 55, 305, 1924.

Lantana camara (The paper was read in the Annual meeting of the National Academy of Science and has been published in the annual issue of the Academy, 1939)

LATHYRUS SATIVUS Linn

The production and pharmacological action of Khesari Amine. Acton, Hugh W. & Chopra R.N., *Ind. Med Gaz*, 1922, 57, 412.

The toxic effects in Guinea-pigs of diets containing large proportion of Lathyrus sativus Bhagwat, K., *Ind. Jour Med Res.*, 1946, 34, 299

The toxic effects of Ticora (Khesari) on man Jacoby, H., *Ind. Med. Gaz.*, 1947 82, 122.

Study on "Lathyrism" a disease produced by Lathyrus sativus Linn. De, H N. & Datta, P.K., *Sci & Cult*, 1948, 14, 159.

LAURUS CAMPHORA Linn

Laurus camphora Nees (Cinnamomum camphora) V E Vorotzov,
Chem Abst 5474, 1933

LAVENDULA BURMANNI Benth

Oil from the flowers and leaves of Lavendula burmanni—D D Kanga,
Journal of the Indian Institute of Science, Bangalore Vol I, Pt VIII,
pp 181 to 189, 1916

LAWSONIA INERMIS Roxb

Essential oil from the flowers of Camphine or Henna plant Antia
M B & Kaushal, R , *Curr Sci* , 1950, 19, 284

LEUCAS ASPERA Spreng.

Studies on Leucas aspera Shirazi, A M J , *Ind Jour Pharm* , 1947,
9, 116.

LEGUMINOSAE

Seed oil of Formosan Plants Constituents of seed oils Leguminosae.
Anona squamosa Kinzo Kafulice and Chintu Hata, *J Chem Soc
Japan* 55, 369, 1934

LENS ESCULENTA Moench

Studies in nutrition value of Indian vegetable foodstuff Pt III
Nutritive values of Lentil--Lens esculenta Moench , Cow pea—
Vigna catjang Walp., and Aconite Bean—Phaseolus aconitifolius
Jacq Niyogi, S P , Narayana, N. & Desai, B G , *Ind Jour Med
Res* , 1932, 19, 859.

LEPIDIUM LATIFOLIUM Linn

The content of Lepidium latifolium and Cochlearia armoracia in
essential oils S Rivos Goday and M Gomez Serranillos, *Chem Abst*
4432, 1945

LEUCAENA GIAUCA Benth

Leucenol, a definite principle obtained from the seeds of Leucaena
giauca Benth Marcel Maseve, *Compt-Rend* 240, 890. 1937

Presence of Stachyose (Manotetrose) in the seed of Leucaena glauca
Benth H Heressey and M Mascré, *J Pharm Chem* 1, 521, 1941

Structure of leucenine (Leucenol) from Leucaena glauca Benth
A F Bichel, *J Am. Chem Soc* 60, 1801, 1947

LINUM USITATISSIMUM Linn.

The hydrolysis of ptyic compounds derived from seeds of hemp,
horse bean, horse chestnut flax, wheat and embryos of rye W.
Jarosza, *Chem Abst.* 5501, 1934.

LITSAEA Lamk.

Actinodaphne and Litsaea fats as raw material for a valuable new
detergent S. V. Puntambekar, *Ind Forester*, 60, 707, 1934.

L. ZEYLANICA Nees.

Oil of *Litsaea zeylanica*, Nees. Rao, B. S.; *Journal of Indian Institute of Science, Bangalore*, **15A**, 1932, 71.

Bellary leaf oil Sharma, M. R. G., Varadhan C., Sastry, S. G. *J. S. I. R.*, 1953, **6B**, 243.

LOBELIA Linn.

Amino nueteric determination of alkaloids in belladonna, lobel herb and harmel roots, G. Ya Kait, *Chem. Abst.* 2996, 1942.

The origin of Haustoria in the value of *Lobelia*. Maheshwari, P. *Curr. Sci.*, 1944, 13, 186

L. INFLATA Linn.

Indian substitute for the drug *Lobelia inflata* *Ind. Jour. Pharm.* 1944, 6, 105.

L. NICOTIANAEOFOLIA Heyne.

Pharmacognostic studies on *Lobelia nicotianaefolia* Heyne. Datta S. C. & Bal, S. N., *Sci. & Cult*, 1944, 10, 260

Lobelia nicotianaefolia Heyne as substitute for *Lobelia inflata* Linn. Mukerji, B. & Ghosh, S. K., *Curr Sci.*, 1945, 14, 198.

L. PYRAMIDALIS Vahl.

Pharmacognostic investigations on *Lobelia pyramidalis* Vahl. substitute for *Lobelia inflata* Linn. Amiya Datta & Datta, S. C. *J. Sci. Indl. Res.*, 1951, **10B**, 218.

LOLIUM TEMULENTUM Linn.

Pisonous foodgrain wheat mixed with *Lolium temulentum*. Greva S. D. S. & Bhandari, P. N., *Ind. Med Gaz*, 1946, 81, 294.

LORANTHUS LONGIFLORUS Den.

Parasitism of *Cuscuta reflexa* Roxb. & *Loranthus longiflorus* Des Sheriar, K. C., *Sci. & Cult.*, 1951, 17, 218.

LUFFA ACUTANGULA Roxb

Luffa acutangula The Chemical and Pharmacological investigation of *Luffa* seeds Grewal, K. A. & Kochhar, B. D., *Ind Jour. Med. Res.* 1943, 31, 63

L. AEGYPTIACA Mill.

A study of the oils from the seeds of *Luffa aegyptiaca*, *Benincas cerifera* and *Allium cepa*. Phadnis, K. D., Rege, A. V., Pishawika D. G., and Shah, S. V., *J Univ. Bombay*, **17A**, 62, 1948.

L. AMARA Roxb.

Chemical constituents of *Luffa amara* Roxb Pt. I. Isolation of a crystalline bitter from the seeds Chaudhry, G. R., Vishwa Nath Sharma & Siddiqui, S.; *J. Sci. Indl. Res.*, 1951, **10B**, 26.

L. CYLINDRICA Linn.

The crystalline principle of *Luffa cylindrica* Linn. Roemer, Martin, De, La Torre and A. C. Santos, *Chem Abst* 5593, 1939.

Saponins of *Luffa cylindrica* Linn. Roemer., Amparo Mendoza, Parahuman Cruz and Alfredo C. Santos, *Chem Abst.* 4915, 1941.

A chemical study of *Luffa cylindrica* with special emphasis on its saponins. Amparo. S. Mendoza and Alfredo C. Santos, *Chem Abst* 7652, 1941.

L. ECHINATA Roxb. & L. GRAVEOLENS Roxb.

Chemical examination of the seeds of *Luffa graveolens* Roxb. & *Luffa echinata* Roxb. Nigam, Ram Gulam Singh, Pandya, K. C & Tayal, J N, *Curr. Sci.*, 1949, 18, 451

LUPINUS (Tourn) Linn

Biological experimental study of the Lupine alkaloids A. Guillaume, *J. Pharm. Chem* 12, 335, 1930.

L. ALBUS Linn.

Phyto-Pharmacological experiments with *Lupinus albus*. Ludwig Johanning, *Chem. Abst* 167, 89, 1930

Hypoglucmic action of the seeds of *Lupinus albus* Orestin, G., *Chem Abst* 1511, 1941

The alkaloids of *Lupinus albus* L and its supposed antimalarial action Liberalli, C H, *Chem Abst* 4101, 1944

LUVUNGA SCANDENS Ham

On the constitution of natural coumarins isolated from *Luvunga scandens* Ham. Bose, P K & Miss Mukerji, A, *J Ind Chem. Soc.* 21 (1944), 181.

LYCOPodium.

Toxicity of the alkaloids of *Lycopodium* P. Oficialski; *Bull Sci. Pharmacol* 33, 470, 1937

Pharmacological action of certain *Lycopodium* alkaloids Henry M. Lee and Chen, K K, *J. Am Pharm. Assoc* 34, 197, 1945

L. CLAVATUM Linn.

Alkaloids of *Lycopodium* (*Lycopodium clavatum* L.). Achmatowie and W Uzieblo, *Chem Abst* 9092, 1938.

The alkaloids of *Lycopodium* species, *Lycopodium clavatum* L. Leo. Marion and Richard H. F. Manske, *Can. J. Res.* 22B, 137, 1944

L. PHLEGMARIA Linn.

Lycopodium phlegmaria Linn., on occurrence of two sporangiate sporophyllus in. Ghosh, A. K., *Sci & Cult.*, 1941, 7, 410.

MABA NIGRESCENS Dalz.

Occurrence of marginal strands in the leaf of *Maba nigrescens* Dalz
Rao, T. A. & Kelkar, S. S , *Sci & Cult* , 1950, 16, 209

MAHONIA Nutt.

Structure of Neprotine A new alkaloid of *Mahonia* species Chatterjee, R & Guha, M P., *Sci. & Cult* , 1950, 16, 119

M. ACANTHIFOLIA Don.

The alkaloids of *Mahonia acanthifolia* Don. Chatterjee, R , Guha, M P , *Sci & Cult* , 1949, 15, 163.

MALLOTUS PHILIPPINENSIS Muell

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency XIII Kamala Cains, J F & Mhaskar, K S , *Ind Jour. Med Res* , 1921, 9, 202

The constitution of Indian Kamala (*Mallotus philippinensis*) Sikhishushan Dutt, *J Chem Soc* 2044, 1925.

The Constitution of Rottlerin from Indian Kamala Part II Dutt, Sikhishushan, Puri, & Dhanraj, *J Ind Chem Soc* 5 (1928), 21.

Rottlerin V Bakshi, H S Ravi, S Jalota, Narang, K S. & Ray, J N., *Curr. Sci.*, 1939, 8, 165.

The constitution of Rottlerin Ray, J. N , Narang, K S. & Roy, B S., *Curr Sci* , 1939, 8, 558

The use of Kamala as an antioxidant of Ghee Govindarajan, S V. & Banerjee, B. N , *Curr Sci* , 1939, 8, 559.

The fatty oil from the seeds of *Mallotus philippinensis* Muell Arg. (N O Euphorbiaceae) Singh, Bawa Kartar, Saran Brij Mohan, *Curr Sci.*, 1942, 11, 360

The fatty oil from the seeds of *Mallotus philippinensis* Muell Puntambekar, *Curr Sci* , 1942, 11, 464

Kamala as a colouring agent for hydrogenated vegetable oil Its toxicity on white Rats. Gupta, J C & Chatterjee, M L ; *Sci & Cultr* , 1946, 11, 375

Quality of commercial Kamala and its treatment Subba Rao, V , *Ind Jour Pharm.*, 1947, 9, 81

The treatment of intestinal worms with the Indigenous drugs— Butea, Embelia and Kamala Mukerji, A. K , Bhaduri, N V., *Ind Med Gaz* , 1947, 82, 66.

Chemical Examination of the seeds of *Mallotus philippinensis* Muell Arg. Pt I. Analysis of drying oil and seed cake Aggarwal, J S , Bhatnagar, S. S , Parkash Narain & Karimullah, *J. Sci Indl. Res* , 1948, 7B, 136.

The Chemical Constitution of the oxy-acid from the seed oil of *Mallotus philippinensis* Muell. Arg. (Kamala) Gupta, S. C , Vishwa Nath Sharma & Aggarwal, J S , *J. Sci Indl Res* 1951, 10B, 76.

Chemical Examination of the seeds of *Mallotus philippinensis* Muell. Arg (Kamala) II Constitution of the unsaturated Hydroxy acid isolated from the oil Gupta, S. C., Sharma, V. N. & Aggarwal J. S., *J. Sci Indl Res*, 1952, **11B**, 463.

Chemical Examination of the seeds of *Mallotus philippinensis* Muell. Arg (Kamala) III—Constitution of Mamlolenic Acid isolated from the oil Gupta S. C., Gupta S. S., Aggarwal, J. S., *J S I R* 1953, **6B**, 240

MALVA ARBOREA Webb.

Oils from *Althea officinalis* and *Malva arborea* H. Ya. Tropp, *Chem. Abst* 1597, 1936

M SYLVESTRIS Linn.

Chemical and pharmacological researches on *Malva sylvestris* Luigi Callegari, *Chem Abst* 1589, 1946.

MANGIFERA Linn

A note on the genus *Mangifera* Linn. Mukerjee, S. K., *Sci & Cultr.* 1942, 8, 92

M. INDICA Linn.

Mangifera indica. Chem. Examination of mango "Chep"—the exudation of the fruit of Vasishta, Shiam Kishore & Siddiqui, Salimuzzaman, *J Ind Chem Soc* 15 (1938), 110

Chemical Examination of Mango blossoms (*Mangifera indica* L., Flora) Jogendra Lal Ball & Salimuzzaman Siddiqui, *J Sci Indl Res*, 1948, **7B**, 100

Wild mangoes of India. Mukerjee, S. K., *Sci & Cult*, 1950, 15, 464.

MANIHOT ESCULENTA Crantz.

Age of Cassava Plant for max. yield Chatterjee, K. N., *Sci & Cult*, 1949, 14, 533.

M UTILISSIMA Pohl.

The poisonous constituents of sweet Cassava (*Manihot utilisissima*) and the occurrence of hydrocyanic acid in foods prepared from Cassava B. J. W. Furnock, *J Trop Med Hyg* 43, 65, 1937

Therapeutically utilisable constituents of *Manihot utilisissima* and *Mercurialis annua* or *perennis* Friedrich W. Friese, *Chem Abst.* 2288, 1938

MANSONIA DIPIKAE C. S. P.

Mansonia dipikae C. S. Purkayastheo sp. Nov. A new species from Assam Jain, B. C., Iyer, B. H., & Guha, P. C., *Sci & Cultr*, 1947, 12, 405

MARRUBIUM VULGARE Linn.

Comparative marrubium content in Marrubium vulgare from European Vs. American seeds Adelia Mccrea; *J. Am Pharm Assoc* 19, 231, 1930

Maruburr, the bitter principle of hove bound (Marrubium vulgare) Alexander Lawson and E. D. Euktice; *J. Chem. Soc.* 587, 1939.

Substance contained in Marrubium vulgare Gualtenio Seppi *Chem Abst.* 4893, 1947

MARTYNIA DIANDRA Glox

Fatty oil from the fruit of Martynia diandra (N. O. Pedaliaceae) Shah, S. V., Airan J. W. & Rege, A. V., *Curr Sci.* 1942, 11, 291.

MATRICARIA Linn.

Biochemical studies of the glucosides of Camomile flowers. Charlev Beguin; *Pharm Acta Helv* 7, 332, 1932

Estimation of essential oil in comomile flowers. F. Gestirner, *Chem. Abst.* 5477, 1933.

Anti-inflammatory action of Camomile oil Wolfgang Henlner and Franz Grabe; *Arch. Expt Path Pharmacol* 171, 329, 1933

Biochemical studies of the glucosides of camomile flowers, comparison of samples of different origin collected in 1932. Ch Beguin, *Pharm. Acta Helv.* 10, 147, 1935

Camomile Orlando Gulminelli, *Chem Abst.* 1517, 1936

M. CHAMOMILLA Linn.

New results for the evaluation of the pure azulene content of Matricaria chamomilla and its preparations Kaiser, H. and Frez, H., *Chem Abst* 1441, 1939.

MELIA AZEDARACH Linn

Fatal intoxication by fruit from Melia azedarach Regelo, L., Cavrrabala, E.; *Chem Abst* 6951, 1939

Chemical Examination of the Bakayan fruit (Melia azedarch Linn.) Amir Chand, Chittaranjan Mitra & Salimuzzaman Siddiqui, *J Sci Indl Res* 1948, 7B, 69

MELILOTUS PARVIFLORA Desf.

Powdery Mildew on Melilotus parviflora Desf. Mahmud, K. A. & Namen, K. G., *Sci. & Cult.*, 1952, 17, 473.

MENTHA AQUATICA Linn.

Studies in the genus Mentha. The non-volatile constituents of Mentha aquatica Linn. Samuel M. Gordon; *Am. J. Pharm.* 100, 433, 1928.

Mentha aquatica L. Crestoforo Masino, *Chem. Abst.* 2741, 1937.

M. VIRIDIS Linn.

Japanese peppermint oil. Oil of Japanese *Mentha viridis* L. Var *Crispa* Benth. Carvone Tetu Nagasawa; *Chem. Abst.* 219, 1940

MERCURIALIS AMNUA Linn.

Therapeutically utilisable constituents of *Manihot utilisima* and *Mercurialis amnua* or *perennis* Friedrich. W. Friese; *Chem. Abst.* 2288, 1938.

MESUA FERREA Linn.

Mesua ferrea Preliminary note on Mesuol. The bitter principle of. Dhanindra Bhushan Dutt, Deb, Narendra Chandra & Bose, Prafulla Kumar, *J. Ind. Chem. Soc.* 17 (1940), 277

Refining of Nageshwar oil Gupta, A. C., *J. Sci. Indl. Res.*, 1951, **10B**, 24.

MILLETTIA PACHYCARPA Benth.

Occurrence of Rotenone in *Millettia pachycarpa* Ghose, T.P. & Krishna, S, *Curr. Sci.*, 1937-38, 6, 57.

MIMOSA PUDICA Linn.

The chemical properties of the hormone from *Mimosa pudica* Hans. Fitting, *Chem. Abst.* 8300, 1936.

Chemical Examination of the seeds of *Mimosa pudica* Linn. Pt I. Joti Sarup Aggarwal & Karimullah, *J. Sci. Indl. Res.*, 1945-46, 4, 80.

MIRABILIS JALAPA Linn.

A note on starch in *Mirabilis jalapa* soluble in cold water Chaudhri, T. C., *Sci. & Cultr.*, 1947, 12, 449

MOMORDICA CHARANTIA Linn.

Balsam pear seed oil (*Momordica charantia* L.) Yoshiyuki Toyama and Tomotaro Tsu Chiya, *J. Soc. Chem. Ind. Japan.* 39, 220, 1936.

Preliminary study of an alkaloid like material obtained from *Cundeamor* or *Momordica charantia*. Luis Torres Diaz, *Chem. Abst.* 6134, 1936.

Chemical Examination of Karla plants Airan, J.W & Ghatge, N D, *Curr. Sci.* 1950, 19, 19.

M. DIOICA Roxb

The fatty oils from the seeds of *Momordica dioica* (N. O. Cucurbitaceae) Airan, J W & Shah, S.V., *Curr. Sci.*, 1942, 11, 246

MONNIERA CUNEIFOLIA Michx.

Investigation of Indian medicinal plants *Hydrocotyle asiatica*, *Vitex negundo* and *Monniera cuneifolia*. Basu, N K, Lamsal, P P., and Singh, G.B., *Quart. J. Phar. Pharmacol.* 20, 135, 1947.

MORINDA CITRIFOLIA Linn

Notes on the constituents of *Morinda citrifolia*. John Lionel Simonsen *J Chem. Soc* 561, 1920

MORINGA CONCANENSIS Nimm

Chemical investigation of seed oil of *Moringa concanensis* Patel, C B, *Curr Sci*, 1943, 12, 272.

M OLEIFERA Lamk

Availability of calcium in Lady's finger (*Hibiscus esculentus*), Cabbage (*Brassica oleracea* var. capitata), Drumstick (*Moringa oleifera*) and Amaranth Tender (*Amaranthus gangeticus*). Pt I. Availability of Calcium in vegetables determined by experiments on growing rats. Basu, K P & Ghosh, D, *Ind. Jour Med. Res*, 1943, 31, 29.

Availability of calcium in Lady's finger (*Hibiscus esculentus*), cabbage (*Brassica oleracea* var capitata), Drumstick (*Moringa oleifera*) & Amaranth Tender (*Amaranthus gangeticus*) Pt II Availability of calcium in vegetables determined by metabolism experiments on a human adult Basu, K P & Ghosh, D, *Ind Jour Med Res*, 1943, 31, 37

M PTERYGOSPERMA Gaertn

A preliminary note on the action of the alkaloids of *Moringa pterygosperma* (N O Moringae) Chopra, R N, *Ind Med Gaz*, 1932, 67, 128

Moringa pterygosperma (N O Moringae). Chopra, R N., Premankur D C & De, Nripendra Nath, *Ind Jour Med Res*, 1933, 20, 533

A Chemical examination of the bark of *Moringa pterygosperma*. Ghosh, S, Chopra, R N & Datta, A *Ind Jour Med Res.*, 1935, 22, 785

Antibiotic principles from *Moringa pterygosperma* Kurup, P A & Rao, P L., *Curr Sci*, 1950, 19, 54

A note of physiological and chemical findings of the active principle (Spirochin) of the *Moringa pterygosperma* Chatterjee, G. S, Mitra, S R., *Sci & Cult*, 1951, 17, 43

Chemical investigation of the fatty oil from the seeds of *Moringa pterygosperma* Subba Rao, B C, Rao, K N., Jois, H S, *J.I.C S*, 1953, 7, 477.

MORINGACEAE

Notes on the affinities of Moringaceae. Chatterjee, D., *Sci. & Cult*, 1948, 14, 253

On the affinities of Moringaceae. Datta, R.M, Mitra, J N. & Chatterjee, D., *Sci. & Cult.*, 1949, 15, 115.

MORUS INDICA Linn

On the Hermaphrodite flowers of *Morus indica* Linn Datta, R M ,
Sci & Cult , 1938, 4, 301.

On the Hermaphrodite flowers of *Morus indica* Linn Datta, R M ,
Sci. & Cult , 1939, 5, 373

Fasciation in the Female inflorescence of *Morus indica* Linn Datta,
R M , *Sci & Cultr.*, 1941, 6, 425

M. NIGRA Linn.

The blood sugar lowering action of the leaves of *Morus nigra* L Henri
Leelvé, *Chem Abst.* 1507, 1935

MUCUNA PRURIENS D C

Indian Medicinal plants Pt V *Mucuna pruriens* (N O Papilionaceae),
Pt I Mehta, J C & Majumdar, D N , *Ind Jour Pharm* , 1944, 6, 92

MUNDULEA SUBEROSA Benth.

The pharmacological action of Cassine Croceum De and *Mundulea*
suberosa Benth N Sapeika, *S African J Med Sci* 10, 51, 1945

MUNDULEA SUBEROSA Benth

Chemical constituents of *Mundulea suberosa* Th M Meyer, *Chem*
Abst 6607, 1947

MURRAYA EXOTICA Linn & M KOENIGI Spreng

Essential oils from the leaves of *Murraya koenigii* (Spreng), *Murraya*
exotica Linn and *Murraya exotica* Var *Ovalifoliata* (Ergler) A R
Penfold and J. L Simonsen, *J Proc Royal Soc* 59, 146, 1925

Murraya exotica Natural glucosides Part I The constitution of the
Glucoside present in *Murraya exotica* Praphela Kumar Bose & Miss
Asima Mukerjee, *J Ind Chem Soc* 14 (1937), 489

MYRISTICA FRAGRANS Hoult

Essential oil from the leaves Nutmeg (*Myristica fragrans*) M Meyer,
Chem Abst 4549, 1941

Chemistry of Nutmeg—Ari (Jay-Patri) Ashok D Pishawikar,
Pishawikar, D. G., *Curr Sci* 1953, 3, 81.

MYRSINE AFRICANA Linn.

Myrsine africana Linn. Active principles of *M. africana* Krishna,
S. & Varma, B. S ; *J. Ind Chem Soc* 13 (1936), 115

The active principle of Kurjan seed (*Myrsine africana*) Annon;
Chem. Abst. 36, 319, 1938

NARCISSUS TAZETTA LINN.

A new alkaloid in the bulbs of *Narcissus tazetta* L Yosiziro Kihara;
J. Agr. Chem. Soc. Japan, 15, 128, 1939

NARDOSTACHYS JATAMANSI D C.

Chemical constituents of Nardostachys jatamansi. Pt. I. Isolation of a crystalline acid and an essential oil Chowdhry G R, Vishwa Nath Sharma & Siddiqui, S., *J Sci Indl Res.*, 1951, **10B**, 48

NERIUM ODORUM Soland.

Chemical examination of the bark of Nerium odorum (Soland.) Gajanan P Pendse and Sikhibhushan Dutt, *Bull Acad Sci. United Provinces.* 3, 209, 1934

Chemical Examination of the root of "Nerium odorum". Part I Garde, S.T., *Journal of the Indian Institute of Science, Bangalore*, Vol I, Part XVIII, pp 181 to 189, 1910

Pharmacological action of odorin, a glucoside of Nerium odorum. Isao Numoto, *Chem Abst.* 4734, 1940

N. OLEANDER Linn

A pharmacological study of folinerin, a new glucoside from Oleander leaves. Italo Simon, *Chem Abst* 4799, 1943.

Medicinal use of the glucoside of Nerium oleander L. Gibert Queralto, *Chem. Abst.* 5786, 1943.

Pharmacology of the new Cardiac glucoside folinerin (From Nerium oleander) Effect on the cardiovascular system. V I Simon, *Chem. Abst* 5299, 1944

Glucosides of Nerium oleander of the Argentine, Oscar A Rossi and Jaun A. Izquierdo; *Chem Abst* 7940, 1948

NICOTIANA Linn

The occurrence of crossing over in Nicotiana species Hybrids Kostoff Dontcho, *Curr Sci*, 1933-34, 2, 370

Tobacco seed oil Fatty acid comp of. Rao, Rao & Venkateswarulu, *J. Ind Chem. Soc.* 20 (1943), 374

Glyceride comp of Tobacco seed oil Venkata Rao, C Narsingarao, M & Venkateswarlu, A, *J. Ind Chem. Soc* 21 (1944), 249

Manufacture of 40% Nicotine as a soln. of nicotine sulphate in water Subramanian, T. S & Varia, M. C, *J Sci Indl Res.*, 1944 45, 3, 143

N. RUSTICA L. & N GLAUCA Grab

Nicotine and citric acid content in the progeny of the Allopolyploid Hybrid Nicotiana rustica L X. N glauca Grab Kustoff, Dontcho; *Curr. Sci*, 1939, 8, 59

N. TABACUM Linn

Australian Tobacco Industry, article on Charles Lynch, *J. Sci Indl. Res.*, 1945-46, 4, 760.

Products of Tobacco industry Tobacco seeds and its utilization. Mahant, S. D. & Pandit P. N., *J. Sci. Indl. Res.* 1948, 229, **7A**.

Cytogenetics of *Nicotiana tabacum* Var. Viru resistant to the common Tobacco Mosaic virus Kostoff, Dontcho, *Curr. Sci.* 1948, 17, 315.

Tobacco Research in India. (Article); *J. Sci. Indl. Res.*, 1950, 9A, 69.

Utilization of Tobacco seed and Safflower seed oil in varnish and paints. Pt I, Sharma, P G, Budhiraja, N C, & Aggarwal, J. S, *J. Sci. Indl. Res.*, 1951, 10B, 33.

NUPHAR LUTEUM Sibth & Sm.

The glucoside of Nuphar luteum Marjan Hulajewski and Jerzy, Modrakowski, *Chem. Abst* 5106, 1937

NYMPHAEA (Tourn) Linn

The pharmacological action of nymphacin Zderek Kocher; *Chem. Abst* 9282, 1938

N ALBA Linn

Alkaloids from flowers of *Nymphaea alba* Jerzy, Modrakowski; *Chem. Abst* 9279, 1938

N. LOTUS Linn.

Axial floral Proliferation and Metamorphosis of stamens in a Flower of *Nymphaea lotus* Linn Bose, P K, *Sci & Cultr*, 1940, 5, 636.

OCIMUM Linn.

Development of essential oil industry (O lemon, *Ocimum basilicum*, *Ocimum canum*, *O sanctum*, Lemon grass, Motia, Palmarosa grass) Jitandra Nath Rikshit, *Sci & Cultr*, 1939, 5, 108.

O BASILICUM Linn

The volatile oil of ocimum of North India—Rakshit J N., *The Perfumery and Essential Oil Record*, March, 1938

Ocimum basilicum Pt I & II A chemical study of the oil and antibacterial properties Khorana, M L, Miss M. B Vangikar, *Ind Jour Pharm* 1950, 12, 132 & 134

O CANUM Sims

Oil of *Ocimum canum*—B. S. Rao and Co-workers; *Perfum. Record*, 28, 1937, 412.

Ocimum canum oil of North India—Rakshit, J N, *The Perfumery and Essential Oil Record*, October, 1938

Chemical examination of *Ocimum canum* Sims. Mehta, R. C, Mehta T. P., *Curr Sci*, 1943, 12, 300

O. GRATISSIMUM Roxb.

Essential oil from *Ocimum gratissimum* Roxb Nayab, U. G. & Guha D. C.; *J. Ind. Chem. Soc.* 29 (1952), 203.

O KILIMANDSCHARICUM Guerke

Camphor production from *Ocimum kilimandscharicum* Guerke
Ribeiro, D J., *J. Sci Indl Res* 1950, **9B**, 230.

O SANCTUM Linn

Ocimum sanctum (*Ind Acad. Sc. Voc. IX*, Jan 1939, pp 72-77
Sikhi Bhusan Dutt,

A seedling blight of *Ocimum sanctum* Linn. caused by *Rhizodonia solani* Kuhn Mohmud K A, *Sci & Cult*, 1950, 16, 161.

ODINA WODIER Roxb.

Gum Jeol (*Odina woder* Roxb) *J Ind Chem Soc* 25 (1948).

- | | |
|--|-----|
| (i) Solution & purification of Mukerjee & Bannerjee, | 59 |
| (ii) Hydrolysis of & sugar in Mukerjee & Banerjee, | 63 |
| (iii) Electrodialysis of. Mukerjee, | 323 |
| (iv) Dependence of Viscous electro-chemical prop of aqueous
solution of, on concentration. Mukerjee & Rohatgi, | 339 |
| (v) Influence of neutral salts on some electrochemical &
Viscous properties of the solution of Mukerjee &
Rohatgi, | 531 |

Aldobionic acid from Gum jeol & its hydrolysis. Mukerjee & Dutta,
J Ind Chem Soc 25 (1948), 113

OLDENLANDIA BIFLORA Hk. f

Chemical examination of *Oldenlandia biflora* R N S Clauba &
Tewari, J D, *J Ind Chem Soc* 29 (1952) 386.

OLEA Linn

On the occurrence of Sclereide in the Genus *Olea* Linn Rao, T A,
Sci & Cultr, 1948, 13, 299

ONONIS SPINOSA Linn

New constituent from the root *Ononis spinosa* L F Neuwald, *Arch Pharm* 277, 130, 1939

ORIGANUM VULGARE Linn

Volatile oil of *Origanum vulgare* B N Rytovskii, K A Guseva and
L Koroleva, *Chem. Abst* 5476, 1933

Essential oils of *Origanum vulgare* Linn Var *formosanum* Yasuji
Fugita, *J. Chem Soc Japan*, 57, 574

Essential oil of the flowering tops of *Origanum vulgare* infected with
Eriophyes thomasi R Salgues, *Chem Abst* 4270, 1936

OROXYLUM INDICUM Vent.

Natural flavones Part II On the colouring matter of the bark of
Oroxylum indicum Vent. Prafulla Kumar Bose & Sachindra Nath
Bhattacharya, *J Ind Chem. Soc* 15 (1938) 311.

Baicalin from the seeds of *Oroxylum indicum* Vent Mehta, C. R. & Mehta, T P, *Curr Sci*, 1943, 12, 274.

Constitution of Oroxylin Sastri, V.D N. & Seshadri, T R., *Curr Sci*, 1946, 15, 235

Tetuin, A Glucoside from the seeds of *Oroxylum indicum* Vent Mehta C R, Mehta T P, *Curr Sci* 1953, 4, 114

ORTHOSIPHON STAMINEUS Benth.

Preliminary study of *Orthosiphon stamineus* Benth. Teng Hang Tang and Chao Hsu, *J Chinese Chem Soc* 7, 111, 1940.

OXYTROPIS THOMSONI Benth

A note on *Oxytropis thomsoni*, Benth. Siddiqui, R.H, Basha, S. K., *Ind Jour Pharm*, 1944, 6, 83

PANDANUS ODORATISSIMUS Roxb.

Kewda Perfumes Vol IV, No 1—Sadgopal,—*Indian Soap Jour.*, July 1936

Kewda—Sadgopal,—*Soap Perfumery and Cosmetics*, May, 1937

Pandanus oil—Sadgopal, *Soap Perfumery and Cosmetics*, 10, 1937, 396
Essential oil from flowers of Kewda, *Pandanus odoratissimus*. S S Deshpande, *Calcutta University Press*, 1938

Kewda (*Pandanus odoratissimus*) Essential oil from S S Deshpande, *J Ind Chem Soc* 15(1938), 509

PAPAVER SOMNIFERUM Linn

A preliminary note on addiction to "Post" (unlanced capsules of *Papaver somniferum*) Chopra, R N, *Ind Med Gaz.*, 1930, 65, 361

Addiction to "Post" (unlanced capsules of *Papaver somniferum*) in India Chopra, R N, Grewal, K S, Chowhan, J S & Chopra, G S, *Ind Jour Med Res*, 1930, 17, 985

Addiction to "Post" Part II Chopra, R. N & Ghosh, N N, *Ind Jour. Med Res*, 1931, 19, 415

Opium and Albuminuria Chopra, R N & Bose, J P, *Ind Med Gaz*, 1931, 66, 299

Administration of Opium to infants in India Chopra, R N, Chopra, G S, *Ind. Med Gaz.*, 1934, 69, 489

Indian Opium Its principal groups of chemical constituent Rakshit, J.N., *Sci & Cult*, 1942, 8, 16

Opium alkaloids Recent development in India Mukhopadhyay, B K, *J Sci. Indl. Res*, 8A, 118, 1949

Papaverine from Indian opium Mukhopadhyay, B. K & Parthasarthy, C, *J. Sci Indl. Res.*, 1949, 8B, 23

PARIS POLYPHYLLIA Sns.

A preliminary note on the Glucosides from *Paris polyphylla* Sns. Chopra, R. N, Gupta, J C, Bose, B. C. & Chopra, I C., *Ind. Jour. Med Res*, 1942, 30, 103.

PEGANUM HARMALA Linn

Oil from Peganum harmala H. Ya Tropp, *Chem Abst.* 1598, 1936.

On Peganine isolated from the flowers of Peganum harmala L. A Rosenfile, D G , Kolesnikov, *Chem Abst* 9306, 1939

Amino neuteric determination of alkaloids in belladonna, lobelia herb and harmel roots, G Ya Kait, *Chem Abst* 2996, 1942

The action of Peganum harmala and of its alkaloids on the circulation. Arsenio Fraile Ovejero; *Chem Abst* 3214, 1947

Peganine from the blossoms and stems of Peganum harmala L. Remarks on the papers of E Spath on Peganine. A D. Rosenfeld and D G Kolesfeld and D.G Kolesnikov, *Ber* 28, 1704.

PENICILLIOPSIS CIAVARIAEFORMIS Solms Laub

Penicilliosis ciavariaeformis Solms Laub, A new record in India, Ganguly, D & Ghosh, T , *Sci & Cult.*, 1947, 12, 553.

PENTAPETES PHOENICEA Linn

A note on the development of the female gametophyte in Abroma angusta L and Pentapetes phoenicea L , Banerji, I , *Curr. Sci* 1941, 10, 30

PERANOSPORA PARASITICA

Medicinal action of Capsella bursa-pastoris and also of its parasites, Cystopus candidus and Peranospora parasitica W Harste, *Arch Pharm.* 266, 133, 1928

PEROWSKIA ATRIPLICIFOLIA Benth.

The Essential oil from the flower heads of Perowskia atriplicifolia, Benth, M G Rao, *Journal of the Indian Chemical Society, Calcutta* 1926, Vol III, p. 141.

PHALARIS MINOR Retz

Phalaris minor Retz A new promising winter fodder grass. Gandhi, R T , Wadhwa, N D , *Sci & Cult* , 1951, 17, 113

PHASEOLUS ACONITIFOLIUS Jacq

Studies in nutrition value of Indian vegetable foodstuff Pt III Nutritive values of Lentil-Lens esculenta, Moench., Cow pea - Vigna, catjang, Walp. and Aconite Bean - Phaseolus aconitifolius, Jacq , Niyogi, S P , Narayana, N & Desai, B G , *Ind Jour Med. Res* 1932, 19, 859

P RADIATUS Linn

Constituents of Phaseolus radiatus L Var Aureus Prain E Miyamichi and H Yamada, *J Pharm. Soc Japan*, 50, 1095, 1930.

Saponin glucoside in adzuki beans (Phaseolus radiatus L, Var. Aureus Prain) E. Miyamichi and S Onishi, *J. Pharm Soc, Japan.* 52, 168, 1932.

PHEBALIUM ARGENTUM Sm.

Phebalium argentum, occurrence of Psoralen in P. K Bose & Finlayson, *J Ind Chem Soc* 15 (1938), 516

PHYLLANTHUS EMBLICA Linn

Ascarbic acid contents of some Indian plant material M Damodaran and M Srinivasan, *Proc Ind Acad Sci* 2B, 377, 1935.

Indian Gooseberries (Phyllanthus emblica Linn) as a source of Vit C Giri, K V, *Ind Jour Med Res*, 1939, 27, 429

A new technique for propagating Aonla (Phyllanthus emblica) Singh, L B, *Sci & Cult*, 1952, 17, 345

P NIRURI Linn

Bitter principles of Phyllanthus niruri G V Krisana Murti and T R Seshadri, *Proc Ind Acad Sci* 24A, 357, 1946

Investigation of quinine in Phyllanthus niruri L Richardo Martin Serra, *Chem Abst* 1812, 1947

PHYSOCHLAINA PRAELTA (Don). Miers

Chemical investigation of Physochlaina praelta Miers Handa, K L, Nazir, B N, Chopra, I C & Jamwal, K S, *J Sci Indl. Res*, 1951, 10B, 182

Chemical investigation of the roots of Physochlaina praelta Miers Handa, K L, *J Sci Indl. Res*, 1951, 10B, 234

Chemical investigation of Physochlaina praelta Pt II Handa, K L & Onkar Nath Channa, *J Sci Indl. Res*, 1952, 11B, 505

PICRASMA EXCELSA Swartz

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum rutae (Ruta graveolens Linn), Butea monosperma Roxb, Melia azadirachta Linn, Punica granatum Linn, Picrasma excelsa Swartz Vernonia anthelmintica Willd, Cocos nucifera Linn (coconut) Cains, J F. & Mhaskar, K S, *Ind Jour Med. Res*, 1923, 11, 353

PICRORHIZA KURROA Royle ex Benth

Some common indigenous remedies, Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum, Chopra R. N. and Ghosh, S., *Ind Med Record* 55, 77, 1935

Comparative pharmacognosy of Gentiana kuroo Royle & Picrorhiza kurroa Royle. ex Benth. Bal, S N. & Datta, S C, *Ind Jour Pharm*, 1945, 7, 109.

Chemical Examination of *Picrorhiza kurroa* Benth. Part I Rastogi, R P, Vishwa Nath Sharma & Siddiqui, S, *J Sci Indl Res*, 1949, **8B**, 173.

Chemical investigation of *Picrorhiza kurroa* Benth Attempted synthesis of Kutkin Rastogi, R P, Sen, A B, *J I C S*, 1953, 7, 514

PIMPINELLA SAXIFRAGA Linn

Constituents of the roots of *Pimpinella saxifraga* Fritz, Wessely and Editha Nadler, *Monatsh*, 60, 141, 1922

Constituents of the roots of *Pimpinella saxifraga* Fritz, Wessely and Ferdinand Kallal, *Monatsh*, 59, 161, 1932

PINUS Linn

A plea for the distillation of Pine needle oil in India Puran Singh, *Indian Forester*, 1914, Vol XL, No 10

Oleum trebinthinae *Ind Jour Med Res*, 1920, 8, 125

Turpentine oil - H M Hulshy and E. P. Watson, *Quarterly Journal of the Indian Chemical Society*, 3, 1926, 258

The constituents of some Indian Essential Oils Part XVII - Derivatives of Abietic Acid Rau M Gopal and Simonsen J L, *Journal of Indian Institute of Science, Bangalore*, 1926, Vol 9, Part IV, pp 111-116 Continued from Indian Forest Records

The Pinene content of Indian Turpentine, Mulany H M & Watson, E R, *J Ind Chem Soc* 3 (1926), 258.

The preparation of Camphor from Pinene Mulany, H M & Watson, E. R, *J Ind Chem Soc*, 3 (1926), 253

The Pinene content of Indian Turpentine Mulany and Watson, *Journal of the Indian Chemical Society, Calcutta*, 1928, Vol III, p. 258.

Some pharmacological action of Sandal wood oil Vitivert oil and Pine needle oil Tameno Okanishi, *Chem Abst* 658, 1930

Vitamin C in pine needles Iyengar, N. K, Bose, B. C & Mukerji, B, *Ind Jour Med Res*, 1944, 32, 165

On the possibility of using oil of Turpentine for the treatment of Scabies Roy, D N & Ghosh, S M., *Ind Med Gaz* 1944, 79, 589.

Abietic acid from Indian Rosin Ghatak, N & Mitra, S P., *J Sci Indl Res* 1945-46, 4, 653

Studies in Indian Turpentine—production of p cymene (3) Sindhi, H K; *Sci. & Cult*, 1949, 15, 202

P EXCELSA Wal.

The constituents of some Indian Essential Oils Part II The Essential oil from the oleo-resin of *Pinus excelsa* Simonsen, J L and Rau M Gopal, *Indian Forest Records*, 1922 Vol IX, Part IV, pp. 6 - 12.

The leaf oil from *Pinus excelsa*. Simonsen, J.L. *Indian Forest Records*, Vol. IX. Part IX, 1922.

●Indian resin oil from *Pinus excelsa*. Patel, I.S. & Guha, P.C., *Curr Sci*, 1950, 19, 128.

P GERARDIANA Wall.

The constituents of some Indian Essential Oils Part IX—The Essential oil from the oleo-resin of *Pinus gerardiana*, Wall Simon- sen, J L, *Indian Forest Records*; 1923, Vol IX, Part VIII, pp. 5-8

Oil from *Pinus gerardiana* Chilgoza oil Part I. Hardikar, S.D. *J. Ind. Chem Soc* 5 (1928), 69.

P KHASYA Royle.

Turpentine oil and resin from *Pinus khasya* (Khasi Pine, Dieng Kesh or Saral). Saikia, B., Das, P.K. & Roy, B.K, *Curr. Sci.*, 1951, 20, 275

P. LONGIFOLIA Roxb.

Turpentine oil from *Pinus longifolia* (Chir)—Puran Singh; *Indian Forest Records*, 1913, Vol. IV, Part I.

The constituents of Indian Turpentine from *Pinus longifolia* Roxb. John Lionel Simonsen, *J. Chem Soc.* 570, 1920.

The Constituents of Indian Turpentine from *Pinus longifolia* Roxb Part I Simonsen, J.L., *Trans Chem Soc.* 1920, Vol. 117, p. 570.

The Constituents of Indian Turpentine from *Pinus longifolia* Roxb. Part II Simonsen, J L and Gopal Rao, M, *Trans Chem. Soc.* 1923, Vol 123, p. 549.

The Constituents of Indian Turpentine from *Pinus longifolia* Roxb. Part III. Simonsen, J L, *Trans Chem Soc.* 1923, Vol. 123, p 2642.

The Constituents of Indian Turpentine, from *Pinus longifolia* Roxb. John Lionel Simonsen, *Jour. Chem Soc.* 2642, 1923

The Constituents of some Indian Essential Oils Part XVII Abietic Acid from the resin of *Pinus longifolia*, Roxb (Preliminary note) Rau M Gopal and Simonsen, J L, *Indian Forest Records*, 1925, (Chemistry Series) Vol. XI Part VI.

The Constituents of Indian Turpentine from *Pinus longifolia*, Part IV Pillay, P P and Simonsen, J.L. *Journal of the Indian Institute of Science, Bangalore*, 1928, Vol. 11A, Part XV pp. 200-205.

The Constituents of Indian Turpentine from *Pinus longifolia* Roxb. John Owen and John Lionel Simonsen, *J. Chem. Soc.* 3001, 1931.

On the carenes in Indian Turpentine Oil from *Pinus longifolia*, Roxb. Verghese, James, *Curr. Sci*, 1950, 19, 212.

Turpentine oil from *Pinus longifolia* Roxb. James Verghese & Gulati, K C., *J. Sci Indl. Res.*, 1951, 10A, 112.

Liquid phase production of *p*-cymene from Indian Turpentine, oil of *Pinus longifolia* (Roxb.) & 3-carene. James Verghese & Lourdu M. Yeddanapalli; *J. Sci Indl. Res.*, 1951 10B, 100.

Studies in oil of *Pinus longifolia* Roxb. II. Aromatization of 3-carene to *p*-cymene James Verghese & Lourdu M. Yeddanapalli, *J. Sci. Indl Res.*, 1952, **11B**, 36.

Studies in oil of *Pinus longifolia* Roxb III—Two directional Cyclopropane ring of 3-carene in Sulphur Aromatization. Yeddanapalli, L.M., James Verghese, *J S I R*. 1953, **3B**, 121.

Studies in oil of *Pinus longifolia* Roxb II—Isomerization of 3-carene & Terpinene. Yeddanapalli L M., Krishnan T S., Verghese J; *J S I R* 1953, 38, 122

PIPER BETLE Linn.

Studies in human nutrition Pt IV Availability of Calcium ingested in the process of Chewing Betel-leaves with lime Basu, K.P.; Basak, M N & De, H.N., *Ind Jour Med Res*, 1942, 30, 309.

Tipburn of Piper betel in the Central Provinces. Asthana, R P. & Mahmud, K A, *Curr. Sci.*, 1944, 13, 234

The areca and the betel. Andre Mericier, *Chem. Abst* 956, 1946.

On the control of *Rhizoctonia*, root-rot of Pan (Piper betel L.) Chowdhury, S, *Sci. & Cult.*, 1948, 13, 507.

Propagation of Pan (Piper betel L.) in the Northern districts of the Central Provinces. Mahmud, K A., *Sci. & Cult*, 1950, 15, 324.

The antibacterial principle of Betel leaf. A preliminary note. Pai, Narayana & (Miss) Irani, R J., *Ind Med Gaz.*, 1950, 85, 302.

P. CUBEBA Linn

Constitution of some Indian essential oils, essential oil from the fruit of Piper cubeba Linn. B S Rao, V P. Shintre and J L. Simonsen, *J Soc. Chem Ind.* 74, 92, 1928.

The constituents of some Indian Essential Oils, Part XXIII—The Essential Oil from the fruits of Piper cubeba, Linn Rao B Sanjiva, Shintre, V P and Simonsen J L, *Journal of the Indian Institute of Science, Bangalore*, 1928, Vol **11A**, Part XV-1, pp 187-194.

P NIGRUM Linn

The Nitrogen complex of Indian foodstuffs condiments. Pt I. Black pepper (*Piper nigrum*) Narasimhamurthy, G. & Ranganathan, S, *Ind. Jour. Med Res*, 1937, 25, 373.

PISTACIA INTEGERRIMA Stew.

Chemical investigation on Kakra Singi Karimullah, Subba Rao, V. & Uma Shankar, *J. Sci. Indl Res.*, 1944-45, 3, 423.

A note on the crystalline constituents of (*Pistacia integerrima* Stew. N O. Anacardiaceae) Kakra Singi. Ghose, S.K., *Sci. & Cultr.*, 1945, 11, 46.

Chemical investigations of Kakra Singi (*Pistacia integerrima*). The essential oil. Karimullah & Uma Shankar; *J. Sci. Indl. Res.*, 1946, 5, 60.

Essential oil from the galls of *Pistacia integerrima*. Baslas & Deshpande, *J. Ind Chem. Soc.* 27 (1950), 441.

P TEREBINTHUS Linn.

Ethereal oil from the resin of *Pistacia terebinthus*. A Tsatsas, *Chem. Abst* 5989, 1939

PITHECOLOBIUM SAMAN Benth

Composition of 'Rain Tree' fruit Rao, V R., Noshir, N, *Curr. Sci*, 1946, 15, 250.

PLANTAGO MAJOR Linn.

A holloside extracted from the seeds of *Plantago major* L and *Plantago ovata* Forsk. (P *Isphagulata* Roxb), N Wathez and H. Hans, *Chem Abst* 4849, 1945.

Wound healing preparations from plantain (*Plantago major*) leaf. R.K Aliev, *Chem Abst.* 2210, 1947

P. OVATA Forsk

Plantago ovata, ispaghul in chronic diarrhoeas and dysenteries. Chopra, R N, *Ind Med. Gaz*, 1930, 65, 428

A holloside extracted from the seeds of *Plantago major* L and *Plantago ovata* Forsk (P *isphagulata* Roxb), N Wathez and H Hans, *Chem Abst* 4849, 1945

PLUMBAGO ROSEA Linn

Chemical examination of the root bark of *Plumbago rosea* Linn. M G Tumin Katti and V N Patwardhan, *J. Ind Inst Sci.* **15A**, 9, 1932

P ZEYLANICA Linn.

The pharmacological action of *Plumbago zeylanica* and its active principle (Plumbagin). Bhatia, B.B & Lal S., *Ind. Jour Med. Res*, 1933, 20, 777

PLUMERIA ACUTIFOLIA Poir

The aerial roots of *Plumeria acutifolia* Poir. Banerji, I, *Curr Sci* 1943, 12, 85

PODOPHYLLUM Linn.

Components of Indian *Podophyllum*. Seshadri, T.R. & Subramanian, S.S., *J. Sci. Indl. Res* 1950, **9B**, 137.

A revision of the structure of Podophyllotoxin and picropodophyllin Chatterjee, R., Chakravarti, S C, *Sci & Cult* 1951, 17, 136.

POLYGALA CHINENSIS Linn

Polygala chinensis Linn as substitute for official senega Hossain, T., Guha, R.C. & Mukerji, B.; *Sci. & Cult.* 1943, 9, 167

The Senegas of Indian Market. Gupta, B. & Bal, S.N., *J. Sci. Indl Res*, 1952, **11B**, 116.

POLYGONUM Linn.

Anthracenic derivatives in the genera Polygonum and Rumex. Maurin, E., *Bull. Sci. Pharmacol.* 33, 138, 1926.

PONGAMIA GLABRA Vent.

The oil of Pongamia glabra. Beal, C D. and Katti, M.C.T.; *J. Am. Pharm Assoc.* 14, 1086, 1925

Pongamia glabra Leaf-Gall-Farmer. Cherian, M C., *Curr. Sci.*, 1934-35, 3, 564.

Some common indigenous remedies, Picrorhiza kurrooa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum. Chopra, R.N. and Ghosh, S., *Ind. Med. Record* 55, 77, 1935.

Pongamia glabra. A note on the occurrence of Behenic acid in the oil from the seed of. Manjunath, B L. & Shanker Rao, M.S., *J. Ind. Chem Soc* 15(1938), 653.

Pongamia glabra. A note on the occurrence of free fatty acid in the cake of Subba Rao, N.V. & Veerbhadra Rao, J., *J. Ind. Chem. Soc.* 175(1940), 26.

POTENTILLA ANSERINA Linn. & P. ARGENTEA Linn

The comparative activity of Potentilla anserina L. and Potentilla argentea L. on the isolated guinea pig uterus. Youngken, H and Fischer, E.B , *Am J. Pharm.* 114, 417, 1943.

PRANGOS PABULARIA Lindl.

Essential oil of Prangos pabularia Lindl. L. Grach , *Chem. Abst.* 7529, 1940.

PREMNA INTEGRIFOLIA Linn.

Premna integrifolia Linn , Chemical investigation of. Basu, N K. & Joneja, A.N., *Ind. Jour Pharm* 1949, 11, 191.

PRINSEPIA UTILIS Royle.

The fatty oil from the seeds of Prinsepia utilis. Puntambekar, S V. ; *J. Ind Chem. Soc.* 19(1942), 183

PRISTIMERA INDICA Willd.

Pristimerin, the antibacterial principle of Pristimera indica Isolation, Toxicity and antibacterial action. Bhatnagar, S.S., Divekar, P V., *J. Sci. Indl. Res* , 1951, 10B, 56.

Pristimera indica—A source of Dulcitol. Bhatnagar, S.S , & Divekar, P.V.; *J. Sci. Indl. Res.*, 1951, 10B, 117.

PROSOPIS JULIFLORA D.C.

Antibiotic activity of ext of Prosopis juliflora. Shankarmurty, P. & Siddiqui, S., *J. Sci. Indl. Res.*, 1948, 7B, 188.

PRUNUS ACUMINATA Hk. f.

Chemical investigation of the bark of *Prunus acuminata*. Chakravarti, D., Kundru, N., *Sci & Cult.* 1948, 14, 36.

P. AVIUM Linn.

Pharmacological studies on *Prunus avium*. Detection and determination of carbohydrates present in an infusion of stems of *Prunus avium* L. Gaetano Di Manggio, *Chem Abst.* 566, 1947

P. CERASUS Linn.

Indian Wild Cherry bark Kapoor, L. D. & Handa, K. L., *Cur. Sci.*, 1948, 17, 54.

P. NEPALENSIS Koch.

Chemical investigation of the bark of *Prunus nepalensis* Chakravarti, D. & Momen, S. A., *Sci. & Cult.*, 1950, 15, 329.

P. PUDDUM Roxb.

Isolation of a new flavone from the bark of *Prunus puddum* (N. O. Rosaceae), Chakravarti, D. & Ghosh, R. P., *Sci. & Cultr.*, 1943, 8, 463.

Isolation of a new iso-flavone from the bark of *Prunus puddum* (N. O. Rosaceae), Chakravarti, D. & Bhar, C. N., *Sci. & Cultr.*, 1943, 8, 498.

Identity of Puddumetin and Genkwanin, Venkataraman, K., *Sci & Cultr.*, 1943, 9, 45.

Crystalline components of the bark of *Prunus puddum* Roxb. Part I. Identity of Puddumetin with Genkwanin. Chakravarti, G. D. & Ghosh, R. P., *J. Ind. Chem Soc* 21 (1944) 171

Prunus puddum Roxb. Crystalline components of Chakravarti & Behari, *J. Ind. Chem Soc* 22 (1945), 301.

Prunus puddum Roxb. Crystalline components of the bark of. Chakravarti, Kundu & Ghosh, *J. Ind Chem Soc* 25 (1948), 329

Prunus puddum. Isolation of Sakuranin from the bark of. Chakravarti & Sen *J. Ind. Chem. Soc* 27 (1950), 148

PSIDIUM GUAJAVA Linn

Effect of hormones on the rootage of *Psidium guajava* L. Singh, S. N., *Sci. & Cult.*, 1950, 16, 198

PSILOTUM TRIQUERUM Sw.

On the occurrence of *Psilotum* Sw. in the East Godawari Distt. Venkateswarlu, V., *Sci. & Cultr.*, 1943, 9, 166.

PSORALEA CORYLIFOLIA Linn.

Psoralea corylifolia (Babchi). Its constituents, their pharmacological action and therapeutic properties. Chopra R. N., Chatterjee, N. R., *Ind. Jour. Med. Res.*, 1927, 15, 11.

Psoralea corylifolia. Chemical examination of the seeds of Part I Jois, H. S., Manjunath, B. L. & Venkata Rao, S., *J. Ind Chem Soc.* 10 (1933), 41

Oil of *Psoralea corylifolia*—T. R. Seshadri and C. Venkat Rao, *Proc. Indian Acad. Sci. Sect 5A*, 1937, 351

Components of *Psoralea corylifolia* Linn. T. R. Seshadri and C. Venkata Rao, *Proc. Ind Acad. Sci 5A*, 1937, 351

Nicotinic acid in Bavchi (*Psoralea corylifolia* - Bharucha, F. R., *Sci & Cult*, 1941, 7, 169

Histological study of fruit and seed of *Psoralea corylifolia* Linn. Prasad, G. S., *Ind Jour Pharm*, 1944, 6, 61.

Chemical examination of the seed of *Psoralea corylifolia* Linn. (Bebchi) Kamler Kinkar Chakravarti, Anil Kumar Bore & Salimuz-zaman Siddiqui, *J. Sci. Indl. Res*, 1948. 7B, 24.

PTERIS ACQUILINA Linn.

Poisonous properties of bracken (*Pteris acquilina*) G. D. Shearar, *Chem Abst* 6643, 1946

PTEROCARPUS MARSUPIUM Roxb

The Phlobatannins of Kino and buteagums G. V. Krishnamurti and T. R. Seshadri, *Proc. Ind Acad. Sci 22A*, 134, 1945

A preliminary note on the action of *Pterocarpus marsupium* Roxb. on blood sugar Ojha, K. N., Parba, P. R. & Venkatachalam K., *Ind Jour Pharm*, 1949, 11, 188

PUERARIA THUNBERGIANA Benth

Kudzu vine (*Pueraria thunbergiana*) as fodder. Dhar, N. D. & Goswami, M. N., *Sci. & Cult*, 1950, 16, 77

PUNICA GRANATUM Linn.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum rutae (*Ruta graveolens* Linn.), *Butea monosperma* Roxb., *Melia azadirachta* Linn., *Punica granatum* Linn., *Picrasma excelsa* Swartz, *Vernonia anthelmintica* Willd., *Cocos nucifera* Linn. (coconut) Cains, J. F. & Mhaskar, K. S., *Ind Jour. Med Res*, 1923, 11, 353.

PUTRANJIVA ROXBURGHII Wall

Putranjiva roxburghii Wall The oil from the seeds of Krishna, S. & Puntambekar, S. V.; *J. Ind Chem Soc* 8 (1931), 301.

Development of the Microspores and the Nuclear behaviour in the Tapetal Cells of *Putranjiva roxburghii* Wall. Dutt, M. K., *Sci. & Cultr.*, 1943, 8, 309.

QUERCUS Linn.

Electric treatment of horse chestnuts and acorus. Hayao Muraoka; *Chem Abst.* 3268, 1936.

Q. DILATATA Lindl., Q ILEX Linn., Q INCANA Roxb.

Quercus incana Roxb , *Q. dilatata* Lindl. and *Q. ilex* Linn. Some Indian acorn oil Puntambekar, S. V. & Krishna, S , *J. Ind Chem. Soc.* 11 (1934), 721.

A note on the variable tannin content of the wood of *Quercus incana* H. S Chaturvedi & Watson, E. R , *J Ind Chem Soc.* 3 (1926), 211

QUISQUALIS INDICA Linn.

Chemical study of the seeds of *Quisqualis indica*. Composition of the crude oil Chit Fang Hsu and Pao Him King, *J. Chinese Pharm. Assoc* 2, 132 1940

RANDIA DUMETORUM Lamk.

Chemical and Pharmacological examination of *Randia dumetorum* N O. Rubiaceae Hardikar, S W & Mohiuddin, M. G , *Ind Jour. Med Res* , 1937, 25. 131.

RAUWOLFIA Linn

Combined Digitalis and Rauwolfia poisoning in a human subject. De, M. M. & Tarapaido Chatterjee, *Ind Med Gaz* 1941, 76, 724.

Rauwolscine & derivatives. Mookerjee, *J. Ind Chem Soc.* 20, (1943), 15

Pharmacognostic studies in the leaves of a few species of Rauwolfia. Determination of Palisade Ratio and Stomatal Index value Datta, S C , *Ind Jour. Pharm* , 1944, 6, 77

Rauwolfia roots, Pharmacognostic studies on commercial varieties Datta, S C.; *Ind Jour Pharm* , 1949, 11, 105

A preliminary note on the excretion of Rauwolfia total alkaloids in urine Gupta, J. C , Roy P K , Ray, G K & Ganguly, S. C., *Ind. Jour. Med Res.*, 1950, 38, 67

Rauwolscine methiodide Chatterjee, *J. Ind Chem Soc* 28 (1951) 31

Iso Rauwolscine & its HCl, perchlorate, picrate and Chloroplatinate Chatterjee, *J. Ind Chem. Soc* 28 (1951) 31, 33

R. CANESCENS Linn.

Alkaloids of *Rauwolfia canescens* Linn. Part I. (Miss) Mookerjee, A., *J. Ind. Chem Soc.* 18 (1941), 33

The Alkaloid of *Rauwolfia canescens* Linn. Part I. (Miss) Mookerjee, A , *J. Ind. Chem. Soc* 18 (1941), 485.

Function of Rauwolscine in *Rauwolfia canescens* Linn. (Miss) Mookerjee, A., *Sci. & Cultr.*, 1942, 8, 40.

The Alkaloids of *Rauwolfia canescens*. Mookerjee, A; *J. Ind. Chem. Soc.* 20 (1943), 11.

Rauwolfia canescens. Constitution of the alkaloid (Rauwolscina) Mookerjee, *J. Ind. Chem. Soc.* 23 (1946), 6.

Studies on the Pharmacology of Rauwolscine, the alkaloid of *Rauwolfia canescens* Linn. Mukherjee, J. N.; *Sci. & Cult.*, 1953, 7, 338

Yobyrine, the selenium dehydrogenation product of Rauwolscine, the alkaloid of *Rauwolfia canescens*. Chatterjee, A., Pakrashi, S. *Sci. & Cult.*, 1953, 9, 443

R. SERPENTINA Benth

Chemical Examination of the Roots of *R. serpentina*. Siddiqui-Siddiqui, *J. Ind. Chem. Soc.*, 8, 667 (1931)

Chemical examination of roots of *Rauwolfia serpentina* Benth By Salimuzzaman Siddique & Rafat Hussain Siddiqui; *J. Ind. Chem Soc.* 8 (1931), 667.

The alkaloids of *Rauwolfia serpentina* Benth. Part I. Salimuzzaman Siddiqui & Rafat Hussain Siddiqui; *J Ind Chem Soc* 9, (1932), 539.

The Alkaloids of *R serpentina* Benth , Part I Siddiqui-Siddiqui, *J. Ind Chem. Soc* 9, 539 (1932),

R. serpentina. van Itallie-Steenhauer, *Arch. Pharm.* 270, 313 (1932),

The pharmacological action of an alkaloid obtained from *Rauwolfia serpentina*, Benth A preliminary note. Chopra, R N., Gupta, J C. & Mukerji, B., *Ind Jour Med. Res* , 1933, 21, 261

The Alkaloids of *R serpentina* Benth , Part II, Studies in the Ajmaline-Series Siddiqui-Siddiqui, *J Ind Chem Soc.*, 12, 37 (1935)

Die Wirkung von Rauwolfia auf das Herz, Hartog *Arch Int. Pharmacodyn Therap* 51, 10 (1935).

The alkaloid of *Rauwolfia serpentina* Benth. Part II Studies in the Ajmaline Series. Siddiqui & Rafat Hussain Siddiqui, *J Ind. Chem. Soc.* 12 (1935), 37

Sur un nouveau paralysant electif des vasoconstricteurs adrenal-inosensible l'ajmaline, alcaloïde cristallise de *R. serpentina* Raymond-Hamet, *Bull. Acad Med* 115, 452 (1936)

The Action of some Drugs on Fibrillation of the Heart (Rauwolfia). van Dongen, *Arch Int Pharmacodyn Therap.* 53, 80 (1936).

Ajmaline, Alkaloid from *R. serpentina* of selective Paralyzing Action on Vaso-constrictors Reactive to Adrenaline Raymond-Hamet, *Bull Sci-pharmacol*, 43, 364 (1936)

The Action of Ajmaline on Nerve Impulses Chopra-Das-Mukerjee, *Ind J Med Res* 24, 1125 (1937).

A note on the Alkaloids of *R. serpentina* Benth. Siddiqui, *J Ind. Chem Soc* , 16, 421 (1939).

Rauwolfia serpentina Benth. A note on the alkaloids of. Salimuzzaman Siddiqui, *J. Ind. Chem Soc* 16 (1939), 421.

Influence de la serpentinine sur les effets de l'adrenaline, de l'occlusion carotidienne et de la faradisation du pneumogastrique. Raymond-Hamet, *C. r.* 211, 414 (1940).

Sur quelques propriétés physiologiques de la serpentine, alcaloïde cristallisé du *R. serpentina* Benth Raymond-Hamet, *C. r Soc. biol.* 134, 94 (1940).

Effets intestinaux d'ajmaline pure. Raymond-Hamet, *C. r Soc. biol.* 134, 369 (1940).

A method of assay for *Rauwolfia serpentina*, Benth. Mahadev Lal Schroff & Rattan Lal Bhatia, *Ind Jour Pharm*, 3, 59, 1941.

A preliminary note on the pharmacological action of the alkaloids of *Rauwolfia serpentina* Chopra, R. N. & Chakravarti, M., *Ind Jour Med Res*, 1941, 29, 763

Alkaloids of *Rauwolfia serpentina*, A comparative study of pharmacological action and their role in experimental hypertension Chopra, R. N., Bose, B. C., Gupta, J. C. & Chopra, I. C., *Ind Jour. Med. Res.*, 1942, 30, 319

Preliminary observations on the use of *Rauwolfia serpentina* Benth. in the treatment of mental disorders Gupta, J. C., Deb, A. K. & Mahali B. S., *Ind Med Gaz*, 1943, 78, 547

Comparative pharmacology of the total alkaloids of *Rauwolfia serpentina* Benth obtained from Bengal, Bihar & Dehra Dun. Gupta, J. C., Kahali, B. S., *Ind Jour Med Res*, 1943, 31, 215.

Hypnotic effect of *Rauwolfia serpentina* The principle underlying this action, its probable nature Chopra, R. N., Gupta, J. C., Bose, B. C. & Chopra, I. C., *Ind Jour Med Res.*, 1943, 31, 71

Pharmacological action of alkaloid of *R. serpentina* Benth. Pt I Neo-ajmaline & iso-ajmaline Bhatia, B. B. & Kapur, R. D., *Ind Jour Med Res.*, 1944 32, 177

The Hypnotic Effect of a Resin-Fraction from the Root of *Rauwolfia serpentina* Benth obtained from Dehra Dun Gupta-Kahali-Dutt, *J Ind Med. Res*, 32, 183 (1944)

Effets tenseurs vasculaires d'un des alcaloïdes colorés du *R. serpentina* Serpentinine Raymond-Hamet, *C. r* 223 927 (1946).

Rauwolfia serpentina Benth Comparative chemical investigation of some of the constituents of the drug obtained from different sources and isolation of the active resin Dutt, A., Gupta, J. C., Ghosh, S. & Kahali, B. S., *Ind Jour Pharm*, 1947, 9, 54.

A note on the Hypnotic Principle of *Rauwolfia serpentina*. Gupta, J. C., S. Ghosh, A. T. Dutta & B. S. Kahali, *J Amer. Pharmac. Assoc Sci Ed* 36, 416 (1947).

Pharmacological action of alkaloids of *R. serpentina* Benth Pt. II Total alkaloidal exts of Bihar & Dehra Dun varieties. Kapur, R. D., *Ind. Jour Med. Res.*, 1948, 36, 57.

Chemistry of Ajmaline (*Rauwolfine* of van Itallie and Steenhauer). Mukherji, Robinson-Schlittler, *Exper.* 5, 215 (1949)

Ajmaline, serpentine and Serpentinine. Raymond-Hamet, *C. R.* 229, 1165 (1949).

A Clinical Trial of *R. serpentina* in Essential Hypertension Wakil, *Brit Heart J* 11, 350 (1949).

Serpentina alkaloids. Bakshi, V. M , *Ind. Jour Pharm* , 1950, 12, 172.

Ueber das alkaloid Serpentin aus *R serpentina* Benth Schlittler-Schwarz, *Helv.* 33, 1463 (1950),

Structure of Reserpine. Neuss, Boaz, Forbes, *J A. C S* 75, 4870 (1950)

Ex-cretion of Rauwolfia total alkaloids in Urine. Gupta-Roy-Ray-Ganguly, *Ind. Jour. Med. Res.* 38, 67 (1950)

Rauwolfia serpentina in essential hypertension Chakravarty, N K , Rai Chaudhuri, M N & Chaudhuri, R N , *Ind Med Gaz* , 1951, 86, 348.

A new alkaloid from the root of Rauwolfia serpentina Benth. Chatterjee, A & Sukumar Bose, *Sci & Cult* , 1951, 17, 139

Ueber Rauwolfia serpentina, eine neue Droge in der Behandlung des Hochdrucks. Mayer, *Deutsche Apothekerzeitung* 1952, 435

Rauwolfinine, the new Alkaloid from the Root of *R serpentina* Bose, *Science and Culture*, 18, 98 (1952)

Ueber *R. serpentina*, eine neue Droge in der Behandlung des Hochdrucks Meyer, *Deutsche Apoth Ztg* 1952, 435

Zur Konstitution des Serpentins. Bader-Schwarz, *Helv.* 35, 1594 (1952).

Rauwolfia in Hypertension Mazumdar-Mukherji, *J. Ind. Med. Ass.* 19, 362 (1952).

The Use of *R. serpentina* in High Blood Pressure. Bhatia, *J. Ind Med. Ass* 21, 262 (1952)

Die Behandlung der chronischen arteriellen Hypertonie (II. Teil) Arnold, *Therapie der Gegenwart* 1952, 167.

Behandlung der Hypertonie mit der Indischen *R. serpentina*. Vida, *Die Medizinische* 1952, 1157

Raupin, ein neues Alkaloid aus *R serpentina* Bodendorf-Eder, *Naturwiss* 40, Heft 12, (1953)

Sarpagin, ein neues Alkaloid aus *R serpentina* Stoll-Hofmann, *Helv.* 36, 1143 (1953)

The Constitution of Ajmaline Chatterjee-Bose, *Exper.* 9, 254 (1953)

The Constitution of Reserpine Furlenmeier, Lucas, MacPhillamy, Mueller Schlittler, *Exper.* 9, 331, (1953)

On the Constitution of Réserpine from *R serpentina* Dorfman, Huebner, MacPhillamy, Schlittler, St. Andre; *Exper* 9, October 15th. (1953).

Alkaloids of Rauwolfia serpentina, The Characterization of Reserpine and its Hydrolysis Products. Klohs, Draper, Keller, Petrcek, *J. A. C. S.* 75, 4867 (1953).

Indian Medicinal Plants in the Treatment of Hypertension
Chakravarti, *Brit. Med J.* 1953, 1390.

Zur Pharmakologie des Reserpins, eines neuen Alkaloids aus *R. serpentina*. Bein, *Exper* 9, 107 (1953)

Sedative and Hypnotic Action of Reserpine in Unanesthetized Animals
Plummer, Barrett, Wagle, Yonkman *Fed Proc* 12, 357 (1953)

Pharmacological Effects of Reserpine, a new Crystallized pure Alkaloid from *R. serpentina* in the Dog Trapold, Osborne, Yonkman, *Fed. Proc* 12, 213 (1953)

Wirkung von Reserpin (Serpasil) auf isolierte Kreislauforgane.
Tripod-Meier, Submitted to *Arch Internat Pharmacodyn*

The Cardiovascular Effects of Reserpine (Serpasil) Trapold, Plummer, Yonkman, Submitted to *J Pharmacol exp. Therap*

Combinations of Drugs in the Treatment of Essential Hypertension (pure Reserpine=Serpasil) Wilkins, *The Mississippi Doctor*, 1953, 359 (April 5th)

The Use of *R. serpentina* in Hypertensive Patients (*Rauwolfia* extracts). Wilkins-Judson, *New England J. Med* 248, 48 (1953)

Medical Management of Arterial Hypertension (*Rauwolfia* extracts), Meilman, *New England J. Med.* 248, 936 (1953)

The Action of *Rauwolfia Serpentina* on Vasomotor Reflexes Ray, Roy, Dasgupta and Werner, *Arch exper Path. u Pharmacol Bd.* 219, S 310-314 (1953)

The Sympathicolytic Activity of *Rauwolfia Serpentina* (Benth) Ray, Roy, Dasgupta and Warner, *Ind. J Med. Sciences*, 7, No 5, 229-235, (5/53).

REVENELIA SESSILIS

Lysis in the germination of the Urediospores of *Revenelia sessilis*
Mathur, R L, *Sci & Cult*, 1952, 17, 380.

RHAMNUS Linn.

Further studies on Cathartic action in mice, Senna, Aloe Cascara and bile salts Lloyd W Hazleton and Kathleen D Talbert, *J Am. Pharm Assoc* 33, 170, 1944.

R. VIRGATUS Roxb

Indian Cascara, *Rhamnus virgatus* Roxb Chopra, I C, Kohli, J D. & Handa, K L, *Ind. Jour Med. Res.*, 1950, 38, 473.

RHEUM Linn.

Assay of anthracene purgatives by the estimation of the contents of Hydroxymethylanthraquinone Pt I Rhubarb. Ghosh, S., Gupta, J. C, & Mahali, B S., *Ind Jour Med Res.* 1945, 32, 129

R. EMODI Wall.

Rhapoticin and anthraquinone derivatives from *Rheum emodi* Wall. (Indian or Himalayan Rhubarb). M. Ghouse and M. C. Tumin Katti, *J. Ind. Inst Sci.*, 16A, 1, 1933.

A note on the essential oil from the Rhizomes of *Rheum emodi* Wall. M. Ghouse Mohiuddin, *J. Ind. Inst. Sci.* **18A**, 134, 1935.

Some common indigenous remedies *Picrorhiza kurroa*, *Erythrina indica*, *Sansevieria zeylanica*, *Pongamia glabra*, *Hygrophila spinosa*, *Bryophyllum calycinum*, *Rheum emodi*, *Solanum indicum* Chopra, R.N. & Ghosh, S, *Ind. Med. Record* 55, 77, 1935

A note on the Essential oil from the Rhizomes of "*Rheum emodi* Wall". Mohiuddin M Ghouse, *Journal of the Indian Institute of Science, Bangalore*, 1938, Vol **18A**, Part XVII, pp 134-135.

Indian Rhubarb as substitute for "official" Rhubarb Mukerji, B., *Curr. Sci*, 1943, 12 275.

Examination of Indian Rhubarb-*Rheum emodi* Wall Ray, G. K., Guhar, R. C., Bose, A. B. & Mukerjee, B, *Ind Jour. Pharm*, 1944, 6, 55.

Indian Rhubarb Hocking, M, *Ind. Jour Pharm*, 1945, 7, 89

Studies on Indian Rhubarb, *Rheum emodi* Heber W Yongken, *J Am. Pharm. Assoc* 35, 145, 1946.

RICINUS Linn.

The Growth of effects of thiamine Hydro chloride ascarbic acid and phytohormones on beladonna and ricinus Loxis C Zoft, *J Am Pharm Assoc.* 29, 487, 1940.

R. COMMUNIS Linn.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency XVIII Cathartics *Oleum Ricini*, *Oleum Tiglli*, *Aloe*, *Succies Acalyphae* Chans, J F, Mhaskar, K S, *Ind. Jour Med Res.* 1923, 11, 103

On the description of the inflorescence of *Ricinus communis* Linn Datta, R M, *Sci & Cult*, 1946, 10, 431.

ROSA Linn

Rose Oil - Chatterjee, N G.; *Indian Soap Journ* 3, 1937, 293.

Rose Distillation - P E O R Jan. 21, page 14 *Journal of Indian Industries and Labour*, Vol 2, Part I, page 86.

RUMEX Linn.

Anthracenic derivatives in the genera *Polygonum* and *Rumex* E. Maurin, *Bull Sci. Pharmacol* 33, 138, 1926

RUTA GRAVEOLENS Linn.

The correlation between chemical composition of anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency *Santonin*, *Oleum rutae* (*Ruta graveolens* Linn), *Butea monosperma* Roxb, *Melia azadirachta* Linn, *Punica granatum* Linn., *Picrasma excelsa* Swartz., *Vernonia anthelmintica* Willd., *Cocos nucrifera* Linn. (coconut). Cains, J F & Mhaskar, K S., *Ind Jour Med. Res.*, 1923, 11, 353.

SACCHARUM SPONTANEUM Linn.

Wilt Rot in grasses I, Involution, *Saccharum spontaneum* L. Srivastava, R K, *Sci. & Cult*, 1952, 17, 529

SALIX Linn

The physiological significance of the glucosides in *Aesculus* and *Salix* Gerhard Kerstan, *Chem Abst* 6772, 1934

SALVADORA OLEOIDES Dcne.

Fat from *Salvadora oleoides*, Khakanfot Patel, C K, Iyer, S N, Sudborough, J. J. & Watson, H E, *J Ind Inst Sci* 9A, 117, 1926

SAMBUCUS Linn

Presence of alkaloids in *Sambucus* species Henri Yardin, *Chem Abst* 5723, 1936

SANSEVIERIA ZEYLANICA Willd

Some common indigenous remedies, *Piciorhiza kurroa*, *Erythrina indica*, *Sansevieria zeylanica*, *Pongamia glabra*, *Hypogophila spinosa*, *Bryophyllum calycinum*, *Rheum emodi*, *Solanum indicum*. Chopra, R N & Ghosh, S, *Ind Med Record* 55, 77, 1935

A Phytochemical study of *Sansevieria zeylanica* Stanley Scheindlin & Austin A Dodge, *Am J Pharm* 119, 232, 1947

SANTALUM ALBUM Linn

A further note on the oil value of some sandalwoods from Madras Puran Singh, *Indian Forester*, 1915, Vol XII, No 4

West Australian Sandalwood Oil Rao B Sanjiva & Sudborough J J, *Journal of the Indian Institute of Science, Bangalore*, 1923, Vol 5 Part XII, pp 163-176

Some pharmacological action of Sandalwood oil, Vitivert oil and Pineneedle oil Tamenno Okanishi, *Chem Abst* 658, 1930

Historical regeneration of Sandal (*Santalum album* Linn) and its significance *Curr Sci*, 1934-35, 3, 251

East Indian Sandal Wood Oil A.E Bradfield, Penfold, A R & Simonsen, J.L, *Journ. Chem Soc* 1935, 309

The wax from the leaves of Sandal (*Santalum album* Linn) Chibnall, A C., Piper, S H, H A Cl, Mangouri, Williams, E F & Iyenger, A V.V., *Biochem J.* 31, 1981, 1937.

Santalum album Chemical examination of the oil from the seeds Madhurnath, M.K & Manjunath, B L, *J. Ind. Chem. Soc.* 15, (1938), 389

The white Sandal (*Santalum album*). Majumdar, G.P., *Sci. & Cultr.* 1941, 6, 492.

Saussurea lappa (Kuth root) in Pharmacology and Therapeutics Chopra, R N. & De, Premankur., *Ind Jour Med Res.*, 1929, 17, 351.

Chemical examination of roots and leaves of Saussurea lappa Part I, Sudhamoy Ghosh, Nibarranjan, Chatterjee & Ashutosh Dutta, *J. Ind. Chem. Soc.* 6(1929), 517.

Root oil of Saussurea lappa, Clarke Establishment Antoine Chris, *Chem. Abst.* 1553, 1937

A histological study of the root of Saussurea lappa C.B. Clarke, Prasad, S, *Ind Jour. Pharm.*, 1945, 7, 81.

Chemical Examination of roots of Saussurea lappa, Clarke, Pt. I On the reported isolation of the alkaloid "Saussurine" Salooja, K C, Vishwa Nath Sharma & Salimuzzaman Siddiqui, *J. Sci. Indl. Res* 1950, 9B, 1.

SCIRPUS GROSSUS Linn

Studies in enzymes. Part II. Purification of amylose from Kaseru (Scirpus grossus Linn) Shukla, J P, *J Ind Chem Soc* 19(1942), 121

SCOPOLIA Jacq.

The differentiation of Scopolia and Belladonna leaves. D Markovic, *Chem Abst* 10, 1, 9, 1944

SECALE CEREALE Linn.

The hydrolysis of ptyic compounds derived from seeds of hemp, horse bean, horse chestnut flax, wheat and embryos of rye W Jarosza, *Chem. Abst* 5501, 1934

SECURIGERA SECURIDACA Linn.

The pharmacological action on the circulatory system of a bitter principle isolated from Securigera securidaca Linn Dagen el Dorfler (N.O. Leguminosae) De, P., *Ind Jour Med Res* 1943, 31, 67.

The constituents of the seeds of Securigera securidaca Linn (Syn S. coronilla D C.). Ghosh, B.K & Dutta, A, *Ind. Jour Pharm.*, 1950, 12, 233.

SEMECARPUS ANACARDIUM Linn.

Constituents of the marking nut, Semecarpus anacardium. D. Satyanarayana Naidu; *J. Ind. Inst. Sci* 8A, 129, 1925.

Semecarpus anacardium Linn. Chemical examination of the Marking Nut. Parameswaran, P. & Salimuzzaman, Siddiqui, *J. Ind Chem. Soc.* 8 (1931), 517.

Studies in the Thermal degradation Prdts of naturally occurring resins, Pt. I. Bhilawan shell liquid. Ram Prakash Rastogi, Vishwa Nath Sharma & Salimuzzaman Siddiqui, *J. Sci. Indl. Res.*, 1948, 7B, 61.

Chemo-therapeutic studies in Bhilawan series, Pt I. Syn of sulphuric acids and arsines from Bhilawanol and its derivations Vishwa Nath Sharma, Ahmed Kamal & Salimizzaman Siddiqui, *J. Sci. Indl. Res*, 1948, **7B**, 67

S. TRAVANCORICA Bed

Chemical Examination of the constituent of Holigarna arnottiana Hook. F Pt I, latex of Semecarpus travancorica Bed Pt II Nair, A V, Poti, D N & Pillay, P P, *J Sci Indl Res.*, 1952, **11B**, 294, 298

SENECIO JACOBAEA Linn. & S RETRORSUS D C

The alkaloids of Senecio species The necines and necic acids from S. retrorsus and S. Jacobaea, Richard, H F Manske, *Can J. Res.* 5, 61, 1931.

S. VULGARIS Linn

Alkaloids of species of Senecio. Alkaloids of Senecio vulgaris. Degradation of Sinecionine R A Konovalova & A P. Onekhov, *Chem. Abst* 7436, 1937

SESELI INDICUM Wight & Arn

Essential oil of Seseli indicum W & A Farooq, M.O., Gupta, G S, *Curr. Sci.* 1953, 2, 46

SETARIA PALMAEFOLIA Stapf

Fodder value of Setaria palmaefolia Stapf Mirchandani, T.J & Dabadghao, P M, *Sci & Cult*, 1949, 14, 432

SIDA CORDIFOLIA Linn

The action of Sympathomimetic alkaloid in Sida cordifolia (Brela). Chopra, R N., Premankur, De, *Ind. Jour. Med Res*, 1930, 18, 467.

Sida cordifolia Linn Chemical examination of Sudhamoy Ghosh & Ashutosh Dutt, *J Ind Chem Soc* 7, (1930), 825.

SISYMBRIUM IRIO Linn

Chemical Examination of the fatty oil from the seeds of Sisymbrium irio Linn. Joti Sarup Aggarwal & Karimulla, *J. Sci Indl Res.*, 1946, 5, 57

SKIMMIA LAUREOLA Sieb & Zucc ex Walp

Chemistry of essential oils Investigation of the oil of Skimmia laureola Heinrich Weinhaus & Tara Chand Rajdhan, *J. Prakt. Chem.*, 147, 113, 1936

A preliminary note on the chemistry and pharmacology of the leaves of Skimmia laureola Chopra, R N, Chatterjee, R G., De, N & Ghosh, S.; *Ind Jour. Med. Res*, 1938, 26, 481.

Studies on the active principles of Skimmia laureola, Asima Chatterjee, Bhattacharya, A., *J.I.C.S* 1953, 1, 33.

SOLANUM INDICUM Linn.

Some common indigenous remedies. Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum Chopra, R N & Ghosh, S, *Ind Med Record* 55. 77, 1935

S NIGRUM Linn.

Chemical examination of the fruits of Part I The comp of the oil from the seeds Pendse, G P, *J Ind Chem Soc* 14 (1937), 366

S PSEUDOCAPSICUM Linn

The action of an alkaloidal product from the leaf of Solanum pseudocapsicum Watt, L J M, Heimann, H L & Mellzer, E; *J Pharmacol* 39, 387, 1930

Chemical Examination of the fatty oil from the seeds of Solanum pseudocapsicum Linn Pt I Husain, R, Siddiqui, S., Warsi, S A, *Ind Jour Pharm*, 1944, 6, 68.

S TORVUM Sw

A note on the chemistry of Solanum torvum Krishnamurti, G V, Sheshadri, T R., *J Sci Indl Res* 1949, **8B**, 97.

S XANTHOCARPUM Schrad & Wendl

Chemical examination of some Indian medicinal plants. Tinospora cordifolia, Solanum xanthocarpum and Fumaria officinalis Pense. G P. & Dutt, S., *Ind Jour Med. Res*, 1932, 20, 663

Fruits of Solanum xanthocarpum Sayed, I Z. & Kanga, D D, *Proc Ind. Acad Sci* **4A**, 255, 1936

The chemical examination of Solanum xanthocarpum Schard & Wendel Part I The constitution of the oil from the seeds Gupta, Mahadeo Prasad & Dutt, Sikhishushan, *J Ind Chem Soc* 13 (1936), 613

Chemical examination of the seeds of. Solanum xanthocarpum. Schard & Wendel Part II The constituents Gupta Mahadeoprasad & Sikhishushan Dutt, *J. Ind. Chem Soc.* 15 (1938), 95

SONNERATIA ACIDA Linn.

Constitution of the colouring matter from Sonneratia acida Linn Govind Rai Chaudhry, Vishwa Nath Sharma; Siddiqui, S, *J Sci Indl. Res*, **7B**, 292, 1949

Chemical Examination of Archa (Sonneratia acida Linn). Pt I. Isolation of three crystalline products from the wood Chaudhry, G R & Siddiqui S; *J. Sci. Indl. Res.*, 1950, **9B**, 137.

Chemical Examination of Archa (Sonneratia acida Linn.) Pt II. Studies in the constitution of Archin and Archiniry Chaudhry, G R., Vishwa Nath Sharma & Siddiqui, S, *J. Sci. Indl. Res.*, 1950, **9B**, 142.

SPARTIUM SCOPARIUM Linn.

Lupine alkaloids. Secondary alkaloids of *Spartium scoparium*
Winterfield, K. and Fritz Nitzsche, *Arch. Pharm.* 278, 393, 1940.

SPHAERANTHUS INDICUS Linn.

Chemical investigation of *Sphaeranthus indicus* Linn. Basu, N K. &
Lamsal, P.P. *J. Am Pharm Assoc* 35, 274, 1946.

SPILANTHES ACMELA Linn

Chemical investigation of *Spilanthes acmela* Linn. Gokhale & Bhide
J. Ind Chem Soc. 22, (1945), 250

SPIRAEA ARUNCUS Linn.

An antibiotic from *Spiraea aruncus* L E P Abraham, N G. Heatley
R. Rolt and E M Osborn, *Nature*, 157, 511, 1946.

STELLARIA MEDIA Linn.

A note on the embryology of *Stellaria media* L. Niranjan Pal
Sci. & Cult, 1950, 16, 37.

STEPHANIA GLABRA Miers

The alkaloids of Indian Stephanias Pt. I Isolation of three crystalline
alkaloids from the tubers of *Stephania glabra* Miers Chaudhry, G R
& Siddiqui S., *J. Sci Indl. Res*, 1950, 9B, 79

The alkaloid of *Stephania glabra* Miers. Identity of Aindarine &
Gindarine with Tetrahydropalmatine and Palmatine respectively
Chaudhry, G.R, Sharma, V.N. & Dhar, M L, *J Sci Indl. Res* 1952
11B, 337.

STRYCHNOS NUX-VOMICA Linn

Nux vomica, chromatographic analysis of its tincture and extract
Notes and News. *Ind Jour Pharm* 1946, 8, 105

SWERTIA CHIRATA Ham.

A modified method for the assay of *Chirata* (*Swertia chirata*) Handa
K L & Vidya Sagar Parbhakar; *Ind Jour Pharm*, 1952, 14, 87.

S. DECUSSATA Nimmo

Chemical investigation of Indian Medicinal plants Part I. Investiga-
tion of *Swertia decussata* Isolation of decussatin and Swertinin
Shah, R.C., Dalal, S R, (Miss) Sethna, S, *J I C S* 1953, 7, 457.

Chemical investigation of Indian Medicinal plants Part II
Structures of decussatin and Swertinin. Shah, R.C, Dalal, S R., (Miss)
Sethna, S, *J I C S*, 1953, 7, 463.

SWIETENIA MACROPHYLLA Linn.

Chemical investigation of the seeds of *Swietenia macrophylla* Linn
Chakravarty, T. & Guha, S.S., *Sci. & Cultr*, 1947, 12, 450

Swietenia macrophylla. Chemical investigation of the non bitter
principle of. Guha, Sircar & Chakravarty, *J. Ind Chem. Soc*
28 (1951), 207.

Swietenin from the seed of *Swietenia macrophylla*. Sircar, Guha, & Chakravarty, J., *Ind. Chem. Soc.* 28 (1951), 208.

SYMPLOCOS RACEMOSA Roxb.

Pharmacognostic studies of *Symplocos racemosa* Roxb. Baddomu, S, Clarke, C B, Chatterjee, K C. & Khorana, M L., *Ind Jour. Pharm.*, 1950, 12, 290.

TABERNAEMONTYNA CORONARIA Br.

Phytochemistry of the bark of *Tabernaemontana coronaria* Br Ratangiriswarn, A.N. and Vankatachalum, K, *Quart J. Pharm. Pharmacol* 22, 174, 1939.

TAGETES ERECTA Linn.

Petaloidy of sepal in a ray floret of *Tagetes erecta*. Farooq, M, *Sci & Cult*, 1951, 17, 92.

T PATULA Linn

Colouring matter of the flowers of *Tagetes patula* Isolation of a new flavonol, patulatin and its constitution. P Suryaprakasa Rao & Seshadri, T.R., *Proc Ind. Acad Sci.* 14A, 643, 1941

TAMARINDUS INDICA Linn.

Tartaric acid from Tamarind fruit Iyenger, A V, Varadaraja, *Curr Sci*, 1937-38, 6, 610

Tamarind seed 'Pectin' Hanji, H R, Savur, G R. & Sreenivasan, A., *Curr Sci* 1945, 14, 129.

Tamarind seed 'Pectin' Damodran, M & Rangachari, P N., *Curr. Sci*, 1945, 14, 203.

Tamarind seed 'Pectin'. Ghose, T P. & Krishna, S, *Curr. Sci.*, 1945, 14, 299.

Tamarind seed 'Pectin', Ghose, T P., Krishna, S & Suryaprakasa Rao, P, *J Sci Indl. Res.*, 1945-46, 4, 705.

Tamarind seed 'Pectin', Damodran, M & Rangachari, P N, *Curr. Sci*, 1946, 15, 20 & 133

Tamarind seed 'Pectin', Rao, P S & Krishna, S, *Curr. Sci.*, 1946, 15, 133 & 163.

Tamarind seed 'Pectin', Savur, G R. & Sreenivasan, A., *Curr. Sci.*, 1946, 15, 43, 134 & 168.

Tamarind seed Polysaccharides Rao, P S, Krishna, S, *Curr. Sci.*, 1947, 16, 256.

Tamarind seed Jellose ("Pectin"), its jellying properties. Rao, P S., *J. Sci Indl Res.*, 1948, 7B, 89.

Tamarind (*Tamarindus indica*) seeds as Protein rich feed for Live-stock, Kehar, N.D. & Sahai, B., *Sci. & Cult.*, 1949, 14, 534.

Jellies and related products from Tamarind seed kernels. Rao, P.S., *J. Sci. Indl. Res* , 1949, **8A**, 354.

Studies on Tamarind seed kernel powder, Pt I. Preparation and utilization as a sizing material in the Jute Industry. Macmillan, W.G. & Chakarvarti, I.B , *J Sci Indl Res* , 1951, **10B**, 13

Studies on Tamarind kernel powder Pt II Physical Prop T K Paste & T K P Sind Yarns Macmillan, W G & Chakarvarti, I B , *J Sci Indl Res* , 1951, **10B**. 270

Studies on Tamarind kernel powder. Pt III Comparative study of antiseptics used in Jute sizing Mixtures Macmillan, W G. Chakarvarti, I B & Pal, P N , *J Sci Indl Res* , 1952, **11B**, 438

TANACETUM. Linn

Oleum Tenaceti, Caius, J.F & Mahaskar, K S , *Ind Jour Med Res*. 1920, 7, 606

TAXUS BACCATA Linn

Physiologically active constituents of the Yew, *Taxus baccata*, Taxine Robert K Callow, John M Gulland & Cyril J Virden; *J Chem Soc* 2138, 1931.

Detection and estimation of the alkaloid Taxine in preparations of *Taxus baccata*, Kuhn, A & Schafer, G , *Chem. Abst* 723, 1938

T. BREVIFOLIA Nutt.

Taxus brevifolia Georges Masson, *J Am. Pharm Assoc.* 28, 493, 1939.

TECTONA GRANDIS Linn.

The oil from the seeds of *Tectona grandis* (Teak) Puntambekar, S V & Krishna, S , *J. Ind Chem Soc.* 10 (1933), 401

TEPHROSIA Pers

Source of Indian *Tephrosia* sp. as rotenone Krishna, S & Ghose. T P , *Curr Sci* , 1937-38, 6, 454

Retenone containing *Tephrosia* as a source of insecticides. M G. Timoshenko , *Chem Abst.* 3530, 1943.

Effect of extracting agents & methods of extraction on the toxicity of *Tephrosia* extracts. Z K. Bogatova, *Chem. Abst.*

T. PURPUREA Pers

Studies on the hypoglycaemic effect of *Tephrosia purpurea* Var pumila Swift, H B N , Sethi, M S , Sareen, K N , *Ind. Med. Gaz* , 1949, 84, 243

T. VILLOSA Pers.

Tephrosia villosa. A plant useful in diabetes Harbhajan Singh; *Ind. Jour. Pharm.*, 1945, 7, 60

TERMINALIA ARJUNA W. & A.

Terminalia arjuna. Its chemistry, pharmacology and therapeutic action. Chopra, R N & Ghosh, S, *Ind Med Gaz.*, 1929, 64, 70

Chemical Examination of *Terminalia arjuna*. Isolation of arjunin. Radharaman Aggarwal & Sikhishushan Dutt, *Proc Acad. Sci. Ind.* 5, 50, 1935

Chemical examination of the bark of *Terminalia arjuna* Isolation of arjunetin from the alcoholic extract Radharaman Aggarwal & Sikhishushan Dutt, *Proc Natl Acad. Sci* 6, 305, 1936

T. BELERICA Roxb.

Preliminary examination of the Fruits of *Terminalia belerica* Roxb Chakravarti, M D & Tayal, J N, *Sci & Cultr*, 1947, 13, 122

T CHEBULA Retz

Distribution of Tannin in different parts of the Myrobalan plant (*Terminalia chebula*) Biswas, H G, *Sci & Cult*, 1944, 9, 399.

Tannic acid B P from Myrobalan (*Terminalia chebula*) Biswas, N G, *J Sci Indl. Res*, 1947, 6B, 122

THALICTRUM FOLIOLOSUM D.C

Chemical examination of *Thalictrum foliolosum* D.C Isolation & characterisation of a new alkaloid Thalictin Vashistha, S K & Siddiqui, S., *J Ind Chem Soc* 18 (1941) 641

Plant alkaloids Part III On *Thalictrum foliolosum* Chatterjee R, Guha, M D & Chatterjee, A, *J Ind Chem Soc* 29 (1952) 371

THEA SINENSIS Linn

The use of tea in the treatment of burns Peiris, M V P, *Ind Med. Gaz*, 1937, 72, 718.

THEOBROMA CACAO Linn

Coca leaves, the determination of ether soluble alkaloids and ecognineesters Notes and News, *Ind Jour Pharm* 1946, 8, 104

THEVETIA NEREIFOLIA Juss.

The Pharmacological action of *Thevetia* glucoside occurring in *Thevetia nereifolia* (Yellow oleander) Chopra, R N & Mukerji, B, *Ind Jour Med Res*, 1933, 20, 903

Chemical examination of the roots of *Thevetia nereifolia* (Juss) N. Ghatak and G P Pendse, *Bull Acad Sci. Ind* 2, 259, 1933

The constituents of be-still nuts, *Thevetia nereifolia* K K Chen. & A. Ling Chen, *J Biol. Chem* 105, 231, 1934

The pharmacological action of Thevetoxin. A second glucoside from *Thevetia nereifolia*. Bhatia, B B. & Lal, S, *Ind. Jour. Med. Res.*, 1934, 21, 605.

Nevefolin, a new digitalic heleroside from *Thevetia nereifolia* Marcel Frerrefacque, *Compt. Rend.* 221, 645, 1945.

THYMUS SERPHYLLUM Linn.

Studies in Indian Essential oils. III—Essential oil from leaves of *Thymus serpyllum*, Linn. Jagjit Singh and Rao B. Sanjiva *Journal of the Indian Institute of Science*, Bangalore, 1932, Vol 15A Part VII, pp 78-83.

THYSANOLAENA MAXIMA Kuntze

Thysanolaena maxima as cattle feed Das, B K & Mukherjee, N.C. *Sci & Cult*, 1951, 17, 137

THYMUS VULGARIS Linn

Thymus vulgaris L Christine Rosenthal, *Pharm Ind.* 10, 22, 1943.

TINOSPORA CORDIFOLIA Miers.

Chemical Examination of *Tinospora cordifolia* Miers Kidwai, A R, Salooja, K C, Vishwa Nath Sharma & Siddiqui, S, *J Sci. Indl. Res.*, 1949, 8B, 115

Chemical examination of some Indian medicinal plants *Tinospora cordifolia*, *Solanum xanthocarpum* and *Fumaria officinalis*. Pense, G.P & Dutt, S., *Ind Jour Med Res*, 1932, 20, 663.

The nature and germination of seeds of *Tinospora cordifolia* Miers Ajrekar, S L. & Oza, J D, *Curr Sci*, 1934-35, 3, 379.

Chemical investigation of *Tinospora cordifolia* Bhide, B V., Phalankar, N.L. and Pranjpe, K, *J. Univ Bombay.*, 3, 89, 1941

Chemical investigation of *Tinospora cordifolia* (Miers) Bhide, B.V, Phalankar, N L and Pranjpe, K, *J Univ. Bombay.* 10, 89, 1941

T. CRESPIA Miers

The wax of *Tinospora crespia* Miers. Lucienene Beauquesne; *Chem. Abst* 868, 1942

TODDALIA ACULEATA Pers

On the failure of *Toddalia aculeata* in the treatment of malaria. Bhatia, B B, *Ind Med Gaz*, 1932, 67, 192

The chemistry and pharmacological action of *Toddalia aculeata*. Dey, B B., Pillay, P P, David, J C. & Rajamanikam, N., *Ind. Jour. Med. Res*, 1935, 22, 765

Botany, Chemistry and Pharmacodynamics of *Toddalia aculeata*. Lobstien, J.E and Hesse, P.; *Chem Abst.* 2811, 1937.

On the constitution of natural coumarins of *Toddalia aculeata*. Phanibhushan Dutta, *Jour Ind. Chem. Soc.*

TRAPA BISPINOSA Roxb.

The food value of the nut of *Trapa bispinosa*. Brahmachari, B.B. & Chatterjee, N K., *Ind. Med. Gaz.*, 1927, 62, 365.

TRESPASIA POPULNEA

Colouring matter of Indian Tulip (*Trespasia populnea*) flowers. Populnin and populnetin. Oelakatham, K & Seshadri, T R, *Curr. Sci.*, 1938, 7, 16

TRIANTHEMA PORTULACASTRUM Linn.

A comparative study of *Boerhaavia diffusa* Linn. and the white and red flowered varieties of *Trianthema portulacastrum* Linn. Chopra, R N, Chatterjee, N R and Ghosh, S, *Ind Jour Med Res*, 1940, 28, 475

TRIBULUS TERRESTRIS Linn

Chemical Examination of the fruits of *Tribulus terrestris* Linn. Ghatak, N, *Bull Acad Sci* 2, 163, 1933.

TRICHOSANTHES ANGUINA Linn

Chemical examination of drying oil from the seeds of *Trichosanthes anguina* Linn Miss Padmini Soni, Aggarwal, J S., *J Sci Indl Res.*, 1949, 8B, 150

T DIOICA Roxb

Chemical examination of the Indian Medicinal plants, *Trichosanthes dioica* Nag, N C, *Trans Bose Research Inst, Calcutta*, 10, 113, 1934-35.

TRIGONELLA FOENUM-GRÆCUM Linn

Composition of Fenugreek seeds Mixture of fenugreek seeds with grain intended for flour E Fleurent, *Compt. Rend* 182, 944, 1926

Proteins of Indian Food Stuffs V Alcohol-soluble protein of fenugreek (*Trigonella foenum-graecum*) Sreenivasa Rao, Y V Sastri, B N and Narayana, N, *J Ind Inst Sci.* 16A, 85, 1933

Thesaponin of fenugreek seeds. Gabra Soliman and Zahira Mustafa, *Nature*, 151, 195, 1943

Study and improvement of fenugreek (*Trigonella foenum graecum* L.). Toao Marquis De Almeida, *Chem Abst* 419, 1943

Sapogenins The sapogenins of *Trigonella foenumgræcum* Russel E Marker, Wagner, R B, Ulshaper and Clarence H Rouf, *J Am. Chem. Soc.* 65, 1247, 1943.

TRITICUM Linn

The hydrolysis of ptytic compounds derived from seeds of hemp, horse bean, horse chestnut, flax, wheat and embryos of rye. W. Jarosza, *Chem. Abst* 5501, 1934

Poisonous food grain wheat mixed with *Lolium temulentum*. Greva, S.D S & Bhandari, P.N, *Ind. Med Gaz*, 1946 81, 294

TRIUMFETTA RHOMBOIDEA Jacq.

The development of the female gametophyte in *Triumfetta rhomboidea* Jacq Bhattacharyya, S.S. & Mitra, J.N., *Sci. & Cult.*, 1952, 137, 45.

TUSSILAGO FARFARA Linn

Plant pigments Taraxanthin from *Tussilago farfara* (Colts foot)
Karner, P and Morf, R., *Helv. Chem Acta* 15, 863, 1932.

TYLOPHORA ASTHMATICA W & A

The chemical examination of *Tylophora asthmatica* and isolation of the alkaloid Tylophorine and Tylophorinine Ratnagiriswaran, A N & Venkatachalam, K, *Ind Jour Med Res*, 1934, 22, 433

Chemical and Pharmacological investigation of *Tylophora asthmatica* Chopra, R N, Ghosh, N N, Bose, J B & Ghosh, S, *Arch Pharm* 275, 236, 1937

URGINEA INDICA Kunth

Chemical Examination of Indian squill Seshadri, T R & Subramanian, S S, *J Sci Indl Res*, 1950, 9B, 114

Optical activity of preparations of Indian Squill Subramanian, S, *Ind Jour Pharm*, 1952, 14, 25

VACCINIUM Linn.

The constituents of diuretic drugs, the flavonol glucoside of *Folium Uvae Ursi* and *Folium Vaccinii-Harnkitu* Nakamura, Tatuo Ota and Genitiro Hukuti, *J. Pharm Soc, Japan* 55, 800, 1935.

VALERIAN Linn

Investigation of the Pharmacological action of fresh and dried Valerian Janina Romanowska Majcherezyk, *Chem Zent* 906, 1939

V OFFICINALIS Linn

An acid ester present in the roots of the *Valeriana officinalis*, Emil Coringa, *Comp Rend* 201, 1152

Pharmacological investigation of four strains of *Valeriana officinalis* var *Labfolia* W Rusieki, *Chem Zent* 1404, 1939

VANDA ROXBURGHII R. Br

Pharmacological action of an active constituent isolated from *Vanda roxburghii* R Br Gupta, J P, Roy, P. K & Sen Gupta, K K, *Ind Jour Med Res*, 1946, 34, 253.

VANGUERIA SPINOSA Roxb

The fat from the seeds of *Vangueria spinosa* (N O. Rubiaceae). Airan, S. W & Shah, S V, *Curr Sci*, 1942, 400, 11

Fat from the seeds of *Vangueria spinosa*. Nadkarni, Airan & Shah, *J Ind Chem. Soc.* 24(1947), 25-30

VERBENA OFFICINALIS Linn.

Constituents of *Verbena officinalis* L Identity of Verbenalin with corin. Benno Reichert *Arch. Pharm.* 273, 357, 1935.

V. VENOSA.

Presence of Stychose in the stems and roots of *Verbena officinalis*, and the underground parts of *Verbena venosa*. J. Cheymol *J. Pharm. Chem.* 25, 110, 1937.

Dessication of *Verbena officinalis* L. Loss of holosides and Verbenalloside J Cheymol, *J. Pharm. Chem.* 25, 581, 1937.

Constituents of *Verbena officinalis* L., constitution of Corn. Benno Richert and Walter Hoffmann, *Arch. Pharm.* 275, 474, 1937

Action of Verbenalloside on two isolated animal organs. J. Cheymol; *J. Pharm. Chem.* 27, 386, 1938

Glucide (total holoside and verbenalloside) contents of different parts of *Verbena officinalis* Jean Cheymol, *Chem Abst* 2977, 1938.

Verbanalin. E. Buris & D. Susterova. Rihova, *Chem Abst.* 6004, 1938.

Chemical and Pharmacological Researches on verbenin, a glucagoc-acting glucocide from *Verbena officinalis* L, Kotoku Kawazima, *Chem Abst* 7396, 1939

Verbenalin (*Verbena officinalis*). P. Karner & H. Salomon, *Helv Chim Acta* 26, 1544, 1946

VERNONIA ANTHELMINTICA Willd.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency. Sentonin, *Oleum rutae* (*Ruta graveolens* Linn.) *Butea monosperma* Roxb., *Melia azadirachta* Linn., *Punica granatum* Linn., *Picrasma excelsa* Swartz, *Vernonia anthelmintica* Willd., *Cocos nucifera* Linn (coconut). Caius, J.F & Mhaskar, K.S, *Ind Jour Med Res.*, 1923, 11, 353.

Anthelmintic properties of *Vernonia anthelmintica* Willd (Syn *Serrantula anthelmintica*) Chopra, R N., Ghosh, N N & Mukerji, A K., *Ind. Jour Med Res.*, 1934, 22, 183.

VETIVERIA ZIZANIODES Stapf

Notes on some Indian Essential Oils "Vetiveria Zizaninoides" Stapf Rao B. Sanjiva, Sudborough, J J & Watson, H F, *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. 8A, Part X, pp. 147-149

Analytical characteristics of Dutch East Indian oils of *Cananga patchouli* and *Vetiver* Koolhas, D R. & Rowan, P.A, *Perfumes France*, 15, 245, 1937.

Cananga patchouli and *Vetiver* Oil. Koolhas, D.R. & Rowan, P.A, *Perfumery essential oil Record* 29, 53, 1938.

Indian *Vetiver*, Khus. Menon, A K, Ittyachan, C.T., *J.Sci. Indl. Res*, 1947, 6, 274.

VIGNA CATJANG Walp.

Studies in nutrition value of Indian vegetable foodstuff. Pt. III. Nutritive values of *Lentil*—*Lens esculenta*; Moench, Cow pea—*Vigna catjang*, Walp and *Aconite Bean*—*Phaseolus aconitifolius*,

Jacq. Niyogi, S.P., Narayana, N. & Desai, B.G , *Ind. Jour. Med. Res.*, 1932, 19, 859,

VINCA ROSEA Linn.

Abnormalities in the flowers of *Vinca rosea* Linn. Kapadia, G A , *Sci. & Cult* , 1950, 15, 402

VIOLA ODORATA Linn.

Viola odorata L. the fragrant violet considered from the old and the new point of view Ludwig Kroeber. *Chem. Abst.* 6022, 1947

VISCUM Linn

Cassytha L. on *Viscum*. Dutt, B M S , *Sci. & Cult* , 1950 16, 258

V. ALBUM Linn.

Active principle of *Viscum album* A Enders, O Feuchtinger & S Jannssin , *Arch. Expl Path. Pharmacol* 196, 290, 1940.

Pharmacology of *Viscum album*. Stamatis C Conninos, *Chem Abst* 5996, 1941

The pharmacologically active substances in *Viscum album*. Alfred A. Andres, *Chem Abst* 5996, 1941.

VITEX NEGUNDO Linn.

An introductory note on the chemical investigation of the plant *Vitex negundo* (N O Verbenaceae) Basu, N K , Singh, G B , *Sci & Cultr.*, 1944, 9, 508

A note on the chemical investigation of *Vitex negundo* Pt I Basu, N K & Singh, G B , *Ind Jour. Pharm* , 1944, 6, 71

Investigation of Indian medicinal plants, *Hydrocotyle asiatica* *Vitex negundo* and *Monniera cuneifolia*, Basu, N.K , Iamsal, P P & Singh, G B , *Quart J. Phar. Pharmacol* 20, 135, 1947

V PEDUNCULARIS Wall.

On the failure of *Vitex peduncularis* in the treatment of malaria Chopra, R N., Knowles, R & Gupta, J.C , *Ind. Med. Gaz.*, 1924, 59, 133.

Vitex peduncularis—an antihaemolytic agent Gupta, J.C , Kahali, B.S. & Ganguly, S.C. , *Trop. Diseases Bull* 40, 677, 1943.

WRIGHTIA TINCTORIA Ber.

Chemical examination of the fixed oil from the seeds of *Wrightia tinctoria*. Parihar & Dutt, *J. Ind. Chem Soc* 23 (1946), 307.

XANTHIUM STRUMARIUM Linn.

Oil from the seed of Gokbru (*Xanthium strumarium*), Shrivastava, R.C., Krishnamurthy, R.S. & Athawale, C R , *J Soc. Indl. Res.*, 1950, 9B, 282.

ZANTHOXYLUM ACANTHOPODIUM D C. & Z. ALATUM Roxb.

The constituents of some Essential Oils. 42 (a), Part V—The Essential Oil from the seeds of *Zanthoxylum alatum*, Roxb. 42 (b). Part VI—The Essential oil from the seeds of *Zanthoxylum acanthopodium* D C 42(c). Part VII —The Essential Oil from the seeds of *Zanthoxylum budrunga*, Wall. Simonsen, J L. & Rau M. Gopal; *Indian Forest Records*, 1922, Vol IX Part IV. pp. 23-26

Z ALATUM Roxb

Zanthoxylum alatum ; —*Proc. Nat. Acad Sc.* Vol. IX, 1939, pp 187-192

Z BUDRUNGA Wall.

The constituents of some Essential Oils. 42(a) Part V—The Essential Oil from the seeds of *Zanthoxylum alatum*, Roxb 42 (b). Part VI—The Essential Oil from the seeds of *Zanthoxylum acanthopodium*, D C. 42(c). Part VII The Essential Oil from the seeds of *Zanthoxylum budrunga*, Wall Simonsen, J L & Rau M Gopal, *Indian Forest Records*, 1922, Vol. IX Part IV, pp. 23-26.

The alkaloids of *Zanthoxylum budrunga* Khastagir, Harinarayan, *Curr Sci* 1947, 16, 185.

A study of the colouring matter of Tambul seeds. Seshadri, T.R , *J. Sci Indl., Res.*, 1949, 8A, 232.

Z OVALTIFOLIUM Wight.

Constituents of some Indian essential oils Essential oil from the seeds of *Zanthoxylum ovaltifolium*, Simonsen, J L, *Ind Forest Records*, 11, 1, 1924.

Constituents of some Indian essential oils from the seeds of *Zanthoxylum ovaltifolium*, *Juniperus communis* Simonsen, J L ; *Ind. Forest Records*, 11, 1, 1924.

The Constituents of some Indian Essential Oils Part XIV—The Essential Oil from the seeds of *Zanthoxylum ovaltifolium* Simonsen, J L , *Indian Forest Records*, 1925, Vol. XI, Part I, pp. 1-5.

Z RHETSEA D.C.

Notes on some Indian Essential oils. *Zanthoxylum rhetsea* D C. Rao, B Sanjiva, Sudborough, J.J. & Watson, H E , *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol 8A, Part X, pp. 174-177

ZINGIBER OFFICINALE Roscoe.

Notes on some Indian Essential Oils *Zingiber officinale*. Roscoe, Rao, B Sanjiva, Sudborough, J J & Watson, H.E., *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. 8A, Part X, pp. 151-153.

Travancore Essential Oils. Part VII—The Essential Oil from the Rhizomes of Ginger, *Zingiber officinale* Moudgill, K.L ; *Journal of the Indian Chemical Society, Calcutta*, 1928, Vol V.

Essential oils of Travancore, Part VII, from rhizome of Ginger, *Zingiber officinale*. Kishori Lal Moudgill, *J. Ind. Chem Soc.* 5 (1928), 251.

A note on the essential oil from Ginger scrapings. Varier, N S, *Curr. Sci*, 1945, 14, 321.

ZIZYPHUS VULGARIS Lam.

Anaesthetics from leaves of *Zizyphus vulgaris* Lam E N. Taran, *Farmatsiya*, 4, 20, 1941.

Z XYLOPYRUS Willd.

Oil from the seeds of *Zizyphus xylopyrus* Willd Airan, J W., *Curr. Sci* 1948, 17, 150

GENERAL

ALKALOIDS.

Estimation of total alkaloid by Chromatography. Datal, V. D. & Khorana, M. L., *Ind. Jour Pharm.* 1950, 12, 173

The formation of alkaloids in plants Chatterjee, R *Sci & Cult*-, 1950, 16, 58.

ALKALOID CHEMISTRY.

Trends in alkaloid chemistry. Chatterjee, R. G., *Sci. & Cult*, 1951, 17, 237.

ANIMALS.

CONORHINUS RUBROFASCIATUS De Geer

Some notes on *Conorhinus rubrofasciatus* (De Geer). Awati, P R. *Ind Jour. Med. Res.* 1921, 9, 371.

LABEO ROHITA.

Bile from *Labeo rohita* Basu, U. P., *Ind. Jour. Pharm.*, 1944, 6, 81

MYLABRIS

Synthesis of Cantharidin. Paranjape, K. (Miss), Phalnikar, N L, Nargund, K. S., *Curr. Sci.*, 1943, 12, 256.

M MACILENTA.

Cantharidin content of *Mylabris macilenta* Beetles. Mukerji, B, Hassain, T. & Karim, B. A., *Curr. Sci*, 1944, 13, 315.

UROLONCHA MALABARICA Linn.

On a Trypanosome of the white throated Munia—*Uroloncha malabarica* Linn Das Gupta, D. M. & Siddons, L B., *Ind Med. Gaz*, 1941, 76, 151.

ANTHELMINTICS.

Correlation between the Chemical composition of Anthelmintics and therapeutic values in connection with the Hookworm enquiry in the Madras Presidency. XXII. Summary and conclusions Caius, J F & Mhaskar, K. S., *Ind. Jour. Med. Res.*, 1923, 11, 371.

ANTIBACTERIAL SUBSTANCES.

Antibacterial substance in plants. Notes & News., *Ind. Jour. Pharm.* 1946, 8, 60.

ANTIBIOTICS.

A search for Antibiotic substances in some Indian Medicinal Plants. Mariam George, Venkataraman, P R & Pandalai, K M , *J Sci. Indl. Res.* 1947, **6B**, 42.

Investigation on plant antibiotics. Pt. IV Further searches for antibiotic substances in Indian medicinal plants George, Mariam & Pandalai, K M , *Ind Jour Med Res.*, 1949, 37, 169

Lichens and Antibiotic Activity Rangaswami, S & Subramanian, S. S., *Ind. Jour. Pharm* , 1952, 14, 214

Antibiotic activity of some Indian Medicinal plants Joshi, C G. & Magar, N. G , *J Sci. Indl Res* , 1952, **11B**, 261

AROMATIC PLANTS.

Supplement. Aromatic Plants of India. Pt IX, X, XI, XII Krishna, S , Badhwar, R. L , **2A, 12A, 10B, 11B.** *J Sci Ind Res.*

Supplement Aromatic Plants of India Pt XV *J Sci Ind. Res*

Aromatic Plants of India XIII Krishna, S & Badhwar, R L , *J. Sci. Ind. Res*

Supplement : Aromatic Plants of India Krishna, S & Badhwar, R. L., *J. Sci. Ind Res.*,

Aromatic supplement, plants of India, I, II, III, IV Krishna, S & Badhwar, R. L., *J. Sci Indl. Res*

CRUDE DRUGS

Report on (the assay of) crude drugs H. W Youngken, *J Assoc Official Agr Chem.* 18, 516, 1935

The present position of the crude drugs used in the Indigenous medicine. Pt. I Handa, K L , Kapoor, L D , Chopra, I C & Som Nath; *Ind. Jour. Pharm.*, 1951, 13, 29

CULTIVATION OF MEDICINAL PLANTS

Cultivation of Medicinal Plants in India Bal, S N , *Ind. Jour. Pharm.*, 1944, 6, 42

Medicinal Plants and their cultivation. Mitra, G C , *J. Sci. Indl. Res.*, 1948, **7A**, 319.

Scope for the cultivation of medicinal plants in India. Chopra, R N , Kapoor, L. D , Handa, K. L., Chopra, I C. & Nayar, S L , *J. Sci, Indl Res.*, 1948, **7A**, 527.

Cultivation of Medicinal plants in Kashmir Chopra, I C., Kapoor, L. D , Handa, K. L., *J. S. I. R* 1953, **7A**, 31.

DEHYDROASCORBIC ACID

Significance of dehydroascorbic acid in plants. Inderjit Babbar, *Ind. Jour. Med. Res* , 1950, 38, 263.

DRUG ADULTERATION

Drug adulteration and spurious drugs in India. Chopra, R. N., *Ind. Med. Gaz.*, 1935, 70, 693.

DRUG RESOURCES

Drug Resources of Himachal Pradesh. Handa, K L., Kapoor, L.D., & Chopra I C, *Ind. Jour Pharm.*, 1951, 13, 118

Drug Resources of Kangra Valley Handa, K L., Kapoor, L.D. & Chopra, I C, *J Sci Indl Res* 1951, 10B, 173.

DRUG SCARCITY.

Scarcity of drugs Mohammad Hamid, *Ind Jour Pharm* 1944, 6, 46.

ESSENTIAL OILS.

Indian Essential oils—Sudborough, J.J., *Journal of the Indian Institute of Science, Bangalore*, Vol. I & II. Part II. pp 13 to 27, 1913.

A further note on the efficacy of essential oils in the prevention and treatment of cholera Tomb, J W, *Ind Med Gaz* 1924, 59, 233.

Notes on Indian essential oils. Rao, B S, Sudborough, J J. & Waston, H E, *J Ind Inst Sci* 8A, 143, 1925

The constituents of some Indian Essential oils Note on the rate of oxidation of d-3 Carene and other Terpenes in the presence of Catalysts Rao, M. Gopal, *Indian Forest Records*, 1925 (Chemical Series), Vol XI, Part V

Observations on the treatment of Cholera with essential oil, mistura pro-diarrhoea and permanganate of potash. Bharti, S R.; *Ind. Med Gaz*, 1926, 61, 596.

The constituents of Indian essential oils. B. Sanjiva Rao & John Lionel Simonsen, *J. Chem Soc* 2496, 1928

Notes on some Essential oils. *Perf. & Ess Oil Rec.* 1937, p. 28.

Notes on some Essential oils—*Perf. & Ess Oil Rec.* Nov. 1937, p 411.

Scent Factor of Flowers—Rakshit, J.N. *The Perfumery & Essential Oil Record*. July 20, 1937, Vol 28, No. 7, page 241.

Essential oils from Flowers, Grasses and Plants—Rakshit, J.N.; *Rakshit Gardens, Bulletin No. 2, Ghazipur, U.P* 1938.

Development of Essential Oil industry—Rakshit, J.N.; *Science & Culture*, Vol V, No 2, August 1939.

Can Indian Natural Essential Oils compete with Synthetic Products of foreign countries—Rakshit, J.N.; *Science & Culture*, Vol. 5, No 6, December 1939, pp. 355-357.

Exploitation of Phytosynthesis of Essential oils—Rakshit, J.N.; Applied Chemistry Students' Reunion, 1940, published from the University College of Science and Technology, Calcutta p. 13-18.

Position of essential oil & Perfumery Trade in India. Rao, P A., *Sci. & Cultr*, 1941, 6, 710.

A preliminary study of the rideal walker-coefficient values of certain indigenous essential oils. Gupta, J C, Bose, B.C., Ganguly, S C, *Ind Med. Gaz.*, 1942, 77, 210.

ESTUARIAL FLORA

Estuarial flora of the Godavari, a preliminary Note on Venkateswarlu, V, *Sci. & Cultr.*, 1943, 8, 351.

A note on the Estuarine Flora of Kistna. Venkateswarlu, V, *Sci. & Cultr.*, 1946, 12, 295.

HEART POISONS

Heart poisons from plants Chakravarti, J.K., *Sci & Cultr* 1952, 17, 282.

INDIAN DRUG INDUSTRY.

Indian Drug Industry and the medical profession Mukerji, B, *Ind. Jour Pharm*, 1949, 11, 120.

INDIAN FOODS

Nicotinic acid contents of Indian foods Saharia, G.S, *Univ Bombay*, 13, 5, 1945.

INDIAN MEDICINE.

A note on Indian Medicine. Clifford, A, *Ind Med Gaz*, 1928, 63, 53.

INDIAN PHARMACOPOEIA.

Indian Pharmacopoeia, Pt. A.; *Indian Jour Pharm.* 1950, 12, 85

INDIAN PHARMACOPOEIAL LIST.

Note on the Indian Pharmacopoeial List Subramanian, S.S., *Ind. Jour. Pharm.* 1948, 10, 4

I.P.L. Banerjee, J.N. Tamhene, R.G & Kulkarni, V P, *Ind Jour. Pharm.*, 1950, 12, 12.

INDIGENOUS DRUGS

The field of research in Indian Indigenous Drugs Chopra, R N & Ghosh, B.N., *Ind Med. Gaz.* 1923, 58, 99

Indigenous substitute for imported drugs Bal, S N & Prasad, S, *Ind Jour Pharm.* 5, 30, 1943

Identification of Indigenous Drugs Sircar, N N, *Ind. Jour Pharm*, 1944, 6, 9.

Symposium on utilization of indigenous drugs of India. Ghosh, S, *Ind. Jour. Pharm.*, 1944, 6, 10

Action of some indigenous drugs on Uterus. A preliminary Note. Kapur, R.D., *Ind. Jour. Med. Res*, 1948, 36, 47

Symposium on utilization of Indigenous drugs. Salimuzzaman Siddiqui; *Ind. Jour Pharm.*, 1950, 12, 221.

Phagocytic response in Guinea pigs to certain indigenous Medicinal extracts. Broker, R., Gaffar, K T., *Curr Sci* 1953, 2, 44.

INDIGENOUS REMEDIES.

Chemistry and pharmacology of some common indigenous remedies. Chopra, I C., Kohli, J.D. & Handa, K.L *Ind Jour. Med. Res.*, 1945, 33, 157

INSECTICIDES OF VEGETABLE ORIGIN.

Insecticidal Plants. Kabir, S A & Ramaswamy, M N., *Curr. Sci* , 1939, 8, 82.

Insecticides of vegetable origin. Siddiqui, S , *J Sci. Indl. Res* , 1950, 9A, 181.

MEDICINAL PLANTS.

Some medicinal plants growing in Himalayas, II Chopra, R N , Ghosh, N.N., Ratnagriswaran, A N , *Ind Jour Med. Res.* 1928, 16, 777.

The scientific and economic importance of researches on Indian Medicinal Plants Ghosh, S , *Ind. Med Gaz.* 63, 650, 1928.

Observations in certain medicinal plants used in indigenous medicine Chopra, R.N & Ghosh, S., *Ind Jour. Med Res* 1929, 17, 377

Medicinal plants ; from Bt Col R N. Chopra's presidential address, National Institute of Science in India at Madras *Sci & Cultr* , 1940, 5, 620.

Research on medicinal and poisonous plants. (Article). *Curr. Sci* , 1940, 9, 94

Medicinal plants in Chitral state. Datta, S C., *Sci. & Cultr.*, 1945, 10, 519

Scope for the cultivation of medicinal plants in India Chopra, R N , Kapoor, L D , Handa, K.L , Chopra, I.C. & Nayar, S.L., *J Sci Indl. Res* , 1949, 8A, 14.

NOMENCLATURE OF DRUGS.

The nomenclature of Drugs. (An Article). *Ind. Jour. Pharm.* 1952, 14, 153.

PECTIN FROM INDIAN PLANT MATERIAL

Pectin from Indian Plant materials. Damodaran, M. & Rangachari, P.N.; *J Sci Indl Res* 1945-46, 4, 298.

PLANT PIGMENTS.

Plant pigments. Datta, C., *Sci & Cult* , 1951, 16, 458 & 512.

POISONOUS PLANTS

The toxicity of known and unknown poisonous plants in the union of South Africa Douw G. Steyn, Onderstepoort ; *J. Vet. Sci Animal. Ind* 9, 573, 1937.

Poisonous plants of India Chopra, R N., *Sci & Cultr* 1939, 5, 9

Research on medicinal and poisonous plants (Article), *Curr Sci.* 1940, 9, 94

SILAJIT

Chemical composition and antidiabetic properties of Silajit, Chopra, R N , Bose, J P & Ghosh, N N , *Ind Jour Med Res* 1926, 14, 145

STERIOD HORMONES

Isolation of steriod hormones from Indian plants Chakravarti, *Ind Jour Pharm* , 1952, 14, 80.

VEGETABLE DRUGS

Vegetable drugs Editor, *Ind Jour Pharm.*, 1951, 13, 81.

VEGETABLE DRUG RESOURCES

Vegetable drug resources of Jammu & Kashmir Pt I Plants yielding B P drug and their substitutes Chopra, R N , Kapoor, L D , Handa, K L. & Chopra, I.C., *J Sci Ind Res* , 1947, 6A, 485.

Supplement to the Bibliography.

ACORUS CALAMUS.

Pharmacological action of some common essential oil bearing plants used in indigenous Medicine Part I Pharmacological action of *Acorus calamus*, *Curcuma zedaria*, *Xanthoxylum alatum* and *Angelica archangelica* Chopra, I C , Jamwal, K S. & Khajuria, B N , *Ind Jour Med Res* , 1954, 42, 381

ADHATODA VASICA.

Antitubercular action of *Adhatoda vasica* N O Acanthacea, Part I. Gupta, K.C , & Chopra, I C , *Ind Jour Med Res* , 1954, 42, 355.

ALPINA GALANGA.

Pharmacological action of some common essential oil bearing plants used in Indigenous Medicine Part II Pharmacological action of *Alpinia galanga*, *Pistacia integrima*, *Piper betel*, and *Nardostychas jatamansi* Chopra, I. C , Jamwal, K S & Khajuria, B. N , *Ind Jour Med. Res* , 1954, 42, 385

ARGEMONE MAXICANA.

Removal of toxic alkaloids from Argemone oil and oils adulterated with argemone oil. Roy, A C., *Jour. Sci. Indl Res* , 1954, **13B**, 376.

BERBERIS ASIATICA Roxburgh

Plant alkaloids Pt. VI, *Berberis asiatica* Roxburgh. Barva, A K & Das Gupta, A K , *Jour Ind Chem Soc* , 1954, 31, 253

CARUM ROXBURGIA

Pharmacological study of an Indigenous Drug 'Ajmod' *Carum Roxburgiana* Benth and Hook Gujral, M L , Bhargava, K P , Srivastava, R S and Kishore, K A., *Ind Jour Med Res* , 1954, 42, 389

CAESALPINIA LIGYNA Rottles

Chemical examination of the roots of *Caesalpinia Ligyna* Rottles Chaudhry, G. R , Sharma, V N & Dhar, M L , *Jour 'Sci. Ind. Res* 1954, **13B**, 150.

CHENOPODIUM AMBERIOSOIDES

A short note on *Chenopodium* raised in Kashmir. Handa, K. L Kapoor, L. D & Abrol Hira Lal. *Ind. Jour Pharm* , 1954, 16.

COFFEE POWDER.

Chemical standards for coffee powder. Narayanaier, S & Ram Chandran. *Curr. Sci* , 1954, 192, 23.

CURCURBITA PEPO

Chemical Examination of the oil from the seeds of Curcubita pepo.
Narayanmurthy, N L., Iyer, B H, Gangadharam P R J & Sirsi, M,
Ind Jour Pharm, 1954, 7, 148

DATURA METEL Linn.

Variation & alkaloid content due to difference in size of the leaves of
Datura Metel Linn. Chaudhuri, B. N., *Ind. Jour. Pharm*,
1954, 16, 34

ENTADA SCANDENS BENTH

A note on sulphur containing glucoside in the seeds of Entada
scandens Benth Rangaswamy and Subba Rao, *Ind Jour. Pharm.*
1954, 7, 152

FERONIA ELEPHANTUM

A new aldobiouronic acid 3—(Glucuronopyranosyl)—D-Galacto-
pyranose from Kata (Feronia elephantum) Gum. Mathur, G. P. &
Mukherjee, S, *Jour Sci Indl Res.*, 1954, **13B**, 452.

HYOSCYAMUS MUTICUS

Chemical investigation of Hyoscyamus muticus, Handa, K. L.,
Abrol, Hira Lal, *Jour Sci Indl Res*, 1954, **13B**, 376.

JATROPHA GLANDULIFERA Roxb.

Chemical examination of the seed oil of Jatropha glandulifera Roxb,
Sheth, M C, Desai, C M., *Jour Ind. Chem Soc*, 1954, 31, 407.

MELIODINUS MONOGYNUS Roxb

Chemical examination of the root bark of Melodinus monogynus
Roxb Part I. Chatterjee, S K, Sharma, V N & Dhar, M L, *Jour.*
Sci Indl Res., 1954, **13B**, 546

MENTHA ARVENSIS.

A short note on Japanese Mint raised in Kashmir Handa, K. L.,
Kapoor, L D & Abrol, H L, *Ind. Jour Pharm*, 1954, 16, 32.

MELIA AZEDARACH.

Chemical examination of the heart wood of Melia azedarach. Bholi
Nath, *Jour. Sci Indl Res*, 1954, **13B**, 740.

MUSUA FERRAE

Refining of Nahor seed oil (Musua ferrae) Kusturi, T. R., Narayana
Murty, N L and Ujer, B. H, *Jour. Sci. Indl Res.*, 1954, **13B**, 453.

OCIMUM KILIMANDSCHARICUM Guerke.

Camphor and camphor oil from Ocimum Kilimandscharicum Guerke.
Choudhry, J. K., *Sci. & Cult*, 1954, 19, 354.

POLIANTHESE TUBEROSA Linn.

Glucofructosan from Polianthese tuberosa Linn and Garlic (*Allium sativum* Linn.), Srinivasan, M., and Bhatia, I. S., *Curr. Sci*, 1954, 23, 192.

PARMELIA KAMTSCHADALIS.

Chemical investigation of the Lichens . *Parmelia kamtschadalis* and *Parmelia arnoldii* Shah, Latika, G (Mrs). *Jour Ind. Chem. Soc*, 1954, 31, 311.

RAUWOLFIA CANESCENS.

The alkaloids of *R. canescens*, Part VI Yobyrine, the Se-Dehydrogenation Product of Rauwolscine. Chatterjee, A & Pakrashi, S; *J. Ind. Chem. Soc* 31, 25, 1954

The Alkaloids of *R. canescens*, Part VII Studies on the I R Spectra of Rauwolscine and its Acetyl derivatives Chatterjee, A. & Pakrashi, S. *J. Indian Chem. Soc* 31, 29, 1954.

The Alkaloids of *R. canescens* Part VIII. SeO_2 -Oxidation of Yobyrine, the Se-Dehydrogenation Products of Rauwolscine. Chatterjee, A. & Pakrashi, S, *J Ind Chem Soc* 31, 31, 1954.

Conformation of Rauwolscine, allo-Yohimbine and their Congeners, Identification of Rauwolscine with α -Yohimbine Chatterjee, A, Bose, A K. & Pakrashi, S. *Chem and Industry* 1954, 491

On the Stereochemistry of Rauwolscine, the Alkaloid of *R. canescens*. A. Chatterjee and S Pakrashi. *Naturwiss.* 41, 215, 1954.

The Isolation of Reserpine from *R. canescens* M W Klohs, Draper, M D. Keller, F. Petracek F J *J A C.S.* 76, 1381, 1954

RAUWOLFIA SERPENTINA.

Reserpin, der sedative Wirkstoff aus *R. Serpentina*. J. M Mueller, E. Schlittler & H.J. Bein. *Exper.* 8, 338, 1952

On the Constitution of Reserpine from *R. Serpentina* L Dorfman, C F. Huebner, H B. MacPhillemy, E Schlittler & A F. St. Andre. *Exper* 9, 368, 1953.

Chemistry of the Rauwolfia Alkaloids, including Reserpine. E Schlittler, H B MacPhillamy, L. Dorfman, A Furlenmeier, C F Huebner, R Lucas, J M Mueller, R Schwyzet and A.F St. Andre *Ann. N Y Acad. Sciences* 59, 1, 1954

Structure of Reserpine. N. Neuss, H E Boaz, J W. Forbes : *J A C S* 75 4870, 1953

Rauwolfia serpentina Alkaloids ; Structure of Reserpine. N Neuss. H E Boaz & J W Forbes : *J A C S* 76, 2463, 1954

Rescinnamine, a new hypertensive and sedative Principle. Klohs, M.W, M.D. Draper & F Keller; *J A C.S.* 76, 2483, 1954,

Reserpinine, ein neues Alkaloid aus *Rauwolfia serpentina*. E Haack, A. Popelak, H. Spingler & F. Kaiser, *Naturwiss* 41, 214, 1954.

- Sarpagin, ein neues Alkaloid aus R. Serpentina. A. Stoll und A. Hofmann *Helv. Chim. Acta* 36, 1143, 1954
- Neue Alkaloids aus R. serpentina. A. Popelak, H Spingler und F. Kaiser *Naturwiss* 40, 625, 1953.
- Essai de Classification des Alcaloides des Rauwolfias (Apocynaceae). Raymond-Hamet *C r Seances hebd Acad Sci.* 237, 1435, 1953
- S Bose . *J. Indian Chem. Soc* 31, 47, 1954 Rauwolfinine, a New Alkaloid of R. serpentina, Part I.
- Rauhimbii und iso-Rauhimbii, zwei neue Alkaloide aus R. serpentina A Hofmann. *Helv Chim Acta* 37, 314, 1954
- Isolation of S-Yohimbine and a New Related Alkaloid from R serpentina F L Weisenborn, M Moore, P A Diassi *Chem. and Industry*, 1954, 375.
- A Quel Type d' Alcaloide la Raubasine appartient-elle Raymond-Hamet : *C R Seances hebd Acad. Sci* 238, 1338, 1954.
- Rauwolfia Alkaloids IX Isolation of Yohimbine from R serpentina F E. Bader, D Dickel & E. Schlittler, *J A.C S* 76, 1695, 1954.
- Alkaloids of R serpentina II The Isolation of Naturally occurring Py-Tetrahydroserpentine (Ajmalicine) and a Contribution towards its structure M W. Klohs, M D Draper, F. Keller, W Malesh & F J Petrcek *J A.C S* 76, 1332, 1954
- A Preliminary investigation of R.serpentina W L Holt & Costello C H , *J. Amer Pharm. Ass* 43, 144, 1954
- Chemistry of Ajmaline, Part I. Comp Chem and Industry, 1952, 442. F A L Anet, D Chakravarti, R Robinson and E Schlittler . *J.C S (London)* 1954, 1242
- The Constitution of Ajmaline, Part I Chatterjee, A. & Bose, S *J. Indian Chem. Soc* 31, 17, 1954
- Reserpinin, ein neues Alkaloid aus R Serpentina E Schlittler, H Saner & J M. Meller . *Exper.* 10, 133, 1954
- The Constitution of Sarpagine A F Thomas . *Chem and Industry* 1954, 488.
- Rauwolfia Alkaloide III Die Isolierung weiterer Alkaloide aus R. serpentina A Hofmann *Helv Chim Acta* 37, 849, 1954.
- Method for Determination of Serpasil H Sheppard, G.L Wagle & A J. Plummer : *Fed Proc.* 13, 404, 1954.
- The Determination of Serpasil in Blood Urine and Tissues (to be published). H. Sheppard, N. Smith, L Stannard & G Wagle.
- Zur Pharmakologie des Reserpins, eines neuen Alkaloids aus R. serpentina H J Bein . *Exper.* 9, 107, 1953
- Pharmacological Effects of Reserpine, a New Crystalline Pure Alkaloid from R serpentina, in the Dog J H Trapold, M.W. Osborne & F F. Yonkman . *Fed. Proc.* 12, 373, 1953.

Sedative and Hypnotic action of Reserpine in Unanesthetized Animals A.J. Plummer, W. Barrett, G. Wagle & F.F. Yonkman: *Fed Proc* 12, 357, 1953.

Pharmakologische Untersuchungen Über verschiedene Alkaloide (Serpasil, Ajmalin, Serpentin and Serpentinin) aus *R. serpentina*. H. J. Bein und F. Gross *Verh Ber. Dtsche Ges Kreislaufforsch* 19, 277, 1953.

Experimentelle Untersuchungen über "Serpasil" (Reserpin), ein neues, sehr wirksames Rauwolfia-Alkaloid mit neuartiger zentraler Wirkung H. J. Bein, F. Gross, J. Tripod und R. Meier: *Schweiz. Med Wschr* 83, 1007, 1953

Wirkung von Reserpin (Serpasil) auf isolierte Kreislauforgane. J. Tripod und R. Meier *Arch internat Pharmacodyn* (In Press)

Cardiovascular and Respiratory Effects of Serpasil, a New Crystalline Alkaloid from *R. serpentina* J. H. Trapold, A. J. Plummer & F. F. Yonkman *J. Pharm. Exper Therap* 110, 205, 1954

Additional Pharmacological studies on Serpasil, a Crystalline Alkaloid from *R. serpentina*, in the Dog J. H. Trapold, M. W. Osborne, A. J. Plummer & F. F. Yonkman *J. Pharm Exper. Therap.* 110, 49, 1954

Sedative and Hypertensive Actions of a Group of Esters Related to Serpasil A. J. Plummer, W. Barrett & R. Rutledge: *Fed Proc* 13, 395, 1954 :

Effects of Serpasil (Reserpine) on Monkeys (Motion Picture). A. E. Earl, R. C. Dibble & R. D. Wolf *Fed Proc* 13, 350, 1954

Action of Serpasil on Gastrointestinal Motility and Secretion W. E. Barrett, R. A. Rutledge & B. Rogie, *Fed Proc.* 13, 334, 1954,

Behavioral and Electroencephalographic Studies with Serpasil (Reserpine), a New Alkaloid from *R. serpentina* J. A. Schneider & A. E. Earl *Fed Proc* 13, 130, 1954

Effet Hypotenseur chez l'Animal de la Reserpine, un nouvel Alcaloïde de la *R. serpentina* R. Meier, H. J. Bein, F. Gross, J. Tripod & H. Tuchmann-Duplessis *Cr. Sceances hebdomadaires Acad Sci.* 238, 527, 1954

Effets de la Reserpine, un nouvel Alcaloïde de la *R. serpentina*, sur le Système nerveux central de l'Animal R. Meier, H. J. Bein, F. Gross, J. Tripod et H. Tuchmann-Duplessis *Cr. Sceances hebdomadaires Acad Sci* 238, 961, 1954.

Hypotensive Agents from *R. serpentina* Reserpine and other Alkaloids G. Croneheim, C. Stipp & W. Brown: *J. Pharm. Exper. Therapy* 110, 13, 1954

Characterization of Central Effects of Serpasil and their Antagonistic Reactions. J. Tripod, H. J. Bein und R. Meier: *Arch. intern. Pharmacodyn* 96, 406, 1954 :

Pharmacology of Rauwolfia Alkaloids including Reserpine. A. J. Plummer, A. E. Earl, J. A. Schneider, J. Trapold & W. Barrett: *Ann. N Y Acad. Sciences* 59, 8, 1954.

- Endocrine Aspects of the Pharmacology of Reserpine Gaunt, F., Renzi, A.A., Antonchak, N., Miller, G. & Gilman, M. *Ann N Y. Acad. Sciences* 59, 22, 1954
- Effects of Reserpine (Serpasil) on behavior and autonomic regulating mechanisms in the monkey. Tripod, J und Meier: *Arch Internat Pharmacodyn* (In Press)
- Cardiovascular and Renal Hemodynamic Response (acute and chronic) to Reserpine. E. Dennis, W Hughes & J Moyer *Fed Proc.* 13, 347, 1954
- Effect of Reserpine (Serpasil) on "sham rage" in the cat J A Schneider (to be published)
- Mouse Prosis bioassay of Rauwolfia serpentina for reserpine-like activity B Rubin & J C Burke, *Fed Proc* 13, 400, 1954.
- Changes in convulsant thresholds after Rauwolfia serpentina, Reserpine and Veriloid E H. Jenney *Fed. Proc* 13, 370, 1954
- Sedative and Antihypertensive effect of Intravenous Reserpine in Man A M Sellers and J H Hafkenschiel, *Fed Proc* 13, 404, 1954
- Pharmacological investigation with Serpine Das Gupta, S R & G. Werner *Bull Calcutta School Trop Med* 1, No, 4, 1954.
- Indian Medicinal Plant in the Treatment of Hypertension. Chakravarti, M D, *Brit Med. J* 1953, 1390
- Neue Alkaloide aus R serpentina J D Achelis und G Kroneberg. *Naturwiss* 40, 625, 1953,
- Alterations in Cardiovascular Responses of the Dog following Rauwiloid, an Alkaloid Extract from R serpentina J T Gourzis, J R Sonnenschein & R. Barden *Proc Soc Exptl Biol Med* 85, 463, 1954.
- A Preliminary Note on the Pharmacological investigation of the Hypotensive Action of R serpentina J W Nelson & C A Schlogel. *J Amer Pharm Ass* 42, 324, 1953
- Hypertensive Action of another R serpentina Alkaloid (Serpasil) R. J Vakil *J Indian Med. Assoc* 23, 97, 1953
- Combinations of Drugs in the Treatment of Essential Hypertension R. Wilkins *Mississippi Doctor* 1953. 359
- Hochdruck behandlung miteinem Rauwolfia alkaloid (Serpasil) W. Loffler, A F Essellier, F. Prott und A Wegmann *Schweiz Med. Wschr* 83, 1012, 1953
- Reserpine and the Alseroxylon Alkaloids of R Serpentina in Hypertension. T. Winsor *Arizona Med.* 10, 419, 1953
- Ueber die Behandlung der arteriellen Hypertonie mit Reserpin G Damm und H. Trautner *Dische Med Wschr* 79, 39, 1954

Clinical usage of Rauwolfia Alkaloids, including Reserpine (Serpasil). Wilins, R.: *Ann N. Y. Acad. Sciences* 59, 36, 1954.

Clinical and Experimental Effects of Reserpine in Patients with Essential Hypertension. E. D. Freis & R. Ari: *Ann. N. Y. Acad. Sciences* 59, 45, 1954

Intravenous Reserpine in patients with Essential Hypertension. H. H. Hafkenschiel & A. M. Sellers *Ann. N. Y. Acad. Sciences* 59, 54, 1954

Human Pharmacology of Reserpine. T. Winsor: *Ann. N. Y. Acad. Sciences* 59, 61, 1954.

Clinical use of Reserpine (Serpasil) in Geriatrics. J. H. Moyer: *Ann. N. Y. Acad. Sciences* 59, 95, 1954.

Use of Reserpine (Serpasil) in Certain Gynaecologic Disorders. R. B. Greenblatt *Ann. N. Y. Acad. Sciences* 59, 133, 1954.

The Modern Treatment of Hypertension *Comp. Med. Times* 81, 741. S. W. Hoobler, *Univ. Mich. Med. Bull.* 20, 1, 1954

Recent Developments in the Treatment of Hypertension. E. D. Freis: *Med. Clin. N. A.* 38, 363, 1954

Reserpine in the Treatment of Hypertension A note on the Relative Dosage and effects R. W. Wilkins, W. E. Judson, W. Hollander, W. E. Huckabee & J. H. Friedman *New England J. Med.* 250, 477, 1954.

Observations of Cardiovascular and Renal Hemodynamic Responses to Reserpine and Clinical Results in Treatment of Hypertension J. H. Moyer, W. Hughes & R. Huggins *Clin. Res. Proc.* 2, 72, 1954.

Essais Cliniques sur l'action du Serpasil dans l'hypertension arterielle grave O. Ritter und J. C. Huguenin (In print)

Treatment of Hypertension with Serpasil, a R. serpentina alkaloid. L. Hensel: *Schweiz. Med. Wschr.* 83, 1162, 1953.

Clinical Experience with Reserpine (Serpasil) A controlled study. H. P. Dustan, R. D. Taylor, A. C. Corcoran & I. H. Page. *Ann. N. Y. Acad. Sciences* 59, 136, 1954.

Reserpine alone and in combination with Hydralazine and Hexamethonium in the Treatment of Hypertension W. Hughes, R. McConn and E. Dennis *Fed. Proc.* 13, 368, 1954.

Studies on "Essential" Hypertension V. An endocrine syndrome. H. A. Schroeder and D. F. Davies: *Ann. Int. Med.* 40, 516, 1954.

Über die perorale medikamentöse Dauerbehandlung des chron. Hochdruckleidens. R. Franke & F. Thiele, *Die. Med.* 50, 1626, 1953.

The Medical Management of arterial Hypertension. E. Meilman: *N. Engl. J. Med.* 248, 894, 1953.

Excretion of Rauwolfia Total Alkaloids in Urine. Gupta, J. C., Roy, P. K., Ray, G.K. & Ganguly, S.C. *Indian J. Med Res.* 38, 67, 1950.

Die Behandlung der chronischen arteriellen Hypertonie. O.H Arnold *Therapie der Gegenwart* 1952, 167.

Behandlung der Hypertonie mit der indischen R serpentina F. Vida *Die Medizinische* 37, 1157, 1952.

A comparison of Results in the Treatment of Hypertension with R. serpentina alone and in combination with Hexamethonium or Hydralazine I H Moyer, R.W Food, W R Livesay & S I Miller *J Lab and Clin Med* 42, 926, 1953

Extracts of R serpentina in Hypertension. R W. Ford & J H. Moyer *General Practitioner* 8, 51, 1953.

Treatment of Hypertension with R serpentina alone and in combination with Hexamethonium or Hydralazine J H. Moyer and W. R. Livesay *Amer J Med.* 16, 605, 1954

The Pharmacodynamics of Postural Hypertension (Rauwolfia) R A Schneider et al. *Clin Res Proc* 2, 69, 1954.

Effects of Orally Administered "Veriloid" and "Rauwiloid" on the Blood Pressure of Hypertensive Individuals C T Bello & L. W Turner *Clin Res Proc* 2, 70, 1954

Response of Metacorticoid Hypertension to Bistrium, Apresoline Veriloid and Serpentina F M Sturtevant, *Proc. Soc. Expt Biol Med* 84, 101, 1953

Preliminary observation on R. serpentina in hypertensive patients R W Wilkins, W E Judson & I R Stanton *Proc New Engl Cardiorasc Soc* 34, 1951

R Serpentina in the Treatment of Arterial Hypertension F Klausgraber, *Wien, Med Wschr* 83, 995, 1953

New Drug Therapies in arterial Hypertension. R W Wilkins *Ann Internal Med* 37, 1144, 1952

Preliminary observation of Rauwolfia serpentina therapy of Hypertension R W Ford, W R Livesay, S I. Miller & J H Moyer *Med Res and Ann.* 47, 608, 1953

Antihypertensive Therapy. R. P. McCombs: *Bull Neu Engl. Med Center* 15, 129, 1953.

The use of Drugs in the Management of Hypertension Samuel S Riven & J G. Wells *J. Tenn. State Med Ass* 47. 1, 1954

Management of Essential Hypertension. W. C. Gettelfinger *J Kentucky State Med Ass.* 51, 434, 1953

Rauwolfia, die Neuentdeckung einer alten Heilpflanze. E. Steinegger *Bulletin Galenica* 17, 50, 1954.

I Soprauna Sostanza ad Azione Sedativa Isolate della R. serpentina. A. Lattanzi *Omnia Therapeutica (Italy)* 5, fasc.

Schneider und A.J. Plummer E Schlittler, J A.

Ueber Rauwolfia-Alkaloide Angew. Chemie (In Print).

Zur Wirkung von R. serpentina. G Werner *Arzneimittelforschung* 4, 40, 1954.

Some Modern Hypertensive Agents. R W Richards : *Pharm J.* 172, 269, 1954

Chemical and Medicinal Aspects of R. serpentina R. Kress : *Pharmazie* 8, 726, 1953.

Indian Medicinal, Plant in the Treatment of Hypertension (R. serpentina) Chakravarti, M D.: *Brit. Med. J.* 1953, 1390.

New Results in the Pharmaceutical Treatment of Arterial Hypertension (Part I & II). O.H. Arnold & K D. Bock . *Dtsche. Med. Wschr* 78, 565, 879, 1953

G L. Maisson : *Lancet*, 1953, 1308 The Treatment of Hypertension.

Essential Hypertension . Editorial *Med Times*, 81, 226, 1953.

Hypertension Editorial *Canad. Med Ass J* 68, 287, 1953.

Drug Therapy in Management of Hypertension. C. Levy . *Delaware State. Med J.* 26, 36, 1954.

R. Serpentina and other species G.E. Trease & W.C Evans *Pharmaceut. J.* 172, 351, 1954

A Pharmacognostical study of Rauwolfia see also Pharm 125, 186, 1953. H.W Youngken Sr *J. Amer Pharm. Assoc.* 43, 70, 1954.

Malabar Rauwolfia, R. micrantha, H.W Youngken Sr. *J. Amer. Pharm. Assoc.* 43, 141, 1954

The pharmacognosy of Rauwolfia T.E. wallis & S. Rohatgi : *J. Pharm Pharmacol.* 69, 334, 1932

Ueber R serpentina, eine neue Droge in der Behandlung des Hochdrucks. G. Meyer *Dtsche. Apoth. Ztg.* 92, 455, 1952.

The use of R. serpentina in Neuropsychiatric Conditions. N.S Kline *Ann N Y. Acad. Sciences* 59, 107, 1954.

Effects of R serpentina on Maniac patients. P.K. Roy . *Indian J. Neurol Psychiat.* 2, 59, 1950.

Two cases of Paranoia treated with Rauwolfia serpentina. C K. Nagendranath : *Ind. J. Neurol. and Psychiatry* 2, 62, 1950.

The use of R. serpentina in Psychiatry. P K. Roy *Ind. J. Neurol. and Psychiatry*, 3, 380, 1952.

Recent development on *Rauwolfia serpentina* Benth. Rajgopalan, S : *Jour. Sci Indl Res.* **13B**, 17, 1954.

Rauwolfia serpentina. Iswariah, V *Curr. Sci* 23, 253, 1954

Rauwolfinine. A new Alkaloid of *Rauwolfia serpentina* Benth. Bose, S.: *Jour Ind Chem Soc* , 31, 311, 1954

RICE HUSK

Studies on rice husk, Betel husk and Bamboo lignins Pt II Nitrolignins Part III. Ali, M. Erfan. & M.H Khundkar *Jour. Ind Chem Soc.*, 31, 471 & 474, 1954.

SCILLA INDICA

A note on the isolation of crystalline Glycoside from commercial Indian Squill Rangaswami, S & Subramanian, Sankara, S *Jour Sci Indl Res* **13B**, 150, 1954

STROPHANTHUS SEEDS

Pharmacognosy of some *Strophanthus* seeds from India Gupta, B. & Bal, S N *Indian Jour. Pharm* , 16, 74, 1954

USNEA HIRTA

Chemical components of *Usnea hirta* Subha Rao, V. . *Indian Jour. Pharm* , 7, 151, 1954.

INDEX

OF

SCIENTIFIC NAMES

	PAGE
A	
Abies pindrow Spach. . .	125
Abroma augusta Linn . . .	21, 112, 125
Abrus precatorius Linn . . .	21, 70, 96, 101, 111, 125
Abutilon avicennae Gaertn. .	126
Abutilon indicum G Don . .	101, 126
Abutilon theophrastic Medic .	101
Acacia Willd .	126
Acacia catechu Willd .	126
Acacia chundra (Willd) DC .	126
Acacia ferruginea DC .	101
Acacia leucophloea Willd . .	126
Acacia pennata Willd .	70, 115
Acalypha Linn . .	126
Acalypha indica Linn. . .	127
Acanthaceae .	254
Achillea millefolium Linn .	127
Achras sapota Linn. .	127
Achyranthes aspera Linn. .	127
Aconitum Linn .	21, 127
Aconitum anthora Linn .	127
Aconitum balfourii Stapf .	21, 62, 128
Aconitum chasmanthum Stapf .	22, 62, 128
Aconitum deinorrhizum Stapf .	22, 62, 128
Aconitum elwesii Stapf .	62
Aconitum falconeri Stapf .	62
Aconitum ferox Wall. . .	22, 62, 109
Aconitum heterophyllum Wall. .	22, 127, 128
Aconitum laciniatum Stapf. .	22, 62
Aconitum laeve Royle . .	62
Aconitum lethale Griff . .	62
Aconitum luridum Hk. f & T .	62
Aconitum moschatum Stapf. .	62
Aconitum napellus Linn . .	128
Aconitum spicatum Stapf .	22, 62
Aconitum soongaricum .	62, 128
Aconitum violaceum Jacq. . .	22, 62
Acorus calamus Linn. . .	22, 93, 101, 115, 254

	PAGE
<i>Acorus gramineus</i> Soland .	93, 115, 128
<i>Acronychia laurifolia</i> Blume .	67, 115
<i>Acronychia pedunculata</i> (Linn) Miq	67, 115
<i>Actaea spicata</i> Linn	62, 128
<i>Actinodaphne hookeri</i> Meissn ..	23, 129
<i>Actinomyces griseus</i>	52
<i>Adansonia</i> Linn .	129
<i>Adenanthera pavonina</i> Linn	129
<i>Adenia (modecca) palmata</i> Engl .	74
<i>Adensonia digitata</i> Linn .	101
<i>Adenia wightiana</i> Engl ..	74
<i>Adhatoda vasica</i> Nees .	19, 23, 96, 101, 129, 254
<i>Adiantum lunulatum</i> Burm	101
<i>Adina cordifolia</i> Benth & Hk f	77, 101, 115, 129
<i>Adonis aestivalis</i> Linn .	63
<i>Adonis chrysocyathus</i> H f & T ..	63
<i>Aegle marmelos</i> Corr	23, 96, 101, 129
<i>Aesculus</i> Linn	130
<i>Aesculus hippocastanum</i> Linn .	130
<i>Aerothercium lanatum</i> Wakker	130
<i>Agaricus compestris</i> Linn ..	55
<i>Agaricus ostreatus</i> (Jacq) Fries .	101
<i>Agave americana</i> Linn	91, 115, 130
<i>Ageratum conyzoides</i> Linn	101, 131
<i>Ailanthus altissima</i> (Mill) Swingle .	68, 101, 112
<i>Ailanthus excelsa</i> Roxb .	101
<i>Ailanthus glandulosa</i> Desf	68, 101, 112
<i>Ailanthus malabarica</i> DC .	101
<i>Alangium begoniaefolia</i> Roxb .	131
<i>Alangium lamarckii</i> Thw .	23, 96, 131
<i>Alangium salviifolium</i> (Linn f) Wang	23, 96
<i>Albizzia chinensis</i> (Osbeck) Merr .	115
<i>Albizzia julibrissin</i> Durazz ..	96
<i>Albizzia lebeck</i> Benth ..	96, 101
<i>Albizzia procera</i> Benth. ..	70, 115
<i>Albizzia stipulata</i> Boiv	115
<i>Algae</i> ..	52
<i>Allamanda cathartica</i> Linn. .	80
<i>Allium cepa</i> Linn. ..	96, 131
<i>Allium minus</i> .	132
<i>Allium odorum</i> Linn. .	131, 132
<i>Allium porrum</i> Linn ..	132
<i>Allium sativum</i> Linn. ..	23, 91, 96, 132, 133, 256

	PAGE
<i>Allium schoenoprasum</i> Linn. .	97
<i>Alocasia indica</i> Schott .	93
<i>Alocasia macrorhiza</i> Schott .	93
<i>Alocasia montana</i> Schott .	93
<i>Alocasia odora</i> (Roxb) C Koch .	93
<i>Aloe abyssinica</i> Lam	133
<i>Aloe barbadensis</i> Mill	24, 111
<i>Aloe species</i>	92
<i>Aloe Tourn ex</i> Linn	133
<i>Aloe vera ex</i> Linn	24, 134
<i>Alpina galanga</i> Willd Swartz	24, 97, 254
<i>Alstonia</i> Br. .	134
<i>Alstonia scholaris</i> R Br	24, 101, 134
<i>Alsophila glabra</i> Hook	134
<i>Althaea officinalis</i> Linn	134
<i>Althaea rosea</i> Linn Cav .	101, 134
<i>Amanita muscaria</i> Pers .	54, 55, 74
<i>Amanita pantheriana</i>	57
<i>Amanita phalloides</i> Secr .	55, 57
<i>Amarantus blitum</i> Linn .	134
<i>Amaranthus gangeticus</i> Linn ..	101, 134
<i>Amaranthus tricolor</i> Linn .	101
<i>Amaryllidaceae</i>	91
<i>Amito pers</i> Kurrum	55
<i>Amito ostreatus</i> Jacq ..	55
<i>Ammannia baccifera</i> Linn	74
<i>Ammannia senegalensis</i> Lamk	74
<i>Amomum subulatum</i> Roxb	24
<i>Amomum xanthioides</i> Wall	101, 134
<i>Amorphonphallus campanulatus</i> (Roxb) Bl.	93
<i>Amorphonphallus lyratus</i> Engl	93
<i>Amorphonphallus sylvaticus</i> (Roxb) Kunth	93
<i>Anaboena</i> sp .	53
<i>Anacardiaceae</i> .	69
<i>Anacardium occidentale</i> Linn. .	69, 112, 115, 134, 135
<i>Anagallis arvensis</i> Linn	79, 112, 115, 135
<i>Ananas comosus</i> Merr.	91
<i>Ananas sativus</i> Schult .	91
<i>Anamirta cocculus</i> (Linn.) W & A ..	64, 115, 136
<i>Anchusa tinctoria</i> Lam. ..	135
<i>Andrachne cordifolia</i> Muell. Arg ..	87, 115
<i>Andrographis echinoides</i> Nees. ..	136
<i>Andrographis paniculata</i> Nees. .	24, 101, 136

	PAGE
Andropogon Linn.	136
Andropogon jwarancusa Jones.	136
Andropogon kuntzeanus Hack.	136
Andropogon nardus Linn	117
Andropogon squarrosus Linn.	136
Anemone obtusiloba D Don.	63
Anethum soa Roxb	137
Angelica archangelica Linn.	137, 254
Anisomeles malabarica Br.	137
Annonaceae	64
Annona muricata Linn.	101, 137
Annona reticulata Linn	64, 102, 115, 137
Annona squamosa Linn.	64, 111, 115, 137
Anogeissus latifolia Wall	137
Anthemis cotula linn	77, 112
Anthemis nobilis Linn	137
Anthocephalus cadamba Miq S & H	102
Anthocephalus indicus A Rich	102
Antiaris toxicaria Leschen	25, 89, 102, 138
Apama tomentosa Engl	115
Aphanizomenon flos-aquae (Linn) Ralfs	53
Apium graveolens Linn	76, 111, 112, 138
Apocynaceae	80
Aquilaria agallocha Roxb	138
Aquilegia vulgaris Linn	63
Araceae	93
Arachis hypogea Linn	25, 138
Araliaceae	76
Archangelica officinalis Hoffm	138
Areca catechu Linn	25, 92, 111, 138, 139
Arenga obtusifolia Mart.	92, 115
Argemone alba Lestib.	139
Argemone mexicana Linn.	60, 64, 139, 254
Argyreia speciosa Sweet	140
Arisaema speciosum Mart	93, 112, 115
Arisaema tortuosum Schott.	93, 112, 115
Aristolochia bracteata Retz.	86, 111, 116
Aristolochia indica Linn	25, 86, 111, 140
Aristolochiaceae	86, 140
Arnica montana Linn	141
Arnica Rupp. ex Linn.	140
Arundo donax Linn.	134
Artabotrys Br.	141

	PAGE
Artabotrys suaveolens Blume. . .	109, 141
Artocarpus hirsuta Lamk . .	142
Artemisia Linn. .	26, 141
Artemisia absinthium Linn. . .	26, 77, 116, 142
Artemisia dracunculus Linn . .	26
Artemisia kurramensis. . .	142
Artemisia maritima Linn . .	77, 142
Artemisia pallens Wall. ex DC .	26
Artemisia sacrorum Ledele. .	26, 142
Artemisia scoparia Waldest. & Kit. .	142
Artemisia vulgaris Linn . . .	26, 77, 111, 116, 142
Asarum europeum Linn . .	143
Asclepiadaceae . .	81, 143
Asclepias curassavica Linn .	81, 97, 116
Asparagus Linn .	143
Asparagus adscendens Roxb .	97, 102, 143
Asparagus officinalis Linn . .	112
Asparagus racemosus Willd . .	97, 102, 143
Aspergillus clavatus . .	54
Aspergillus fumigatus . .	54
Aspidium filixmas . .	58
Asteracantha longifolia Nees . .	27, 102, 143
Asystasia coromandeliana Nees . .	143
Atalantia monophylla Corr . .	143
Athyrium filix-foemina Roth .	144
Atriplex hortensis Linn .	144
Atrocarpus integrifolia Linn . .	142
Atropa Linn .	144
Atropa acuminata Royle . . .	27, 144
Atropa belladonna Linn . . .	83, 144, 145, 146, 147
Averrhoa carambola Linn . .	102
Avena fatua Linn . .	94
Avena sativa Linn. . .	94
Azadirachta Linn . .	147
Azadirachta indica A. Juss. . . .	27, 68, 97, 116, 147, 148

B

Bacitracin subtilis. . .	52
Bacopa monnieri (Linn.) Pennell. . .	27
Balanites aegyptiaca (Linn) Delile. Wall. . .	68, 116, 148, 149
Balanites roxburghii Planch . . .	68, 102, 116
Balanophora polyandra Griff. . .	149

	PAGE
<i>Baliospermum axillare</i> Blume. .	88
<i>Baliospermum montanum</i> muell Arg. .	88
<i>Balsamodendron mukul</i> Hook .	97, 149
<i>Bambusa</i> Schreb.	149
<i>Bambusa arundinacea</i> Willd. ..	27, 94, 116, 149
<i>Bambusa bambos</i> Druce .	27, 94, 116
<i>Barringtonia acutangula</i> (Linn) Gaertn .	73, 102, 116, 150
<i>Barringtonia asiatica</i> (Linn) Kurz .	73, 116
<i>Barringtonia racemosa</i> Bl (Linn.) Roxb.	73 116
<i>Barringtonia speciosa</i> Forst	73, 116
<i>Basella rubra</i> Linn.	102
<i>Bassia latifolia</i> Roxb ..	118, 150
<i>Bassia longifolia</i> Linn ..	41, 119, 150
<i>Bauhinia acuminata</i> Linn .	150
<i>Bauhinia racemosa</i> Lam.	97, 102
<i>Bauhinia tomentosa</i> Linn	102
<i>Bauhinia variegata</i> Linn ..	97, 102, 151
<i>Begoniaceae</i> .	76
<i>Begonia rex</i> Putzeys.	76
<i>Belamcanda chinensis</i> Leman ..	151
<i>Benincasa cerifera</i> Savi ..	151
<i>Berberidaceae</i> ..	64
<i>Berberis</i> Linn	151
<i>Berberis aquifolium</i> Pursh .	151
<i>Berberis aristata</i> DC ..	28, 64, 116, 152
<i>Berberis asiatica</i> Roxburgh .	28, 254
<i>Berberis coriaceae</i> Brandis. .	28
<i>Berberis floribunda</i> Hort Wall ..	28, 152
<i>Berberis himalaica</i> Arndt .	28, 152
<i>Berberis insignis</i> Hook f. & B	28, 152
<i>Berberis lycium</i> Royle .	28
<i>Berberis nepalensis</i> Spreng	41
<i>Berberis umbellata</i> Wall ..	29, 152
<i>Berberis vulgaris</i> Linn ..	29, 152
<i>Bergenia ligulata</i> (Wall.) Engl	102
<i>Betula</i> Tourn.	153
<i>Betula utilis</i> D Don. ..	153
<i>Beta vulgaris</i> Linn.	153
<i>Bidens tripartita</i> Linn ..	102
<i>Bignoniaceae</i> ..	84
<i>Bixaceae</i> ..	65
<i>Bixa orellana</i> Linn. .	153
<i>Blepharis edulis</i> Pers. .	29, 154

	PAGE
<i>Blumea balsamifera</i> DC	154
<i>Blumea densiflora</i> DC	154
<i>Blumea eriantha</i> DC	154
<i>Blumea lacera</i> DC	109, 154
<i>Blumea malcomii</i> Hook f	154
<i>Boerhaavia diffusa</i> Linn	154
<i>Boerhaavia repens</i> Linn	29
<i>Bombax malabaricum</i> DC	154
Boraginaceae	82
<i>Borassus flabellifer</i> Linn	155
<i>Boswellia serrata</i> Roxb	102
<i>Botrychium lunaria</i> sew	102
<i>Botrychium virginianum</i> Sew	155
<i>Bragantia tomentosa</i> Blume	115
<i>Bragantia wallichii</i> R Br	29, 155
<i>Brassica campestris</i> Linn	155
<i>Brassica cernua</i> (Thunb) Forbes et Hemsl	65, 102
<i>Brassica integrifolia</i> (West) O E Schulz	65
<i>Brassica juncea</i> Hook f & T	65, 155
<i>Brassica napus</i> Linn	65
<i>Brassica nigra</i> (Linn) Koch	65
<i>Brassica oleracea</i>	134, 156
Bromeliaceae	91
<i>Brosimum galactodendron</i> D Don. ex Sweet.	156
<i>Bryonia alba</i> Linn	156
Bryophyta	57
<i>Bryophyllum calycinum</i> Salisb	156
<i>Bucea amarissima</i> (Lour) Merr.	68, 102
<i>Bucea sumatrana</i> Roxb	68, 102, 156
<i>Bursera</i> Linn	156
<i>Butea</i> Roxb	156
<i>Butea frondosa</i> Koen ex Roxb	70, 116, 156, 157
<i>Butea monosperma</i> (Lam) Kuntze . .	29, 70, 97, 102, 116, 156
<i>Buxus sempervirens</i> Linn.	88
<i>Butea superba</i> Roxb.	29, 157

C

<i>Caesalpinia bonducella</i> Fleming .	30, 157
<i>Caesalpinia crista</i> Linn .	30
<i>Caesalpinia digyna</i> Rottl. .	158
<i>Caesalpinia jayabo</i> Maza. .	102

	PAGE
<i>Caesalpinia nuga</i> (Linn) Ait. . .	116
<i>Caesalpinia pulcherrima</i> Swartz. .	110
<i>Calendula officinalis</i> Linn.	158
<i>Callicarpa longifolia</i> Lam var <i>lanceolaria</i> C B. Clarke .	84, 116
<i>Callitris rhomboidea</i> R Br	158
<i>Calonyction muricatum</i> (Linn) G Don	82, 116
<i>Calophyllum inophyllum</i> Linn .	66, 116, 158
<i>Caloplaca elegans</i>	158
<i>Calotropis</i> Br .	158
<i>Calotropis gigantea</i> Linn.	
Dryand R Br	30, 81, 132, 111, 113, 158, 159
<i>Calotropis procera</i> Br. (Linn) Dryand	81, 111, 113, 159
<i>Caltha palustris</i> Linn .	63
<i>Calycopteris floribunda</i> Lam .	30, 159
<i>Camellia sinensis</i> Linn. (O Kunfize). .	30
<i>Campanulaceae</i>	78
<i>Cananga odorata</i> Hk F & T	159
<i>Cananga patchouli</i>	159
<i>Canarium commune</i> Linn .	159
<i>Canarium strictum</i> Roxb .	160
<i>Canavalia ensiformis</i> D C. .	70, 160
<i>Canavalia virosa</i> W & A	70
<i>Cannabis Tournef</i> .	160
<i>Cannabis indica</i> Lamk .	160
<i>Cannabis sativa</i> Linn	30, 89, 113, 116, 160
<i>Canscora decussata</i> Schult	97
<i>Cantharellus cibarius</i> Fr	55
<i>Capparis aphylla</i> Roth .	65
<i>Capparis decidua</i> Edg	65
<i>Capparis sepiaria</i> Linn . .	160
<i>Capparis spinosa</i> Linn .	97, 160
<i>Capparis zeylanica</i> Linn .	109
<i>Capparidaceae</i> . .	65
<i>Caprifoliaceae</i> .	76
<i>Capsella bursa-pastoris</i> Moench . .	161
<i>Capsicum annuum</i> Linn. .	83, 161
<i>Capsicum frutescens</i> Linn . .	83, 109
<i>Capsicum minimum</i> Roxb. . .	83, 109
<i>Carapa moluccensis</i> Lam. . .	103, 109
<i>Cardaria draba</i> Desv. . .	116
<i>Cardenia campanulata</i> Roxb. . .	118
<i>Cardiospermum halicacabum</i> Linn. . .	69
<i>Carex cernua</i> Boott. .	94

	PAGE
Careya arborea Roxb	73, 116
Caricaceae	75
Carica papaya Linn	31, 74, 111, 161, 162
Carica xanthin	161
Carthamus tinctorium Linn	162
Carum bulbocastanum Koch	103
Carum carvi Linn	31, 103, 162
Carum copticum Benth	50, 110, 162
Carum roxburgia ..	254
Caryophyllaceae ..	66
Caryophyllus aromaticus Linn ..	38, 163
Casearia graveolens Dalz	74, 116
Casearia tomentosa Roxb	75, 116, 164
Cassia absus Linn	31, 71, 163
Cassia acutifolia Delile .	71
Cassia alata Linn ..	71
Cassia angustifolia Vahl	31, 71, 109, 163
Cassia auriculata Linn	103
Cassia fistula Linn	71, 97, 100, 164
Cassia obovata Collad .	71
Cassia obtusifolia Linn	97
Cassia occidentalis Linn ..	164, 175
Cassia tora Linn .	97, 103, 164
Casuarina equisetifolia Linn ..	103, 165
Cassytha Linn	164
Cassytha filiformis Linn	87, 103, 116, 164
Catha edulis Forsk. ..	165
Ceasalpinia digyna Rottles .	254
Cedrela toona Roxb .	31, 103, 165
Cedrus deodara Loud .	165
Celastraceae	68
Celastrus paniculatus Willd	32, 111, 165
Celsia coromandeliana Vahl	103
Cenchrus pauciflorus Benth .	165
Centella asiatica (Linn) Urban. .	32, 76, 103
Centipeda orbicularis Lour .	165
Centratherum anthelminticum Kuntze .	32, 78, 116
Cephaelis ipecacuanha A Rich .	32, 77, 165
Cephalandra indica Naud .	34, 166
Cerbera manghas Linn. .	32, 80, 116
Cerbera odollam Gaertn. ..	32, 80, 116, 166
Ceropegia tuberosa Roxb .	103

	PAGE
<i>Chamaenerium angustifolium</i> Schur	167
<i>Cheiranthus cheiri</i> Linn	167
<i>Chenopodiaceae</i> . .	85
<i>Chenopodium</i> Linn	167
<i>Chenopodium album</i> Linn.	167
<i>Chenopodium ambrosioides</i> Linn	32, 33, 85, 167, 168
<i>Chenopodium botrys</i> Linn	33, 85
<i>Chionachne koenigii</i> (Spreng) Thw	97
<i>Chloroxylon swietenia</i> Dc	168
<i>Chrozophora rottleri</i> A. Juss ex Spreng	88
<i>Chrozophora tinctoria</i> Hook f . .	88
<i>Chrysanthemum coccineum</i> Willd	78, 116
<i>Chrysanthemum cinerariaefolium</i> Vis.	78, 116, 168
<i>Chrysanthemum roseum</i> Adam	78
<i>Cicer arietinum</i> Linn	16
<i>Cichorium intybus</i> Linn	33, 196
<i>Cicuta virosa</i> Linn . .	76, 169
<i>Cimicifuga foetida</i> Linn.	63, 97, 116
<i>Cimicifuga racemosa</i> Nutt	169
<i>Cinchona</i>	33
<i>Cinchona</i> Linn.	169, 170
<i>Cinchona calisaya</i> Wedd.	77, 111, 116
<i>Cinchona febrifuga</i>	170
<i>Cinchona ledgeriana</i> Moens ex Trimen	170
<i>Cinchona officinalis</i> Linn	77, 116
<i>Cinchona succirubra</i> Pavon	77, 117
<i>Cinnamomum</i> Bl	33, 170
<i>Cinnamomum camphora</i> Nees & Eberm	33, 87, 111, 117, 170
<i>Cinnamomum iners</i> Reinw	103
<i>Cinnamomum japonicum</i> Siebold	171
<i>Cinnamomum zeylanicum</i> Breyn . .	33, 171
<i>Citrum asiaticum</i> Linn	91
<i>Citrum lotifolium</i> Linn	91
<i>Cissampelos pareira</i> Linn . .	19, 33, 171
<i>Cissus setosa</i> Roxb . .	113
<i>Citrullus colocynthis</i> Schrad.	34, 75, 97, 111, 171
<i>Citrullus vulgaris</i> Schrad. . .	75, 171
<i>Citrus</i> Linn. . .	172
<i>Citrus acida</i> Roxb. . .	172
<i>Citrus aurantiifolia</i> (Christm.) Swingle.	34
<i>Citrus aurantium</i> Linn.	173
<i>Citrus bigaradia</i> Loisel. . .	172

	PAGE
<i>Citrus decumana</i> Linn	172
<i>Citrus limon</i> (Linn) Burm f	34, 172
<i>Citrus medica</i> var <i>limonis</i>	34
<i>Citrus poonensis</i> Hort ex Tanaka	173
<i>Claviceps purpurea</i> (Fries) Tulasne	173
<i>Clausena willdenowii</i> Wight & Arn	173
<i>Cleistanthus collinus</i> Benth & Hook f	88, 117
<i>Clematis gouriana</i> Roxb	63
<i>Clematis graveoleus</i> Lindl .	63
<i>Clematis napaulensis</i> DC	63
<i>Clematis orientalis</i> Linn.	63
<i>Clematis triloba</i> Heyne . .	63
<i>Clematis wightiana</i> Wall	63
<i>Cleome felina</i> Linn	65
<i>Cleome icosandra</i> Linn	34
<i>Cleome viscosa</i> Linn	65, 174
<i>Clerodendron infortunatum</i> Gaertn	34, 174
<i>Clerodendron serratum</i> . (Linn) Moon.	97
<i>Clitoria ternatea</i> Linn	71, 97, 103
<i>Coccinia cordifolia</i> Cogn	34, 97
<i>Coccinia indica</i> W & A	97
<i>Cocculus</i> DC .	174
<i>Cocculus hirsutus</i> Diels .	97
<i>Cocculus laurifolius</i> DC .	174
<i>Cocculus sarmentosus</i> Diels	174
<i>Coffea arabica</i> Linn	77
<i>Cochlearia armoracia</i> Linn .	174
<i>Cocos nucifera</i> Linn .	97, 174, 175
<i>Coffea</i> Linn	175
<i>Coffea robusta</i> Linn. .	175
<i>Colchicum</i> Linn .	175
<i>Colchicum autumnale</i> . .	14, 92
<i>Colchicum luteum</i> Baker .	14, 34, 92
<i>Colocasia virosa</i> Kunth	93
<i>Collybia albuminosa</i> (Berk) Betch	55
Combretaceae .	73
<i>Combretum pilosum</i> Roxb .	175
<i>Commelina asiatica</i> Linn . .	175
<i>Commiphora agallocha</i> Engler . .	97
<i>Commiphora mukul</i> Engl . .	35, 97
Compositae.	77
Coniferae. .	90
<i>Conorhinus rubrofasciatus</i> De Geer. .	248

	PAGE
Convolvulaceae ..	82
Convolvulus arvensis Linn	82
Convolvulus hamadae (Vved) Petrov	175
Convolvulus pluricantes Chois	175
Convolvulus scammonia Linn .	82
Coprinus comatus (Battara) Fr .	55
Coptis teeta Wall ..	35, 176
Corallocarpus epigaeus Benth ex Hook. f	75, 103
Corchorus Linn.	176
Corchorus capsularis Linn	103
Corchorus fascicularis Lam ..	103
Corchorus olitorius Linn	176
Cariandrum sativum Linn	97, 176
Coriariaceae ..	70
Coriaria nepalensis Wall ..	70
Corypha umbraculifera Linn. ..	92, 117
Coscinum fenestratum Colebr .	35, 176
Crassulaceae ..	73
Crataegus oxycantha Linn ..	176, 177
Crataeva nurvala Ham .	98, 177
Cressa cretica Linn .	98
Cretaria sp.	57
Crinum asiaticum Linn ..	177
Crocus sativum Linn	35, 91, 111, 177
Crotalaria juncea Linn	177
Crotalaria medicaginea Lam	177
Crotalaria spectabilis Roth .	177
Croton sp	89
Croton oblongifolius Roxb	88, 117
Croton tiglium Linn.	88, 117, 178
Cryptostegia grandiflora R Br	81, 178
Cubeba officinalis Rafin	178
Cucumis stativus Linn ..	75, 117, 178
Cucumis trigonus .	75, 111
Cucumis utilissimum Roxb .	178
Cucurbitaceae ..	62, 74, 179
Cucurbita pepo D. C ..	178, 255
Cuminum cyminum Linn ..	98, 103, 179
Cupressus torulosa Don. ..	179
Curcuma Linn ..	179
Curcuma amada Roxb. ..	179
Curcuma angustifolia Roxb. ..	98
Curcuma aromatica Salisb. ..	179

	PAGE
<i>Curcuma caesia</i> Roxb	179
<i>Curcuma domestica</i> Valetton ..	35, 179
<i>Curcuma longa</i> Linn. ..	35, 117, 179, 180
<i>Curcuma zedoaria</i> Rose .	35, 98, 180
<i>Cuscuta reflexa</i> Roxb ..	36, 82, 111, 180
<i>Cycas revoluta</i> Thunb ..	180
<i>Cyclamen persicum</i> Miller ..	79
<i>Cydonia oblonga</i> Mill. .	103
<i>Cydonia vulgaris</i> Pers. ..	103, 181
<i>Cylista scariosa</i> Roxb.	103
<i>Cymbopogon</i> Hack	181
<i>Cymbopogon caesius</i> Stapf ..	181
<i>Cymbopogon citratus</i> Stapf .	181
<i>Cymbopogon clandestinus</i> Stapf ..	181
<i>Cymbopogon coloratus</i> Stapf	181
<i>Cymbopogon martinii</i> Stapf .	182
<i>Cymbopogon nardus</i> (Linn) Rendle	117
<i>Cynanchum arnottianum</i> Wight	81, 117
<i>Cynanchum vincetoxicum</i> Pers .	81
<i>Cynodon dactylon</i> Pers	103
<i>Cynomarathrum nuttallii</i> A Gray .	182
Cyperaceae	94
<i>Cyperus esculentus</i> Linn .	182
<i>Cyperus scarisous</i> R. Br ..	103, 182
<i>Cyperus longus</i> Linn .	94
<i>Cyperus rotundus</i> Linn .	103, 182
<i>Cystopus candidus</i> .	182
<i>Cytisus scoparius</i> Linn .	71

D

<i>Daemia extensa</i> (Linn) R Br .	36, 183, 117
<i>Dalbergia stipulacea</i> Roxb .	71, 117
<i>Daphne cannabina</i> Wall. ..	87
<i>Daphne oleoides</i> Schreb. ..	87
<i>Datura</i> Linn .	187
<i>Datura alba</i> Nees. ..	183
<i>Datura fastuosa</i> Linn. .	83, 183
<i>Datura innoxia</i> Mill. ..	183
<i>Datura metel</i> Linn. ..	83, 183, 255
<i>Datura stramonium</i> Linn .	36, 83, 113, 183
<i>Datura tatula</i> Linn. ..	183, 184

	PAGE
<i>Daucus carota</i> Linn .	76, 104, 111, 113, 184
<i>Decalepis hamiltonii</i> W. & A. ..	184
<i>Delphinium ajacis</i> Linn .	113, 184
<i>Delphinium brunonianum</i> Royle .	63, 117
<i>Delphinium caeruleum</i> Jacquem ex Cambess	63, 117
<i>Delphinium elatum</i> Linn	63, 117, 184
<i>Delphinium vestitum</i> Wall. .	63
<i>Dendrocalamus strictus</i> (Roxb) Nees .	94
<i>Derris</i> Lour	184
<i>Derris elliptica</i> (Roxb.) Benth	71, 117, 185
<i>Derris ferruginea</i> (Roxb.) Benth	71, 117, 185
<i>Derris scandens</i> Benth	71, 185
<i>Derris trifoliata</i> Lour var <i>uliginosa</i> (Roxb ex Willd)	117
<i>Derris uliginosa</i> Benth	71, 117
<i>Desmodium gangeticum</i> DC.	104, 109
<i>Descurainia sophia</i> Linn	104
<i>Dichroa febrifuga</i> Lour .	36, 185
<i>Dichanthium annulatum</i> Stapf	185
<i>Dictamnus albus</i> Linn	113
<i>Dictamnus fraxinella</i> Pers .	185
<i>Didymocarpus pedicellata</i> Br .	36, 185, 186
<i>Digitalis</i> (Tourn) Linn .	186
<i>Digitalis lanata</i> Ehrh. .	36, 186
<i>Digitalis purpurea</i> Linn. .	84
<i>Dioscorea bulbifera</i> Linn.	91
<i>Dioscoreaceae</i> ..	91
<i>Dioscorea daemona</i> Roxb	91, 117
<i>Dioscorea deltoidea</i> Wall .	91, 117
<i>Dioscorea hispida</i> Dennst.	91, 117, 186
<i>Dioscorea prazeni</i> Prain & Burk .	91, 117
<i>Diospyros ebenum</i> Koenig	80, 117
<i>Diospyros embryopteris</i> Pers	186
<i>Diospyros melanoxylon</i> Roxb. ,	98, 104
<i>Diospyros montana</i> Roxb.	80, 117
<i>Diospyros paniculata</i> Dalz. ..	80, 117
<i>Diospyros peregrina</i> Gurkeim.	104
<i>Dipterocarpus indicus</i> Bedd .	186, 187
<i>Dodonaea viscosa</i> (Linn.) Jaxq .	69, 117, 187
<i>Dolichandrone falcata</i> Seem ..	84, 111, 117
<i>Dolichos lablab</i> Linn ..	187
<i>Dregea volubilis</i> Benth. ex Hook	37
<i>Droseraceae</i> .	73
<i>Drosera burmanni</i> Vahl. .	73

	PAGE
<i>Drosera peltata</i> Sm. var <i>lunata</i> Clarke	73
<i>Drosera spathulata</i> Labill.	73
<i>Drynaria quercifolia</i> J. Sm	98, 109
<i>Dryopteris barbigera</i>	58
<i>Dryopteris blanfordii</i>	58
<i>Dryopteris calcarata</i>	58
<i>Dryopteris marginata</i>	58
<i>Dryopteris odontoloma</i>	58
<i>Dryopteris schimperianum</i>	58
<i>Duranta plumieri</i> Jacq	84, 117
<i>Duranta repens</i> Linn.	84, 117

E

Ebenaceae	80
<i>Edgeworthia gardneri</i> Meissn	87, 117
<i>Eichornia</i> Kunth	187
<i>Elaeodendron glaucum</i> Pers	68
<i>Elaeocarpus serratus</i> Linn	104
<i>Elaeis guineensis</i> Jacq	187
<i>Elaeocarpus tuberculatus</i> Roxb	109
<i>Eleusine coracana</i> Gaertn	188
<i>Elettaria cardamomum</i> Maton	37, 98, 188
<i>Embelia</i> Burm	188
<i>Embelia tsjeriam-cottam</i> A DC.	98
<i>Embelia ribes</i> Burm	188
<i>Emblica officinalis</i> Gaertn.	188
<i>Enhydra fluctuans</i> Lour	188
<i>Enicostema littorale</i> Blume	188
<i>Entada phaseoloides</i> (Linn) Merr	37, 71
<i>Entada pursaetha</i> DC.	37, 117, 189
<i>Entada scandens</i> Benth	37, 71, 117, 255
<i>Entoloma microcarpum</i> Berk & Broome.	55
<i>Ephedra</i> Linn	189
<i>Ephedra gerardiana</i> Wall.	37
<i>Ephedra nebrodensis</i> Tin.	189
<i>Ephedra vulgaris</i> Rich.	37, 189
<i>Equisetum arvense</i>	58
<i>Eremostachys acanthocalyx</i> Boiss.	85
<i>Eremostachys superba</i> Royle ex Benth	117
<i>Eremostachys vicaryi</i> Benth.	85, 118
Ericaceae	79

		PAGE
<i>Erigeron candensis</i> Linn.	..	78, 104, 113, 100
<i>Ervatamia dichotoma</i> (Roxb) Blatter	..	80
<i>Erycibe paniculata</i> Roxb		109
<i>Erythrina indica</i> Lam.	..	190
<i>Erythroxylum coca</i> Lam		37, 67
<i>Erythroxylum monogynum</i> Roxb		109, 190
<i>Eucalyptus</i> Lher	..	190
<i>Eucalyptus</i> L W Erit		38
<i>Eucalyptus citriodora</i> Hook	.	190
<i>Eucalyptus globulus</i> Labill		73, 118, 190, 191
<i>Eugenia aromatica</i> Kuntze.	.	38
<i>Eugenia jambolana</i> Lam	.	191
<i>Eugenia operculata</i> Roxb.		108
<i>Eulophia</i> R Br		191
<i>Eulophia nuda</i> Lindl	.	98
<i>Euonymus tingens</i> Wall	..	38
<i>Eupatorium odoratum</i> Linn		78, 118
<i>Eupatorium uticifolium</i> L f		78
<i>Euphorbia</i> Linn		191
<i>Euphorbia acaulis</i> Roxb	.	88, 113
<i>Euphorbiaceae</i>		87
<i>Euphorbia antiquorum</i> Linn		88, 104, 113, 118
<i>Euphorbia cattimandoo</i> W Elliot		88, 113
<i>Euphorbia helioscopia</i> Linn.	..	88, 109, 113, 191
<i>Euphorbia hirta</i> Linn	.	38, 88, 104
<i>Euphorbia hypericifolia</i> Linn		88, 104
<i>Euphorbia nerifolia</i> Linn		88, 113, 118
<i>Euphorbia nivulea</i> Buh Ham		88, 113
<i>Euphorbia peplis</i> Linn		113, 191
<i>Euphorbia pilulifera</i> Linn		38, 291
<i>Euphorbia rothiana</i> Spreng.		113
<i>Euphorbia royleana</i> Boiss		88, 113, 118
<i>Euphorbia thomsoniana</i> Boiss	.	88, 113
<i>Euphorbia thymifolia</i> Linn.	.	88, 104, 118, 191
<i>Euphorbia tirucalli</i> Linn		88, 111, 113, 118, 192
<i>Euphorbia trigona</i> Haw.	..	88, 113
<i>Evolvulus alsinoides</i> Linn	..	104
<i>Excoecaria agallocha</i> Linn	.	88, 111, 113, 118
<i>Exidea polytricha</i> Mont.	.	55

F

<i>Fagonia cretica</i> Linn.	..	104, 109
<i>Fagopyrum esculentum</i> Moench	.	62, 86, 113
<i>Fagopyrum cymosum</i> Meissn.	..	192

	PAGE
<i>Fagopyrum tataricum</i> Gaertn . .	86
<i>Feronia elephantum</i> Corr.	104, 192, 225
<i>Feronia limonia</i> (Linn) Swingle	104
<i>Ferula</i> Linn .	192
<i>Ferula foetida</i> Regel.	38, 192
<i>Ferula jaeschkeana</i> Vatke. .	192
<i>Ferula narthex</i>	41
<i>Ficus anthelmintica</i> Martius .	192
<i>Ficus bengalensis</i> Linn	90, 105, 192
<i>Ficus carica</i> Linn .	193
<i>Ficus glomerata</i> Roxb. .	104
<i>Ficus heterophylla</i> Linn .	104
<i>Ficus hispida</i> Linn.	104
<i>Ficus indica</i> Linn	193
<i>Ficus laurifolia</i> Hort et Lam. .	130
<i>Ficus racemosa</i> Linn. .	194
<i>Ficus religiosa</i> Linn. . .	193
<i>Ficus</i> sp .	90
<i>Fimbristylis junciformis</i> Kunth	104
<i>Fistulina hepatica</i> Fr .	55
<i>Flacourtia indica</i> Merr.	109
<i>Flacourtia ramontchi</i> L' Herit .	109
<i>Flemingia tuberosa</i> Dalz	104
<i>Fleurya interrypta</i> Gaud	90, 113
<i>Fluggea leucophyrus</i> (Koen) Willd . .	88, 118
<i>Fluggea microcarpa</i> Blume	88, 118
<i>Fluggea virosa</i> (Roxb ex Willd) Baill	88, 118
<i>Foeniculum capillaceum</i> Gilb	104
<i>Foeniculum panmorium</i> DC.	193
<i>Foeniculum vulgare</i> Mill. .	38, 104
<i>Fritillaria cirrhosa</i> Don Prodr .	98
<i>Fritillaria imperialis</i> Linn. . .	92
<i>Fritillaria roylei</i> Hook	98, 193
<i>Fumaria officinalis</i> Linn.	193

G

<i>Garcinia indica</i> Chois. .	193
<i>Garcinia mangostana</i> Linn. .	104, 194
<i>Garcinia morella</i> Desr.	66, 111, 104
<i>Gardenia campanulata</i> Roxb. . .	118
<i>Gastrochilus pandurata</i> Ridley. . .	104

	PAGE
<i>Gaultheria fragrantissima</i> Wall.	38, 79, 118, 194
<i>Gaultheria procumbens</i> .	194
<i>Gentiana kurroo</i> Royle. . .	14, 194
<i>Gentiana lutea</i>	14
<i>Geranium macrorrhizum</i> Linn .	194
<i>Geranium robertianum</i> Linn	98
<i>Ginkgo biloba</i> Linn .	113
<i>Girardinia heterophylla</i> Decne	113
<i>Girardinia leschenaultiana</i> Decne	90
<i>Girardinia zeylanica</i> Decne .	90
<i>Gloriosa superba</i> Linn	92, 111, 118, 194
<i>Glycosmis pentaphylla</i> Correa	194
<i>Glycyrrhiza glabra</i> Linn	39, 98, 195
<i>Gmelina arborea</i> Linn	98
<i>Gnaphalium luteo-album</i> Linn .	78
<i>Gnetaceae</i> .	90
<i>Gnetum scandens</i> Roxb	118
<i>Gossypium</i> species	66
<i>Gossypium arboreum</i> Linn	98
<i>Gossypium herbaceum</i> Linn	104, 111, 195
<i>Gramineae</i>	94
<i>Graptophyllum pictum</i> (L) Griff	98
<i>Grewia asiatica</i> Linn	98
<i>Grewia hirsuta</i> Vahl. Symb	104, 109
<i>Grewia microcos</i> Linn . .	104, 110
<i>Gymnema aurantiacum</i> Wall. .	195
<i>Gymnema sylvestre</i> R Br	39, 195
<i>Gynandropsis gynandra</i> (Linn) Merr .	65, 118
<i>Gynandropsis pentaphylla</i> DC	65, 118
<i>Gynocardia odorata</i> R. Br.	65, 118

H

<i>Hagenia abyssinica</i> J F Gmel .	195
<i>Haloxylon recurvum</i> Benge ex. Boiss . .	85
<i>Haloxylon salicornicum</i> Bunge. ex Boiss.	85
<i>Hardwickia pinnata</i> Roxb . .	195
<i>Harpullia cupanioides</i> Roxb. . .	69, 118
<i>Hauclea sessilifolia</i> Roxb . .	98
<i>Hedera helix</i> Linn. . .	76, 113, 119
<i>Hedyotis auricularia</i> Linn. . .	42
<i>Helianthus annuus</i> Linn. . .	195

	PAGE
<i>Helicteres isora</i> Linn	39, 105
<i>Heliotropium eichwaldii</i> Steud	82
<i>Heliotropium indicum</i> Linn.	82
<i>Helminthosporium tetramera</i> McKinney.	195
<i>Helvella crispa</i> Fr	55
<i>Hemidesmus indicus</i> R Br.	39, 105, 196
<i>Heracleum nepalense</i> D Don	196
<i>Herpestis monniera</i> (Linn) H B & K	27, 196
<i>Hibiscus cannabinus</i> Linn	105, 196
<i>Hibiscus esculentus</i> Linn	134, 196
<i>Hibiscus sabdariffa</i> Linn	196, 197
<i>Hibiscus syriacus</i> Linn	197
<i>Hibiscus vitifolius</i> Linn	197
<i>Hippomane mancinella</i> Linn.	113
<i>Hippocratea indica</i> Willd	45
<i>Hirneola polytricha</i> Mont	55
<i>Holarrhena febrifuga</i> Klotzsch	40, 199
<i>Holarrhena mitis</i> R Br	105
<i>Holarrhena antidysenterica</i> Wall	19, 39, 80, 105, 197, 198, 199
<i>Holigarna arnotiana</i> Hook	69, 113, 199
<i>Holigarna ferruginea</i> March	69
<i>Holigarna grahamii</i> (Wight) Hook	69, 113
<i>Holigarna longifolia</i> Buch Ham ex Roxb	69, 114
<i>Homalomena rubescens</i> Kunth	93
<i>Humulus lupulus</i> Linn	114
<i>Hura crepitans</i> Linn	89
<i>Hydnocarpus anthelmintica</i> Pierre	40, 200
<i>Hydnocarpus Gaertn</i>	199
<i>Hydnocarpus kurzii</i> (King) Warb	65, 118
<i>Hydnocarpus laurifolia</i> (Dennst) Sleumer	65, 118
<i>Hydnocarpus wightiana</i> Brume	40, 65, 118, 200
<i>Hydnum caralloides</i> Scop	55
<i>Hydnum repandum</i> Linn	56
<i>Hydrocotyle asiatica</i> Linn.	32, 76, 103, 200
<i>Hydrocotyle javanica</i> Thunb	76, 105, 118
<i>Hygrophila spinosa</i> T Anders	27, 200
<i>Hyoscyamus</i> Linn	201
<i>Hyoscyamus muticus</i> Linn	83, 255
<i>Hyoscyamus niger</i> Linn	40, 83, 201
<i>Hyoscyamus pusillus</i> Linn.	83
<i>Hyoscyamus reticulatus</i> Linn	83
<i>Hypericaceae</i>	66

	PAGE
<i>Hypericum japonicum</i> Thunb . .	105
<i>Hypericum perforatum</i> Linn .	66, 114, 201
<i>Hypholoma fasciculare</i> (Huds) Fr	54, 57
<i>Hyptis suaveolens</i> Poit. . .	201
I	
<i>Illex vomitoria</i> (Soland) Hort .	201
<i>Illicium anisatum</i> Linn. .	63
<i>Illicium griffithii</i> Hk f & T .	63
<i>Illicium parviflorum</i> Michx .	201
<i>Illicium religiosum</i> Sieb & Zucc. .	63, 201
<i>Illicium verum</i> Hook f . .	63
<i>Indigofera enneaphylla</i> Linn . .	201
<i>Indigofera linifolia</i> Retz . .	201
<i>Indigofera oblongifolia</i> Forsk . .	105
<i>Indigofera tinctoria</i> Linn .	98
<i>Inula graveolens</i> Desf .	78
<i>Inula helenium</i> Hk f & T .	201
<i>Inula royleana</i> DC .	202
<i>Ipomoea</i> Linn .	82, 83, 202
<i>Ipomoea aquatica</i> Forsk .	82
<i>Ipomoea hederacea</i> Jacq .	40, 82, 202
<i>Ipomoea muricata</i> Jacq .	82, 116, 202
<i>Ipomoea pescaprae</i> Roth .	202
<i>Ipomoea pulchella</i> Roth .	202
<i>Ipomoea purga</i> Heyne .	82
<i>Ipomoea reptans</i> (Linn) Poir .	82, 102
<i>Ipomoea turpethum</i> R Br .	40, 83
Iridaceae .	90
<i>Ixora nigricans</i> Br. .	105
J	
<i>Jasminum</i> Linn. .	202
<i>Jasminum auriculatum</i> Vahel .	98
<i>Jateorrhiza palmata</i> (Lam) Miers . .	105, 203
<i>Jatropha curcas</i> Linn .	89, 105, 203
<i>Jatropha glandulifera</i> Roxb .	89, 203, 255
<i>Jatropha gossypifolia</i> Linn. .	89
<i>Jatropha multifida</i> Linn. . .	89
Juglandaceae . .	90
<i>Juglans regia</i> Linn. . .	90, 105, 118, 203
Juncaceae . .	92

		PAGE
<i>Juncus effusus</i> Linn.	..	92
<i>Juniperus communis</i> Linn	..	40, 203
<i>Jussiaea suffruticosa</i> Linn	..	105
<i>Justicia gendarussa</i> Burm	..	105
K		
<i>Kaempferia galanga</i> Linn		303
<i>Kalanchoe spathulata</i> (Poir.) DC	..	110, 118
L		
<i>Labeo rohita</i>	..	248
Labiatae	..	85
<i>Lactarius deliciosus</i> Fr	..	56
<i>Lactuca tatarica</i> C A Meyer var. <i>tibetica</i> C B Clarke.		78
<i>Lactarius vellerens</i> Fr.	.	54, 57
<i>Lactuca scariola</i> Linn	..	98, 204
<i>Lagenandra ovata</i> (Linn) Thw.	..	93, 118
<i>Lagenandra toxicaria</i> Dalz	..	93, 118
<i>Lagenaria siceraria</i> Standl	.	75
<i>Lagenaria vulgaris</i> Ser	..	75, 204
<i>Lagerstroemia flosreginae</i> Retz	..	74
<i>Lagerstroemia indica</i> Linn	..	74
<i>Lagerstroemia speciosa</i> (Linn) Pers		74
<i>Lamium amplexicaule</i> Linn	.	85
<i>Lansium anamalayanum</i> Bedd	.	204
<i>Lansium domesticum</i> Correa		204
<i>Lantana aculeata</i> Linn.		85
<i>Lantana camara</i> Linn	..	85, 204
<i>Laportea crenulata</i> Gaud	..	90, 114
<i>Laportea terminalis</i> Wight	..	90, 114
<i>Lasiosiphon eriocephalus</i> Decne	.	87, 114, 118
<i>Lathyrus aphaca</i> Linn	.	71
<i>Lathyrus sativus</i> Linn.		61, 71, 204
Lauraceae	..	87
<i>Laurus camphora</i> Linn.	..	205
<i>Lavendula burmanni</i> Benth.	..	205
<i>Lawsonia inermis</i> Roxb.	..	205
<i>Leea aequata</i> Linn.	..	99
<i>Leea indica</i> Merrill.	..	105
Leguminosae.	..	70, 205
<i>Lens esculenta</i> Moench.	..	105, 205
<i>Lens culinaris</i> Medic.	..	105
<i>Lentinus subnudus</i> Berk.	..	56

	PAGE
Leonurus cardiaca Linn .	114
Lepidium draba Linn .	65, 116
Lepidium latifolium Linn.	205
Lepidium sativum Linn	111
Lepiota sp.	57
Lepiota albuminosa Berk.	55
Lepiota cristta Quel	55, 57
Lepiota mastoidea Fr	56
Lepiota procera (Scop) Sacc	56
Leucaena glauca Benth.	205
Leucas aspera Spreng	205
Leshmania tropica .	29
Liliaceae	91
Linaceae	67
Linostoma decandrum Wall .	118
Linum usitatissimum Linn	61, 67, 99, 205
Litsaea Lamk. .	205
Litsaea chinensis Lam.	99, 205
Litsaea zeylanica Nees .	206
Lobelia Linn	206
Lobelia excelsa Leschen.	78, 114
Lobelia inflata Linn . .	206
Lobelia nicotianifolia Heyne .	78, 114, 206
Lobelia pyramidalis Vahl .	206
Lochnera pusilla K Schum	81
Lochnera rosea (Linn) Reichb .	81
Lodoicea maldivica Pers . .	110
Lodoicea seychellarum Labill	110
Loganiaceae	82
Lolium perenne Linn .	94
Lolium temulentum Linn	54, 94, 95, 906
Loranthaceae	87
Loranthus asper Lour. . .	99
Loranthus longiflorus Den	206
Luffa acutangula Roxb. .	40, 75, 99, 105
Luffa aegyptiaca Mill & Hook f. .	75, 206
Luffa amara Roxb . .	206
Luffa cylindrica Linn . .	207
Luffa echinata Roxb . .	75, 29, 207
Luffa graveolens Roxb. .	207
Lupinus (Tourn.) Linn. . .	207
Lupinus albus Linn. . .	207
Luvunga scandens Ham. .	40, 99, 207

	PAGE
<i>Lycium barbarum</i> Linn .	83
<i>Lycopodon</i> sp	56
<i>Lycopodium clavatum</i> Linn	207
<i>Lycopodium phlegmaria</i> Linn .	207
<i>Lythraceae</i> .	74

M

<i>Maba nigrescens</i> Dalz . .	208
<i>Machilus macrantha</i> Nees .	99
<i>Madhuca longifolia</i> (Linn.) Macbride	41, 80, 119
<i>Madhuca latifolia</i> (Roxb) Macbride	41, 118
<i>Madhuca indica</i> J F Gmel	41, 80
<i>Maesa indica</i> Wall	80, 119
<i>Magnoliaceae</i> .	63
<i>Mahonia acanthifolia</i> Don	208
<i>Mahonia nepalensis</i> DC .	41
<i>Mahonia</i> Nutt .	208
<i>Malanogaster durissiums</i> Cooke. . .	56
<i>Mallotus philippinensis</i> Muell -Arg	41, 208, 209
<i>Malvaceae</i> .	66
<i>Malva arborea</i> Webb . .	209
<i>Malva parviflora</i> Linn	67
<i>Malva rotundifolia</i> Linn	105
<i>Malva sylvestris</i> Linn . .	209
<i>Mandragora caulescens</i> Clarke .	83
<i>Mangifera</i> Linn . .	209
<i>Mangifera indica</i> Linn .	105, 209
<i>Manihot esculenta</i> Crantz	89, 209
<i>Manihot utilisima</i> Pohl.	89, 209
<i>Mansonia dipikae</i> C S. P	209
<i>Marrubium vulgare</i> Linn .	110, 210
<i>Martynia annua</i> Linn. .	99
<i>Martynia diandra</i> Glox. . .	210
<i>Matricaria chamomilla</i> Linn .	210
<i>Matricaria</i> Linn. .	210
<i>Meconopsis aculeata</i> Royle .	64
<i>Meconopsis napaulensis</i> DC	64
<i>Melaleuca leucadendron</i> Linn.	73, 119
<i>Melanogaster durissiums</i> Cooke .	56
<i>Melastoma malabathricum</i> Linn. . .	105
<i>Meliaceae</i> .	68
<i>Melianthus major</i> Linn. .	69

	PAGE
<i>Melia azadirachta</i> Linn ..	97, 116, 148
<i>Melia azedarach</i> Linn. ..	41, 68, 210, 255
<i>Melilotus alba</i> Desr ..	71
<i>Melilotus parviflora</i> Desf. .	210
<i>Melissa parviflora</i> Benth. .	99
<i>Melodinum monogynus</i> Roxb. .	81, 119, 255
<i>Menispermaceae</i> .	64
<i>Mentha arvensis</i> Linn. ..	42, 255
<i>Mentha aquatica</i> Linn. .	210
<i>Mentha longifolia</i> Hunds ..	105
<i>Mentha sylvestris</i> Linn. ..	105
<i>Mentha viridis</i> Linn. .	211
<i>Mercurialis annua</i> Linn. ..	211
<i>Mesua ferra</i> Linn. .	105, 211
<i>Microcystis flos-aquae</i> (Witttr) Kirch .	53
<i>Millettia auriculata</i> Baker ex Brand. .	71, 119
<i>Millettia pachycarpa</i> Benth .	71, 119, 211
<i>Millettia piscidia</i> Wight & Arn ..	71, 119
<i>Mimosa pudica</i> Linn ..	106, 211
<i>Mimusops hexandra</i> Roxb. ..	99
<i>Mimusops elengi</i> Linn .	106
<i>Mirabilis jalapa</i> Linn. .	211
<i>Momordica balsamina</i> Linn ..	75
<i>Momordica charantia</i> Linn .	75, 110, 111, 211
<i>Momordica cymbalaria</i> Fenzl ex Naud .	75, 111
<i>Momordica dioica</i> Roxb ..	99, 211
<i>Momordica tuberosa</i> Cogn ..	75, 111
<i>Monniera cuneifolia</i> Michx. .	211
<i>Morchella esculenta</i> Pers .	56
<i>Morinda citrifolia</i> Linn ..	106, 212
<i>Morinda umbellata</i> Linn. .	106
<i>Moringaceae.</i> .	70, 212
<i>Moringa concanensis</i> Nimm. .	212
<i>Moringa oleifera</i> Lamk .	42, 70, 99, 111, 134, 212
<i>Moringa pterygosperma</i> Gaertn. .	42, 70, 111, 212
<i>Morus indica</i> Linn. ..	213
<i>Morus nigra</i> Linn. .	213
<i>Mucuna atropurpurea</i> DC. ..	114
<i>Mucuna gigantea</i> DC ..	114
<i>Mucuna hirsuta</i> Wight & Arn. ..	114
<i>Mucuna monosperma</i> DC ..	114
<i>Mucuna pruriens</i> DC. ..	114, 213
<i>Mucuna prurita</i> Hook. .	106

	PAGE
Mucor sp. .	54
Mundulea sericea (Willd) Greenway. .	119
Mundulea suberosa Benth .	71, 119, 213
Murraya exotica Linn	213
Murraya koenigii Spreng	106, 213
Murraya paniculata Jack	106
Musa sapientum O Kuntze	99, 106, 110
Musa paradisiaca Linn var sapientum Kuntze.	110
Musua ferreae	255
Mylabris macilenta.	248
Myricaceae.	90
Myrica nagi Thunb .	90, 106, 119
Myristicaceae.	86
Myristica fragrans Houth. .	42, 87, 213
Myristica malabarica Lamk . .	87
Myrsinaceae . .	79
Myrsine africana Linn .	42, 213
Myrtaceae .	73
Myrtus communis Linn .	106
N	
Nannorhops ritchieana H Wendl .	106
Narcissus tazetta Linn . .	91, 213
Nardostachys jatamansi DC	214
Nepeta elliptica Royle ex Benth . .	106
Nerium indicum Mill	81, 111
Nerium odorum Soland .	81, 111, 214
Nerium oleander Linn	114, 214
Nicandra physaloides Gaertn. . .	83, 119
Nicotiana Linn .	214
Nicotiana glauca Grab. . .	214
Nicotiana rustica Linn . .	83, 119, 214
Nicotiana tabacum Linn. .	83, 119, 214, 215
Nigella sativa Linn .	63, 112, 119
Nodularia sp .	53
Nuphar luteum Sibth. & Sm . .	215
Nymphaea (Tourn) Linn .	215
Nymphaea alba Linn .	106, 215
Nymphaea lotus Linn . .	215
Nymphaea pubescens Willd. .	106
O	
Ochna pumila Ham ex D Don. .	99
Ocimum Linn .	215

	PAGE
<i>Ocimum americanum</i> Linn . .	106
<i>Ocimum basilicum</i> Linn.	106, 203, 215
<i>Ocimum canum</i> Sims .	106, 203, 215
<i>Ocimum gratissimum</i> Linn .	119, 215
<i>Ocimum kilimandscharicum</i> Guerke. .	216, 255
<i>Ocimum lemon</i> .	203
<i>Ocimum sanctum</i> Linn . .	203, 216
<i>Odina wodier</i> Roxb .	216
<i>Oldenlandia auricularia</i> K. Schum .	42, 110, 106
<i>Oldenlandia biflora</i> Hk. f .	216
<i>Oldenlandia umbellata</i> Linn. . .	99
<i>Olea</i> Linn. .	216
<i>Ononis spinosa</i> Linn .	216
<i>Operculina turpethum</i> (Linn) Manso .	83, 99
<i>Origanum vulgare</i> Linn . .	216
<i>Oroxylum indicum</i> Vent .	106, 216, 217
<i>Orthosiphon stamineus</i> Benth .	217
<i>Oscillatoria</i> . .	53
<i>Ougenia dalbergioides</i> Benth . .	71, 119
<i>Oxalis corniculata</i> Linn . .	106
<i>Oxytropis thomsoni</i> Benth .	217

P

<i>Paeonia emodi</i> Wall. .	63
<i>Pachygone ovata</i> (Poir.) Miers .	64, 119
Palmaceae. .	92
<i>Pandanus odoratissimus</i> Roxb .	217
<i>Panicum maximum</i> Jacq. .	94
Papaveraceae. .	64
<i>Papaver dubium</i> Linn. .	64
<i>Papaver nudicaule</i> Linn. .	64
<i>Papaver rhoeas</i> Linn. .	64
<i>Papaver somniferum</i> Linn. . .	42, 64, 217
<i>Paris polyphylla</i> Sns. . .	217
<i>Parmelia</i> sp . .	57
<i>Parmelia kamtschadalis</i> . . .	256
<i>Parmelia molliuscula</i> . . .	57
<i>Paspalum scrobiculatum</i> Linn. . .	95
Passifloraceae. . .	74
<i>Pavonia odorata</i> Willd. . .	107
Pedaliaceae. . .	84

	PAGE
<i>Peganum harmala</i> Linn	43, 67, 112, 119, 218
<i>Penicillium ciavariaeformis</i> Solms Laub	218
<i>Penicillium notatum</i> .	54
<i>Pentapetes phoenicea</i> Linn	125, 218
<i>Peranospora parasitica</i> .	218
<i>Perowskia atriplicifolia</i> Benth	218
<i>Peucedanum graveolens</i> Benth.	43, 107
<i>Phalaris minor</i> Retz.	218
<i>Phaseolus aconitifolius</i> Jacq	99, 108
<i>Phaseolus lunatus</i> Linn	72
<i>Phaseolus radiatus</i> Linn.	218
<i>Phaseolus trilobus</i> Art	99, 218
<i>Phebalium argenteum</i> Sm.	219
<i>Phyllanthus emblica</i> Linn	107, 219
<i>Phyllanthus niruri</i> Linn	107, 219
<i>Phyllanthus urinaria</i> Linn	89, 107, 119
<i>Physochlaima praelata</i> (Don) Miers	83, 219
<i>Phytolaccaceae</i>	85
<i>Phytolacca acinosa</i> Hook f B I non-Roxb	86
<i>Phytolacca latbenia</i> (Buch Ham) H Walt	86
<i>Picrasma excelsa</i> Swartz	14, 219
<i>Picrasma javanica</i> Blume var <i>nepalensis</i> Badhwar nov comb	119
<i>Picrasma napalensis</i> Benn.	68, 119
<i>Picrasma quassioides</i> .	14
<i>Picroena quassioides</i> Benth	43
<i>Picrorhiza kurroa</i> Royle ex Benth	43, 219, 220
<i>Pieris ovalifolia</i> D Don	78, 119
<i>Pimpinella anisum</i> Linn	43
<i>Pimpinella saxifraga</i> Linn	220
<i>Pinus</i> Linn	220
<i>Pinus excelsa</i> Walk	220, 221
<i>Pinus gerardiana</i> Wall	221
<i>Pinus khasya</i> Royle	221
<i>Pinus longifolia</i> Roxb	43, 99, 221, 223
<i>Pinus roxburghii</i> Sargent	43, 99
<i>Piper</i> sp	86
<i>Piperaceae</i> .	86
<i>Piper betle</i> Linn.	44, 86, 222, 254
<i>Piper chaba</i> Hunter.	99
<i>Piper cubeba</i> Linn.	44, 222
<i>Piper longum</i> Linn.	99, 107

	PAGE
<i>Piper nigrum</i> Linn. .	86, 107, 222
<i>Pistacia integerrima</i> Stew	44, 99, 107, 222, 223, 254
<i>Pistacia terebinthus</i> Linn .	223
<i>Pistia stratiotes</i> Linn	99, 107
<i>Pithecelobium bigeminum</i> Benth	72, 119
<i>Pithecelobium saman</i> Benth .	223
<i>Plantago ciliata</i> Desf	107
<i>Plantago major</i> Linn .	99, 107, 223
<i>Plantago ovata</i> Forsk	19, 44, 107, 323
<i>Plantanus orientalis</i> Linn	197
<i>Plesmonium margaritiferum</i> Schott	93
<i>Pleurotus cretaceus</i> Massee	56
<i>Pleurotus fimbriatus</i> Bolt	56
Plumbaginaceae	79
<i>Plumbago indica</i> Linn	44, 76, 112
<i>Plumbago rosea</i> Linn	44, 79, 112, 223
<i>Plumbago zeylanica</i> Linn .	45, 79, 99, 197, 112, 223
<i>Plumeria acuminata</i> Ait	81, 112
<i>Plumeria acutifolia</i> Poir	81, 112, 223
<i>Plumeria rubra</i> Linn var <i>acutifolia</i> Bailey	112
<i>Podophyllum</i> Linn	223
<i>Podophyllum emodi</i> Wall ex Hook f. & Thoms	64, 114
<i>Podophyllum hexandrum</i> Royle	45, 64, 114
<i>Podophyllum paltatum</i> .	45
<i>Pogostemon heyneanus</i> Benth . .	85, 119
<i>Pogostemon patchouli</i> Hook f . .	85, 119
<i>Poinciana pulcherima</i> Linn . .	110
<i>Polianthese tuberosa</i> Linn . .	256
<i>Polygala chinensis</i> Linn .	66, 223
<i>Polygala crotalaroides</i> Buch Ham . .	66
Polygalaceae. . .	66
<i>Polygala telephioides</i> Willd. . .	66
Polygonaceae . .	86
<i>Polygonum</i> Linn. . .	224
<i>Polygonum aviculare</i> Linn . .	86
<i>Polygonum flaccidum</i> Meissn. .	86, 119
<i>Polygonum hydropiper</i> Linn. . .	86, 114, 119
<i>Polygonum orientale</i> Linn. . .	86
<i>Polygonum persicaria</i> Linn . .	86
<i>Polygonum tomentosum</i> Willd . .	86
<i>Polyporus squamosus</i> (Huds) Fr . .	56
<i>Polytoca barbata</i> Stapf. .	97
<i>Pongamia glabra</i> Vent. . .	45, 72, 119, 224

		PAGE
<i>Pongamia pinnata</i> (Linn) Merr	..	45, 72, 119
<i>Potentilla anserina</i> Linn	.	224
<i>Potentilla argentea</i> Linn	.	224
<i>Prangos pabularia</i> Lindl.	..	224
<i>Premna integrifolia</i> Linn.	.	45, 224
Primulaceae	.	79
<i>Primula reticulata</i> Wall.	..	79
<i>Prinsepia utilis</i> Royle.	.	234
<i>Pristimera indica</i> Willd.	.	45, 224
<i>Prosopis juliflora</i> DC	.	224
<i>Prunus acuminata</i> Hk f	.	225
<i>Prunus amygdalus</i> Batsch	.	72
<i>Prunus armeniaca</i> Linn	.	72
<i>Prunus avium</i> Linn		72, 225
<i>Prunus cerasus</i> Linn.	..	72, 225
<i>Prunus mahaleb</i> Linn.		72
<i>Prunus nepalensis</i> Koch.	.	225
<i>Prunus padus</i> Linn	.	72
<i>Prunus persica</i> Stokes	.	72
<i>Prunus puddum</i> Roxb	.	72, 225
<i>Prunus undulata</i> Buch Ham	.	72
<i>Psalliotia xanthoderma</i>		57
<i>Psidium guajava</i> Linn	.	107, 110, 225
<i>Psilotum triquetrum</i> Sw	.	225
<i>Psoralea corylifolia</i> Linn.	.	19, 45, 225, 226
<i>Psychotria ipecacuanha</i> Stokes	..	32
Pteridophyta (Vascular Cryptogams)		57
<i>Pteris aquilina</i> Linn	.	58, 226
<i>Pterocarpus marsupium</i> Roxb	..	226
<i>Pueraria thunbergiana</i> Benth	..	226
<i>Pulicaria dysenterica</i> Gaertn.	..	107
<i>Punica granatum</i> Linn	..	107, 226
<i>Putranjiva roxburghii</i> Wall	..	226
<i>Pygeum gardneri</i> Hook f	.	72, 19
<i>Pyrethrum</i>	..	168, 18
<i>Pyrus aucuparia</i> Linn.	.	72
<i>Pyrus malus</i> Linn	..	72

Q

<i>Quercus</i> Linn.	..	266
<i>Quercus dilatata</i> Lindl.	..	227
<i>Quercus ilex</i> Linn	..	227

	PAGE
<i>Quercus incana</i> Roxb.	227
<i>Quisqualis indica</i> Linn	227
R	
<i>Randia dumetorum</i> Lamk	46, 77, 112, 119, 227
<i>Randia uliginosa</i> DC.	77, 119
Ranunculaceae	62
<i>Ranunculus arvensis</i> Linn	63
<i>Ranunculus falcatus</i> Linn.	63
<i>Ranunculus laetus</i> Wall	63
<i>Ranunculus lingua</i> Linn	63
<i>Ranunculus pensylvanicus</i> Linn f.	63
<i>Ranunculus sceleratus</i> Linn.	63, 114
<i>Rauwolfia</i> Linn.	227
<i>Rauwolfia canescens</i> Linn.	46, 227, 228, 256
<i>Rauwolfia micrantha</i>	262
<i>Rauwolfia serpentina</i> Benth	19, 46, 120, 228, 229, 230, 231, 255, 257, 258
<i>Revenelia sessilis</i>	231
<i>Rhamnus</i> Linn.	231
<i>Rhamnus virgatus</i> Roxb	231
<i>Rheum</i> Linn	231
<i>Rheum emodi</i> Wall	46, 86, 107, 231, 232
<i>Rhizodonia solani</i> Kuhn	216
<i>Rhododendron arboreum</i> Sm	79
<i>Rhododendron anthopogon</i> D Don.	79
<i>Rhododendron barbatum</i> Wall ex G Don	79, 120
<i>Rhododendron campanulatum</i> D Don...	79, 100
<i>Rhododendron cinnabarinum</i> Hook f	79
<i>Rhododendron falconeri</i> Hook f.	120
<i>Rhododendron setosum</i> D. Don	79
<i>Rhus insignis</i> Hook. f.	69, 114
<i>Rhus punjabensis</i> J L. Stew. ex Brand.	69, 114
<i>Rhus succedanea</i> Linn	69, 100, 114
<i>Rhus wallichii</i> Hook. f.	69, 114
<i>Ricinus</i> Linn.	232
<i>Ricinus communis</i> Linn.	46, 89, 120, 232
<i>Robinia uliginosa</i> Roxb ex Willd	117
<i>Rosa</i> Linn.	232
<i>Rosa damascena</i> Mill	47
<i>Rourea santaloides</i> W. & A.	100
Rubiaceae.	77
<i>Rubia cordifolia</i> Linn.	107

		PAGE
Rubus moluccanus Gaertn	..	73, 112
Rumex Linn	..	232
Rumex acetosa Linn.	.	86, 114
Rumex acetosella Linn	..	86, 114
Rumex scutatus Linn.	..	107
Rumex vasicarius Linn	.	107
Rutaceae	..	67
Ruta graveolens Linn var angustifolia Hk f	..	67, 112, 114, 120, 232
Ruta tuberculata Forsk.	..	67

S

Saccharum spontaneum Linn	..	233
Salicornia brachiata Roxb	..	85, 112
Salix Linn	..	233
Salix alba Linn		107
Salsola kali Linn.	..	85
Salvadoraceae	..	80
Salvadora oleoides Dcne	.	80, 233
Salvadora persica Linn.	.	80
Sambucus Linn	.	233
Sambucus ebulus Linn	..	77
Sambucus nigra Linn.	.	77
Samydaceae	..	74
Sansevieria roxburghiana Schult	..	100
Sansevieria zeylanica Willd.	..	233
Santaloides minus Schellenb.	..	100
Santalum album Linn.	..	47, 120, 233, 134
Sapindaceae	..	69
Sapindus Plum	..	234
Sapindus laurifolius Vahl.	..	234
Sapindus mukorossi Gaertn.	..	69, 120, 234
Sapindus trifoliatus Linn.	..	69, 100, 107, 110, 112, 120
Sapium indicum Willd.	..	89, 120
Sapium insigne Trimen.	..	89, 114
Saponaria Linn.	..	234
Saponaria vaccaria Linn.	..	66
Sapotaceae	..	80
Sphaeranthus indicus Linn.	..	100
Saraca indica Linn.	..	47, 107, 234
Sarcococca pruniformis Lindl.	..	234
Sarcostema acidum (Roxb.) Voigt.	..	82, 120
Sarcostema brevistigma W. & A.	..	82, 120

	PAGE
<i>Sauromatum guttatum</i> Schott. . .	93
<i>Saussurea lappa</i> Clarke .	47, 78, 120, 234, 235
<i>Sachima wallichii</i> Choisy. . .	115
<i>Schleichera oleosa</i> (Lour) Merr. . .	69, 110, 120
<i>Schleichera trijuga</i> Willd . . .	69, 110, 120
<i>Schweinfurthia spherocarpa</i> Br. .	110
<i>Scilla indica</i> Baker. . .	14, 47, 92, 263
<i>Scilla maritima</i> .	14
<i>Scindapus officinalis</i> Schott. .	107
<i>Scirpus corymbosus</i> Heyne .	9
<i>Scirpus grossus</i> Linn .	235
<i>Scleria pergracilis</i> (Nees) Kunth. .	120
<i>Scopolia</i> Jacq .	235
<i>Scopolia anomala</i> (Linn et Otto) Airy-Shaw. .	83
<i>Scopolia lurida</i> Dunal .	83
Scrophulariaceae .	84
<i>Secale cereale</i> Linn .	235
<i>Secamone emetica</i> R Br. .	82
<i>Securigera securidaca</i> Linn. .	235
<i>Semecarpus anacardium</i> Linn. .	447, 70, 108, 112, 115, 235
<i>Semecarpus travancoricus</i> Bedd. .	70, 115, 236
<i>Senecio</i> sp .	78
<i>Senecio jacobaea</i> Linn .	236
<i>Senecio retrorsus</i> DC .	236
<i>Senecio vulgaris</i> Linn. .	78, 236
<i>Sesbania aegyptiaca</i> Pers . .	100
<i>Sesbania sesban</i> (Linn.) Merr. . .	100
<i>Sesamum indicum</i> Linn .	84, 112
<i>Sesamum orientale</i> Linn .	84, 112
<i>Seseli indicum</i> Wight & Arn. .	236
<i>Setaria palmaefolia</i> Stapf .	236
<i>Shorea robusta</i> Gaertn f. .	108
<i>Sida cordifolia</i> Linn .	48, 100, 236
<i>Sida rhombifolia</i> Linn .	67, 100
<i>Sida spinosa</i> Linn. .	108
Simaroubaceae .	68
<i>Sinapis alba</i> Linn. .	65
<i>Sisymbrium irio</i> Linn. .	236
<i>Skimmia laureola</i> Sieb. & Zucc. ex Walp .	48, 68, 236
<i>Smilax prolifera</i> Roxb. . .	108
Solanaceae. . .	83
<i>Solanum coagulans</i> Forsk. . .	84
<i>Solanum dulcomara</i> Linn. . .	84

	PAGE
<i>Solanum incanum</i> Linn.	84
<i>Solanum indicum</i> Linn. . .	237
<i>Solanum melongena</i> Linn . .	110
<i>Solanum nigrum</i> Linn.	84, 108, 237
<i>Solanum pseudocapsicum</i> Linn .	237
<i>Solanum spirale</i> Roxb.	84
<i>Solanum trilobatum</i> Linn	100
<i>Solanum torvum</i> Sw.	237
<i>Solanum tuberosum</i> Linn	62, 84
<i>Solanum xanthocarpum</i> Schrad & Wendl	237
<i>Sonneratia acida</i> Linn	48, 237
<i>Sophora mollis</i> R Grah	72, 120
<i>Sophora tomentosa</i> Linn	72, 110
<i>Sorghum halepense</i> Pers .	61, 95
<i>Sorghum saccharatum</i> Pers	95
<i>Sorghum vulgare</i> Pers	61, 95
<i>Soyimida febrifuga</i> A Juss . .	108
<i>Spartium scoparium</i> Linn .	238
<i>Sphaeranthus indicus</i> Linn	78, 120, 238
<i>Spilanthes acmella</i> Murr .	108, 238
<i>Spiraea aruncus</i> Linn	238
<i>Spondias pinnata</i> Kurz	100
<i>Stachytarpheta jamaicensis</i> Vahl var. <i>indica</i> Lam.	85, 108, 112
<i>Stachytarpheta indica</i> Vahl. .	85, 108, 112
<i>Stelaria media</i> Linn .	238
<i>Stephania glabra</i> Roxb	48, 100, 238
<i>Stephania hernandifolia</i> (Willd.) Walp. . .	120
<i>Steudnera virosa</i> (Kunth) Prain .	93
<i>Stipa</i> sp	95
<i>Stipa siberica</i>	61
<i>Streblus asper</i> Lour	100
<i>Streptomyces auriofociens</i> .	52
<i>Streptomyces griseus</i> . .	52
<i>Streptomyces rimosus</i> .	52
<i>Streptomyces venezuelae</i>	52
<i>Stropharia semiglobata</i> (Batsch) Quel .	54, 57
<i>Strychnos colubrina</i> Linn.	82, 120
<i>Strychnos nux-vomica</i> Linn . .	48, 82, 87, 108, 110, 120, 238
<i>Suaeda fruticosa</i> Forsk.	85
<i>Sunantherias sylvatica</i> Schott. .	93
<i>Swertia chirata</i> Ham.	48, 238
<i>Swertia decussata</i> Nimmo. . .	238
<i>Swietenia macrophylla</i> Linn. . .	238, 239

	PAGE
<i>Symplocos racemosa</i> Roxb.	49, 108, 239
<i>Syzygium cumini</i> (Linn.) Skeels	108
<i>Syzygium jambos</i> (Linn) Alst.	108
<i>Syzygium operculatum</i> Gamble	108

T

<i>Tabernaemontyna coronaria</i> Br	.	239
<i>Tabernaemontana dichotoma</i> Roxb.	..	81
<i>Taccaceae</i>	.	91
<i>Tacca pinnatifida</i> Forsk.	..	91
<i>Tagetes erecta</i> Linn.	..	239
<i>Tagetes patula</i> Linn.	.	239
<i>Tamarindus indica</i> Linn	.	239, 240
<i>Tanacetum</i> Linn	.	240
<i>Taraktogenos kurzii</i> King	.	65, 118
<i>Taxus baccata</i> Linn	.	90, 112, 120, 240
<i>Taxus brevifolia</i> Nutt	..	240
<i>Tectona grandis</i> Linn f		108, 240
<i>Tephrosia</i> Pers.	.	240
<i>Tephrosia candida</i> (Roxb) DC		72, 120
<i>Tephrosia purpurea</i> (Linn.) Pers.		72, 120, 240
<i>Tephrosia villosa</i> Pers.	.	240
<i>Tephrosia vogelii</i> Hook f	.	120
<i>Teramnus labialis</i> Spreng	.	100
<i>Terminalia arjuna</i> W & A	..	49, 108, 241
<i>Terminalia bellerica</i> Roxb.		73, 120, 241
<i>Terminalia chebula</i> Retz	.	73, 108, 110, 241
<i>Terminalia citrina</i> Fleming.	.	108, 110
<i>Terminalia paniculata</i> Roth.		110
<i>Ternstroemiaceae</i>	.	66
<i>Thalictrum foliolosum</i> DC	.	94, 241
<i>Thea sinensis</i> Linn.	..	66, 241
<i>Thevetia nereifolia</i> Juss.	.	49, 81, 112, 120, 241, 242
<i>Thevetia peruviana</i> (Pers) Merr.	..	49, 81, 112, 120
<i>Thomsonia nepalensis</i> Wall.	..	94
<i>Thymeliaceae</i> .	..	87
<i>Thymus serpyllum</i> Linn.	..	242
<i>Thymus vulgaris</i> Linn.	..	242
<i>Thysanolaena maxima</i> Kuntze.	..	242
<i>Tinospora crespa</i> Miers.	..	242
<i>Tinospora cordifolia</i> Miers.	..	49, 242

	PAGE
<i>Toddalia aculeata</i> Pers	49, 242
<i>Toddalia asiatica</i> Lam.	49
<i>Trachyspermum ammi</i> (Linn) Sprague	50, 110
<i>Tragia bicolor</i> Miq.	89, 115
<i>Trapa bispinosa</i> Roxb.	242
<i>Tragia involucrata</i> Linn.	89, 115
<i>Trespasia populnea</i>	243
<i>Trianthema monogyna</i> Linn.	76, 112
<i>Trianthema pentandra</i> Linn.	76, 112
<i>Trianthema portulacastrum</i> Linn.	76, 112, 243
<i>Tribulus terrestris</i> Linn.	50, 67, 243
<i>Trichosanthes anguina</i> Linn.	243
<i>Trichosanthes bracteata</i> Voigt.	75
<i>Trichosanthes cucumerina</i> Linn	75
<i>Trichosanthes dioica</i> Roxb.	75, 243
<i>Trichosanthes palmata</i> Roxb.	75
<i>Trifolium repens</i> Linn	72
<i>Trigonella foenum-graecum</i> Linn	120, 243
<i>Triticum</i> Linn	242
<i>Triticum aestivum</i> Linn.	95
<i>Triumpheta bartramia</i> Linn	108
<i>Triumfetta rhombodea</i> Jacq	243
<i>Tussilago farfara</i> Linn	244
<i>Tylophora asthmatica</i> W. & A	50, 82, 108, 244
<i>Tylophora fasciculata</i> Buch	82
<i>Tylophora indica</i> (Burm f) Merr.	50, 82
<i>Typha elephantina</i> Roxb	108
<i>Typhonium trilobatum</i> Linn. Schott	94

U

<i>Umbelliferae</i>	76
<i>Uraria lagopoides</i> DC.	108
<i>Urena lobata</i> Linn.	112
<i>Urginia coromandeliana</i> Hook. f.	92
<i>Urginia indica</i> Kunth.	50, 92, 244
<i>Urginia scilla</i>	92
<i>Urticaceae</i>	89
<i>Urtica dioica</i> Linn	90, 100, 115
<i>Urtica hyperborea</i> Jacq.	90, 115
<i>Urtica parviflora</i> Roxb.	90, 115
<i>Urtica pilulifera</i> Linn.	90, 115

		PAGE
Uroloncha malabarica Linn	.	248
Usnea hirta		263
V		
Vaccinium Linn	.	244
Valariana Linn.	.	244
Valariana wallichii DC.	..	50
Valariana officinalis Linn	.	244
Vanda roxburghii R. Br		50, 244
Vanda spathulata Spreng	.	100
Vangueria spidosa Roxb	.	244
Vateria indica Linn	.	100, 109
Verbascum thapsus Linn	.	84, 100, 121
Verbenaceae	.	84
Verbena officinalis Linn.		85, 245
Verbena venosa	..	245
Vernonia anthelmintica Willd	.	32, 78, 116, 245
Vernonia cinerea Less	.	100
Vernonia roxburghii Less	..	100
Vetiveria zizanioides Stapf	..	245
Vicia sativa Linn	.	72
Vigna catjang Walp.	.	245
Vinca pusilla Murr.	..	81
Vinca rosea Linn		81, 246
Viola odorata Linn	..	246
Viscum Linn.	.	246
Viscum album Linn	..	87, 246
Vitex negundo Linn	..	121, 246
Vitex peduncularis Wall.	..	20, 51, 246
Volvaria diplasia Berk & Broome.	..	56
Volvaria glocephala Gill	..	55, 57
Volvaria terrestria Berk & Broome.	..	56
W		
Wallichia disticha T Anders	..	93, 115
Walsura piscidia Roxb.	..	68, 121
Wikstroemia indica (Linn) C.A. Mey.		
var viridiflora (Meissn) Hook f	..	87, 121
Withania somnifera Dunal.	..	51, 84, 100, 112
Woodfordia fruticosa Kurz	..	109
Wrightia tinctoria R. Br.	..	109, 246

X

Xanthium strumarium Linn.	..	51, 78, 115, 246
Xanthoxylum alatum.	..	254

Z

Zanonia indica Linn.	.	75
Zanthoxylum acanthopodium DC.	..	247
Zanthoxylum alatum Roxb.	..	68, 121, 247
Zanthoxylum budrunga Wall.	..	110, 247
Zanthoxylum hamiltonianum Wall	.	121
Zanthoxylum ovaltifolium Wight.	..	247
Zanthoxylum rhesa DC.	.	247
Zea mays Linn	.	95
Zingiber officinale Rosc.	..	51, 247
Zizyphus jujuba Lam	..	100
Zizyphus vulgaris Lam.	.	248
Zizyphus xylopyrus Willd.	.	248
Zygophyllaceae	..	67

INDEX

OF

VERNACULAR NAMES

	PAGE		PAGE
A		Alarī (M)	81
Aal (Bo.)	106	Alarka (S)	81, 100
Aaraar (H)	40	Alasī (Bo)	67
Ach (H & B)	106	Algusī (B.)	36, 82
Acha (M)	80	Alkaloids (English)	248
Ada (B.)	51	Allī (M)	106
Adhatodai (M)	23	Allipallī (P , H & Kash)	112
Adrak (H)	51	Alombe (Bo)	55
Adrakam (S)	51	Alpayushī (S)	92
Adulsa (H & B.)	23	Alshivīral (M)	67
Adutina-palai (M)	86	Amanakham chedī (M)	46, 89
Afim (H & B.)	42, 64	Amaravela (S)	36, 82
Agave Family (English)	91	Amaryllis Family (English)	91
Agla (Kumaon.)	70	Amb (H , B & Bo)	105
Agnigarva (S)	74	Ambarī (Bo)	105
Agnisikha (M)	92	Amku lang-kalang (M)	51, 84
Agnad (B)	120	Amla (Bo)	107
Ahiphena (S)	42, 64	Amlīka (S)	106
Ailanto (English)	101	Amlukī (B)	115
Ajakarua (S)	100	Amra (H , B & Bo)	100
Ajmod (Vern)	254	Amrataka (S)	100
Ajmoda (S. & H.)	76	Amritphala (S)	103
Ajowan (H , Bo & Vern)	50, 163	Amrud (H)	110
Ak (H)	30, 81	Amrul (H & B)	106
Akanda (B & Bo.)	30, 81	Amsania (Vern)	37
Akanadī (Vern.)	33	Amudanda (P)	41
Akar Kanta (B)	23	Amumpatchatarissī (M.)	38
Akarkara (Bo & P.)	108	Amva (S)	105
Akasbel (H)	36, 82	Anantamul (B.)	39
Akhroot (H. & B.)	90	Anannas (H)	91
Ak-Madar (Vern.)	30	Ananta (S.)	39
Akola (H)	23	Anar (H)	107
Akoria (H.)	69	Anaras (B)	91
Akroda (Bo.)	90	Anasha pazham (M.)	91
Akrottū (M.)	90	Anashuppu (M)	63
Akshota (S.)	90	Anasphal (H)	63
Aku-jemudu (M.)	88	Ankra (H.)	72
Al (Bo.)	106	Ankarī (B.)	72
Alangi (M.)	23	Ankola (Bo.)	23

	PAGE		PAGE
Ankota (S)	23	Ati vadayam (M)	22
Antamul (H. & B)	50, 82	Atibala (S)	67
Anthalmul (Bo.)	50, 82	Atimaduram (M)	39
Anthelmintics (English)	248	Atis (H & Vern.)	22
Antibiotics (English)	249	Ativisha (S)	22
Aoula (H)	107	Atmagupta (S.)	106
Aparajita (S.)	71	Atropa (English)	83
Aparajit (H & B.)	71	Aurhur (Vern)	34
Aphu (Bo)	42, 64	Avaram (Tam.)	103
Arand (H)	89		B
Archaka (B)	48	Babachi (H & B)	45
Ardana (H)	109	Babla (B)	101
Ardawal (P)	79	Bach (H & Vern.)	22, 93
Ardisia Family (English)	79	Bachata (H.)	112
Arfu (Nep)	70	Bachnag (Vern)	21
Arghavan (Arab.)	120	Bacteria (English)	52
Aripu (M.)	85	Badam (H , B & Bo)	72
Arishta (S.)	51, 78	Badbar (H.)	98
Arjun (H & B)	49	Badari (S.)	100
Arjuna (S. & Bo.)	49, 74	Badian (Bo)	63
Arka (S)	30, 81	Badrang (H)	110
Arlu (H)	106	Bael (H , B & Bo.)	23
Arni (H & Bo.)	45	Bael fruit (English)	23
Aroid Family (English)	93	Baer (H)	100
Aromatic plants (English)	249	Baghnoki (B)	99
Aru (H)	72	Bagua (H)	84
Arusha (Vern)	23	Bahera (H & B)	73
Arvada (M.)	67	Bahira (S.)	73
Asek (M)	57	Baigun (H.)	110
Aseru (Nep)	36	Bakas (B)	23
Asgandh (H)	51, 84	Bakra (H)	68
Ashvakatri (S)	109	Bakul (S , H & Bo)	106
Ashwagandha (S., B. & Bo)	51, 84	Bala (S & B)	48, 107
Asita kutanja (S)	109	Balraksha (P)	78
Asok (H , B & Bo)	47	Balunt-shep (Bo)	43
Asoka (S)	47	Banbainag (Kash)	22, 62
Aspergillus (English)	54	Banda (P.)	76
Asundro (Bo)	102	Bandurgi (Bo)	69
Aswamantaka (S)	102	Bangab (B.)	117
Ata (B)	64	Ban-gal (B.)	80
Ataicha (B.)	22	Banger chhata (B.)	55
Atasi (S)	67	Bankhur-i-Mirya (Ind Baz.)	79
Athiballachetu (M)	67	Banlaunga (H.)	105

	PAGE		PAGE
Bannatia (H.)	86	Bhat (H & B.)	34
Bans (B & H.)	27, 94	Bhela (H & B.)	47, 70
Bans kaban (H.)	94	Bhimb (Bo)	34
Bar (B)	104	Bhui-chane (Bo)	25
Bara-elachi (H. & B)	24	Bhuidari (Bo)	82
Bara-kanur (B.)	91	Bhui-dumur (B.)	104
Bara-keru (B)	38	Bhui-kanda (Bo)	47
Barbatti (Bo)	98	Bhumbi-jambu (S)	108
Bare kulinjan (H)	24	Bhumya-amalaki (S)	107
Barhanta (H)	89	Bhunimba (S)	24, 101
Barī-saunf (H)	38	Bhuta-pala (Bo)	68
Barī sophā (Bo)	38	Bhutia-Tsallu (Vern)	79
Barnagi (H)	97	Bhutta (H & B.)	95
Barola (B.)	69	Biba (Bo)	47, 70
Barphali (Vern.)	38	Bibu (B)	69
Bartaku (S)	119	Bichu (H & P)	90, 99, 100
Bartang (Bo)	99	Bichuti (B.)	89
Baru (H)	61, 95	Bignonia Family (English)	84
Barun (H)	98	Bihī (H)	103
Bas (Bo)	94	Bijindak (Afgh)	65, 116
Basak (H)	36	Bikh (Vern)	22, 62
Basant (H & P.)	66	Bilnalita (B)	103
Basingh (Bo)	109	Bililotan (H)	99
Bassu (Vern)	48	Bimba (S)	34
Bayabirang (H)	98	Bincha (B)	109
Bean-caper and Guaicum Family (English)	67	Bindimuthi (Santh.)	104
Beet Family (English)	85	Birama-dandu (M.)	25, 64
Begonia Family (English)	75	Birbanka (Nep.)	112
Behaira (Bo)	73	Birmova (Bo.)	104
Belatijau (B)	103	Birch-wort Family (English)	86
Belladonna (English)	168	Bis (Vern & P)	62, 107
Bell-flower Family (English)	78	Bish (H & B)	22, 62
Berberry Family (English)	64	Bishcopra (Kumaon)	79
Bet (H., B & Bo)	102	Bishkapra (P & Bo)	76
Betel leaf (English)	44	Bislambhi (H)	75
Bhallatamu (S)	47, 70	Bishlanguli (B)	92
Bhallava- anga (S.)	105	Biswul (H)	70
Bhang (M)	30	Bitter-bari Family (English)	68
Bhangī (M)	30, 89	Black mustard (English)	65
Bhant (H. & B.)	34	Blue gum tree (English)	38
Bharangi (Vern , S. & Bo.)	43, 97	Bobawachi (Bo)	45
Bharphand (Vern)	25	Bonmethi (B)	180

	PAGE		PAGE
Bon-nil (B.)	72	Champa (H)	66
Bon-okra (B)	51, 78, 112	Champabaha (Santh)	99
Bonpatol (B)	75	Champa Family (English)	63
Bor (H)	104	Chandla (Bo)	25, 89
Borage Family (English)	82	Chandra (B. & Bo)	46, 81
Borasali (Bo)	106	Chandrasura (S)	111
Braehimbhi (H)	27	Chansaur (H)	111
Brahmi (Vern & S)	27, 32	Chanduka (Vern)	25
Brahmo manduki (H.)	32	Chanu (B)	76
Brela (B)	48	Chapra (H)	42
Brihat-upa-kunchika (S)	24	Charayotah (H)	48
Brihmisak (B)	27	Chatium (H)	24
Brimposh (Kash)	106	Chatri (Nep)	41
Broyphyta (Livervorts & Mosses)		Chattrak (S)	55
(English)	57	Chaulmoogra Family (English)	65
Buchanaka (S)	25	Chaulmoogra (H , B & Bo)	65
Buck-wheat (English)	62	Cheraelu (H.)	79
Buck-wheat Family (English)	86	Cherailu (H)	100
Buin (P & Kash)	107	Chian (H)	71
Bulu (Nep)	79	Chikana (Bo)	48
Buna (P)	120	Chikri (Kash)	88
Bun-akora (B)	108	Chikti (H)	108
Burundi (Bo)	82	Chilla (M)	88
Buttercup Family (English)	62	Chillara (H. & Bo)	74, 116
C		Chilli (H)	74
Camphor (English)	255	Chhatim (B)	24
Caper Family (English)	65	Chhota-gokru (H)	51, 78
Carnation Family (English)	66	Chhoti dudhi (H)	88
Carrot Family (English)	76	Chiner badam (B.)	25
Casearia Family (English)	74	Chir (Vern)	43
Cashew Family (English)	69	Chiraita (Bo.)	48
Cassarva (Baz)	89	Chiraputi (Bo.)	75
Castor oil Family (English)	87	Chireta (B)	48
Castor seeds (English)	46	Chiretta (Vern.)	48
Cha (Vern.)	30	Chirghas (Kash)	82
Chagul-banti (B)	36	Chribenda (M.)	48
Chaksu (H.)	31, 71	Chiror Anrudana (Vern.)	41
Chakramards (S.)	97	Chir Pine (English)	43
Chakrani (Vern)	29	Chirpoti (H.)	75
Chakunda (H. & Bo)	97	Chirval (H)	99
Chambul (P.)	63	Chita (H.)	45, 79

	PAGE		PAGE
Chitaro (Bo.)	45, 79	Dalme (H.)	88
Chitra (Vern. & P)	28, 73	Damahan (H)	109
Chitraka (S)	44, 45, 79	Danti (S , H. & B.)	88
Chitro (Nep)	28	Dantimul (Bo)	88
Chittira (M)	45	Dar-hald (H & B)	28, 46
Chittur-mol (M)	79	Darimba (S)	107
Cholam (M)	95	Darmiya Gobriya (Nep.)	21
Chota chand (H)	46, 81	Dapoli (Bo)	110
Chota gokhru (H)	50, 67	Daru haridra (S)	28, 64
Chota pilu (H & B)	80	Daru haridraka (S)	35
Choti-elachi (H & B)	37	Darya kanariyal (H & Bo)	110
Chuka (H., B & Bo)	107	Datura Family (English)	83
Chukapalam (B)	86	Davana (Vern)	26
Chukra (S)	107	Deo-dhan (H & Bo.)	95
Chumbi Aconite (English)	128	Derris (English)	168, 185
Chumlani (Nep)	48, 67	Devadarum (M)	109
Chutrika (S)	86	Devils cotton (English)	21
Cinchona (English)	33	Dhai (H & B)	109
Clathroxystis (English)	53	Dhakangu (P)	63
Coffee (H)	77	Dhakur (B)	32, 80
Coffee Family (English)	77	Dhan (B)	95
Coffee (English)	175	Dhania (H & B)	97
Coffee powder (English)	245	Dhanyaka (S)	97
Colombo (Bo)	105	Dharakadamba (S)	77
Coriander Family (English)	76	Dhatraphala (S.)	107
Convannamilpori (M)	46	Dhatraphal (S)	73
Coriaria Family (English)	70	Dhatakı (S)	103
Cotton Family (English)	66	Dhavala (Bo.)	78, 141
Covannamilpori (M)	81	Dhingri (Vern.)	56
Croton Family (English)	87	Dhub (H , B. & Bo.)	109
Cruciferae (English)	65	Dhudı (Vern)	39
Crude Drugs (English)	249	Dhumrapatra (S)	86
Cryptogamic flora (English)	59	Dhundul (B)	75
Cubebs (English)	44	Digitalis Family (English)	84
Cucumber Family (English)	74	Dikoor (English)	94
Cultivation of Medicinal Plants (English)	249	Dirghapattra (S.)	75
Custard apple Family (English)	64	Dita-bark (English)	24
		Diva (Bo)	91
D		Dochunty (B.)	101
Dahar karanja (B.)	45, 72	Dodbi (Bo)	37
Daka (Bo)	119	Dog-bane Family (English)	80
Dalchini (Vern.)	33	Dogwood (English)	38
Dama (P.)	109	Don (P)	112

	PAGE		PAGE
Dowla (Bo.)	108	Gajar (H., B. & P)	76
Drek (Vern)	41	Gajega (M.)	30
Drug adulteration (English)	250	Ganjara kelangu (M.)	76
Drug resources (English)	250	Galot (P)	103
Drug scarcity (English)	250	Gam (Bo)	95
Dryopteris (English)	187	Gamboge Family (English)	66
Dudhi (H)	38	Gangwa (B)	88
Dudiya (B.)	88	Ganja (H , B & Bo.)	30, 89
Dudla (Bo.)	89	Ganjika (S)	30, 89
Durga chhata (B)	55	Ganjni (H)	117
E		Ganiari (B)	45
Ebony Family (English)	80	Ganikarika (S)	45
Edakula-pala (M)	24	Garbijaur (H.)	99
Ela (S)	37	Gardal (Bo)	71
Elachi (Bo)	37	Garlic (English)	23
Ellakai (M)	37	Garudu-mukku (Tel.)	99
Elam (Tam)	101	Gatta colla (Vern)	42
English-Maiden-hair-tree	113	Gaungchi (H)	21, 70
English-Manchineal tree	113	Gelaphal (Bo)	46, 77
English-Moonwort	102	Geva (B)	88
Equisetum (Horsetail) (English)	58	Geyapal (Bo)	88
Eranda (S)	46, 89	Ghet-kachu (B)	94
Erendi (Bo)	46, 89	Ghetuli (Bo.)	29
Ergot (English)	54, 173	Ghiatarui (H)	75
Ermul (B)	67, 114	Ghi kanvar (H)	24
Erukku (M.)	30, 81	Ghirna (Garhwal)	102
Essential oils (English)	250	Ghosalata (B)	75
F		Ghosi (Nep)	69
Ficoideae (English)	76	Ghosali (Bo)	75
Faughimehandi (P)	74	Ghrita Kumari (S & B.)	24
Flax Family (English)	67	Gigantic swallow root (English)	30
Foxglove (English)	37	Gilagach (B.)	71
French sorrel (English)	107	Giloe (Vern.)	49
Fungi (English)	53	Girbuti (Bo.)	83
Fusarium (English)	54, 193	Gnetum Family (English)	90
G		Goagarilakei (Bo.)	82
Gab (H. & B)	104	Gobar champa (B.)	81
Gadab (H)	114	Gobari (Nep)	21, 62
Gadadhar (S)	26, 77	Goda (B)	51
Gaggar (Kash.)	79, 100	Godhum (S.)	95
Gajapipal (H & B.)	107	Godumal (M)	95
		Gokhuri (B.)	50, 67
		Gokhru (Vern.)	50

	PAGE		PAGE
Gokshura (S)	50, 67	Gurgur (B.)	97
Golapphul (B)	47	Guttiferae (English)	66
Golappu (M)	47		
Golmarich (B)	107	H	
Golmirch (H)	107		
Gorakh-amli (H. & Bo.)	101	Haldi (H)	35
Gorakmundi (H & Bo)	78, 100	Halgach (Vern)	35
Goruma (H)	98	Haldi-gach (B.)	35
Goshtam (M.)	47, 78	Halim (B)	111
Guttibira (M.)	75	Halood (B)	35
Gowali (Bo)	109	Hansraj (Bo)	101
Grass Family (English)	94	Harara (H)	73
Grecian Foxglove (English)	36	Haridra (S)	35
Ground Nut (English)	25	Hardu (H)	77
Gubak (S)	25, 92	Harira (H)	110
Guchhian (P)	56	Haritaki (S & B)	73, 110
Gudatreka (S)	33	Harivera (S)	107
Guduchi (S)	49	Hastisunda (S)	82
Guggul (B , Bo. & H)	35, 97	Hatisura (H & B)	82
Gugul (Vern)	42	Hattut-tumatti (M)	75
Guggula-dhup (Bo)	101	Hawar (H)	84
Guggulu (S)	35	Hayer (B)	97
Gukhulu (M)	35	Hazardana (P)	88
Gul (Bo)	47	Hazarmani (H & B.)	89
Gulab ka phul (H)	47	Heart poisons (English)	251
Gulabjaman (H. & Bo)	108	Helga (B)	108
Gulancha (H & B)	49	Hemp Family (English)	89
Gulari (H)	104	Henna Family (English)	74
Gulsakari (H.)	108	Hijal (B.)	73, 102
Gulwail (Bo)	49	Hijjal (H.)	102
Gumbhari (S)	98	Hijli badam (B.)	69
Gum gugul (English)	35	Hindisana (H)	31
Gandhagatra (S.)	64	Hing (H & B)	38
Gundumani (M.)	21, 70	Hingan (H & B.)	68
Guniyan (Kashm.)	103	Hinger (Bo)	68
Gunj (P)	71	Hingu (S)	38
Gunja (S & Bo)	21, 70	Hiranpadi (H)	82
Guras (Nep)	79, 120	Hirankhori (Bo.)	103
Gurbiani (P)	49	Hiranpag (Bo)	82
Gurdli (P.)	72	Hirida (Bo.)	73
Gurguli (P.)	87	Hirtiz (Kash)	113
Gur mar (Vern.)	39	Hmyaseik (Burm.)	25, 89
Gurgunna (P.)	85	Honey-suckle Family (English)	76

	PAGE		PAGE
Horse-radish Family (English)	70	Ishappukol-virai (M)	44
Horsetail (English)	58	Isharmul (H & B.)	25, 86
Hsathanpaya (Burm)	118	Ispaghul (H.)	44
Hulugiri (Bo)	69	Ivy Family (English)	76
Hurmal (H. & Bo.)	43, 67		
Hurua (B)	89		
Hurua (Bo.)	89		
I			
Ichchura-mula (M.)	25	Jadikkay (M.)	42
Ijjul (H)	116	Jafran (H , B & Bo)	35, 91
Ilaikal (M)	88	Jahari sontakka (Bo)	80
Ilayechi (H.)	101	Jalqumbhi (H)	99
Indian Drug Industry (English)	251	Jam (H & B.)	108
Indian foods (English)	251	Jamalgota (H.)	88
Indian or wild Liquorice root (English)	21	Jamana (H.)	72
Indian medicine (English)	251	Jamma (P.)	72
Indian Napellus (English)	22	Jamti-ki-bel (H)	97
Indian Pharmacopoeia (English)	17	Jamun Family (English)	73
Indian Pharmacopoeia Codex (English)	18	Janglibadam (Vern)	40
Indian Pharmacopoeia List (English)	17, 251	Jangli-chichonda (H.)	75
Indian Rubarb (English)	46	Jangli darchini (H.)	103
Indian Sarsaparilla (English)	39	Jangali-kanda (Bo.)	50, 92
Indian winter green (English)	38	Janglimendi (H)	74
Indigenous drugs (English)	251, 252	Janglimudrika (Bo)	64
Indigenons remedies (English)	252	Jangli piyaz (H & B.)	50, 92
Indra varuni (S.)	34, 75	Janglisaru (H.)	103
Indrajau (Bo)	109	Jar Gandapuro (Vern.)	38
Indrayan (H & Bo.)	34, 75	Jar-amla (H)	107
Ingudi (S)	68	Jarul (H & B)	74
Inji (M)	51	Jashti madhu (B & Bo)	39
Insecticides of vegetable origin (English)	252	Jatifalam (S)	42
Ipecac (English)	165, 166	Javanala (S.)	95
Ipecacuanha (English)	166	Jaya (S)	101
Ipomoea (English)	82	Jayanti (H & B.)	100
Iris Family (English)	90	Jayphal (H , B & Bo.)	42
Isabghol (Bo.)	44	Jethimadh (H.)	39
Isabgul (B. Vern.)	19, 44	Jhilla (S.)	105
Isband (B.)	43, 67	Jhinge (B)	40, 75
		Jira (H. & B.)	31, 98
		Jiraka (S.)	98
		Jirkivirai (M.)	40, 82
		Jiunti (P.)	63, 97
		Jivanti (Vern.)	40
		Jonkhmari (H.)	79

	PAGE		PAGE
Jowan (B.)	50	Kalajira (H. & B.)	63, 102
Jowar (H., B. & Bo.)	95	Kalambi (S)	82
Joypal (B.)	88	Kala mucha (B.)	61, 95
Julpai (B)	104	Kalasaka (S)	130
Jungli anarash (B.)	91	Kala tulshi (H & B.)	106
Jun-li-kali-mirchi (Bo.)	49	Kala vala (Bo)	107
K		Kalenjire (Bo)	63
Kabab-chini (H , B & Bo.)	44	Kali-Jhant (H. & B.)	101
Kabiraj (Pers.)	36	Kaliharī (H)	92
Kabra (H. & P.)	97	Kalijiri (Bo.)	32
Kachlora (H & Bo)	72	Kalkora (B)	96
Kachnar (H)	97, 102	Kalikutki (Bo.)	43
Kachura (H. & Bo)	35	Kalmegh (B)	24
Kadamba (S & H)	102	Kalmisak (B.)	82
Kadatathie (M)	84	Kalo bikhmo (Vern)	22, 62
Kadavanchi (Bo)	75, 111	Kalo bikhoma doughi (Vern)	22, 62
Kadukkay-pu (M.)	73	Kalu-kerā (B.)	109
Kafi (B)	77	Kamala (H. & B.)	41
Kahira champa (S.)	112	Kambari (H)	98
Kahoo (H)	98	Kambila (Vern.)	41
Kaiphal (H , B & Bo.)	90, 106	Kamini (B.)	106
Kajra (Bo.)	48, 82	Kamkher (Bo)	117
Kaju (H. & Bo)	69	Kamugu (M)	92
Kakadani (S.)	97	Kanakaphala (S.)	88
Kakadumbura (S)	104	Kanaphata (H)	69
Kakajangha (S , H & B.)	99	Kanchana (S)	49
Kakaphala (S)	64	Kanchkuri (Bo.)	89
Kakatundi (H)	81	Kanchnine (M.)	102
Kakilakshya (S.)	27, 102	Kanchnal (H)	97
Kakkanan (M.)	71	Kanchuri-vayr (M.)	89
Kakkar tamaku (P.)	83	Kanda (S & Bo)	96, 97
Kakkata shingi (M.)	44	Kandari (Vern)	25
Kakkay-kolli-virai (M.)	64	Kanduri-ki-Bel (H.)	34
Kakmachi (S. & B)	84, 108	Kanga (Vern.)	45
Kakmani (H. & B.)	64	Kanghi (H.)	101
Kakra singi (Vern.)	69, 222	Kangunī (S.)	32
Kakra shingi (H. & B.)	44	Kanj (H)	49
Kakumullu (Vern)	70	Kanhera (Bo)	81
Kakurjiwan (H. & B.)	105	Kanka (M.)	94
Kaku-vali (Malay)	114	Kankhina (Bo.)	80
Kaladana (H., B & Bo.)	40, 82	Kankri (Bo.)	75
Kala Dhatura (H. & B.)	83	Kansa (H.)	97
Kakra (Vern.)	44	Kanta (Simla)	64

	PAGE		PAGE
Kantabans (Vern.)	27	Katnim (H.)	106
Kantala (H.)	91	Katphala (M)	90, 106
Kapas (H., B. & Bo.)	104	Katson (Ku.)	73
Kapası (Vern.)	39	Kattalai (M)	24
Kapıla (M.)	41	Kattu-Shiragam (M.)	32
Kapıttıha (Vern.)	104	Kattuvaları (Vern.)	70
Karabı (B)	81	Katuka (S)	43
Karachunai (M)	91	Katuka-roğani (M)	43
Karail (B)	94	Kaundal (Bo.)	75
Karala (B.)	110	Kava (Bo.)	40
Karanja (S , H. & Bo.)	45, 72	Kavalı (Bo.)	39
Karavı (S _r)	81	Kavıtha (H.)	140
Karber (H.)	81	Kayam (Bo.)	38
Karchı (H.)	39	Kela (H. & Bo)	110
Karela (H.)	75, 110	Kelı kadam (B)	77
Karer (H)	65	Kend (B)	98
Karı (Bo & Santh)	65, 109	Kendu (H.)	98
Karıa (P)	65	Ker (Bo.)	101
Karıanag (Bo.)	92	Kesar (Vern.)	35
Karınthuvarı (M)	80, 117	Khadıra (H.)	101
Karıra (S)	65	Khair (H)	101
Karıvana (Bo.)	32	Khair Champa (H. & Bo.)	81, 112
Karkanı (Bo)	105	Khaj-golı-cha-vel (Bo)	113
Karkı (Bo)	81	Khappar kadu (Bo.)	103
Karkata srıngı (S.)	44, 69	Khardı (Bo.)	98
Karla (Bo.)	75	Kharetı (Vern)	48
Karmal (H)	102, 113	Khesarı (H & B)	71
Karmora (Kash)	76	Khesarı dal (Vern.)	61
Karpas (S.)	104	Khınna (H.)	89
Karpo-Karışı (M.)	45	Khıra (H.)	75
Karpura maram (M & Vern.)	38, 73	Khırnı (H)	99
Karnaspota (S)	69	Khorasani ajowan (B.)	40, 83
Karunaıkkızhangı (M)	94	Khorasani-owa (Bo.)	83
Karun Kanam (N)	31	Khorasani-yomam (M.)	40, 83
Karun-shıragam (M)	63	Khour (Nep.)	101
Karu umattai (M.)	83	Khumb (H. & P.)	55
Karval (M.)	65	Khubanı (H)	72
Kashappuvetpalarışı (M.)	39, 80	Khubakallana (H)	104
Kashıra champa (S)	81	Khubazı (H. & Bo)	105
Kashmal (H & P.)	28, 29	Khumbı (H & P.)	55
Kasrı (Vern)	33	Khurasani-ajvayan (H)	40, 83
Kataralı (M.)	32, 80	Khushing (P.)	31
Kat-Karanj (H.)	30	Kıkar (H.)	101
Katkı (H & B)	43	Kılluk-kay-velai (M.)	72

	PAGE		PAGE
Kilmora (H)	28	Krishna dhatura (S)	83
Kinai tihiri (Bo)	70	Krishnajiraka (S)	63
Kindal (Bo.)	110	Kua (Tam)	98
Kinsuk (S)	70	Kuberakshi (S)	30
Kiralu (P)	93	Kuchla (H.)	48, 82
Kiramani owa (Bo)	26, 77	Kuchila (B)	48, 82
Kiraruga (M)	95	Kuchila-lata (H & B)	82
Kirata (S)	48	Kukarchita (B)	99
Kirithi (M)	79	Kukarlata (H.)	75
Kiriwolla (Sinhalese)	105	Kukarnicha (H)	109
Kirmala (Vern & H.)	26, 77	Kukarwele (Bo)	75
Kirmar (H)	86	Kukronda (P)	88
Kirtana (Bo.)	71	Kukseem (B).	100
Kiwach (H.)	106	Kukursunga (B)	109
Kiryat (H)	24, 101	Kul (B)	100
Knapa chettu (M)	50	Kulahala (S)	103
Kodalı (M)	74	Kuljud (H)	94
Koda-todalı (B)	49	Kuliakhara (B)	27
Kodicpalay (M)	37	Kumbhi (H & B)	73
Kodo (H)	95	Kumkuma (S)	35, 91
Kodoa (B)	95	Kunch (B.)	21, 70
Kodra (Vern & Bo)	95	Kundu (Bo.)	80
Kodrava (S)	95	Kungku (H)	38
Koh-tor (Baluchi)	83	Kungumuapu (M)	35, 91
Kokoranj (H)	30	Kungyi (H)	48
Kolamavu (Tam)	99	Kurchi (English & B)	80
Kolhal (Bo.)	103	Kurelo-jangro (Bo)	75
Kolka-phul (B)	49, 81	Kurling (Nepł)	120
Kolumbu (Tam)	105	Kurlinga (Nep)	79
Kobu-Kalli (M)	88	Kushtha (S)	47, 78
Kombupudalai (M)	75	Kusturi (Vern)	255
Konda vaghe (M)	70	Kut (H.)	47, 78
Kondapala (M)	82	Kutaja (S.)	39, 80
Konea-dumbar (H.)	104		
Korattai (M)	75	L	
Kori (B.)	70	Labangaphal (B.)	40
Koshatakı (S)	40, 75	Lablab (H.)	76
Kosum (H)	69	Ladaki-revanda Chini (Bo)	46, 86
Kottak (M.)	78	Laghukarnı (S)	63
Kotteı (M)	110	Lahana-gokhru (Bo)	50, 67
Kovidara (S.)	97	Lahuriya (H.)	99
Kowtı (Vern.)	40	Lajja (S)	105
Krishnachura (B.)	110	Lajalu (H. & Bo.)	105
		Lakh (Bo)	71

	PAGE		PAGE
Lalbachlu (H)	102	Mainphal (H)	77
Lal chitra (H , B. & Bo.)	44, 79	Makal (B.)	75
Lal-indrayan (H)	75	Makai (Bo)	95
Lalpost (H)	64	Makhal (B)	34, 75
Lalsag (H.)	101	Makka-scholar (M.)	95
Lal siris (H)	96	Mako (Bo)	84, 108
Lankasij (B)	88	Makoi (H)	84
Lasan (H & Bo)	23, 91	Malabar nut tree (English)	23
Laskar (P.)	63	Malkanguni (H)	32
Lasuna (S)	23, 91	Mamekh (P)	63
Lata palas (S)	29	Mamira (H.)	35
Laural Family (English)	87	Mamiran (Bo)	49
Lavangalata (S)	40	Manaka (S)	93
Lichens (English)	57	Manattak kali (M)	84
Life-Plant-Family (English)	73	Manchingi (Bo)	84
Lil kathi (Santh)	66	Manda (M)	103
Lily Family (English)	91	Mandara (Bo)	28, 94
Lodh (H , B. & Bo)	49	Mandgay (Bo)	27, 94
Lodhra (S)	49	Mangal (M)	94
Lohari (H)	80	Mango Family (English)	69
Lohori (H)	117	Mangustan (H , B & Bo)	104
Lona (H)	64	Maniphal (H)	45
Long (Vern)	38	Manith (H)	107
Luban (H & B)	102	Manja kadambe (M)	77
Ludduga (M)	49	Manjistha (S & B)	107
Lycopodium (English)	207	Manjit (Bo)	107
M		Mankachu (B)	93
Machni (H.)	94	Mankenda (H)	93
Machutie (B)	86	Mansa-sij (B)	88
Madan (S)	45, 77	Manskhel (Kash)	55
Madar (H)	81	Manya (S)	98
Madder Family (English)	77	Mara-manjal (M.)	35
Maddi-palu (Tel)	101	Maravara (M)	118
Madhu malati (S)	37	Maravara Tsjembu (M)	93
Madhurika (S)	38	Maravuli (M.)	89
Magadhi (S. & Tel.)	98	Marchula (H)	106
Magnolia Family (English)	63	Maricha (S)	107
Magrabu (H.)	39	Marisha (S.)	101
Mahakal (S)	75	Marlu-mutta (M)	51, 78
Mahalimbu (Vern)	31	Marudam-pattai (M.)	90
Mahmira (Bo.)	35	Maruk-kallan-kai (Bo. & M.)	46, 77
Mahanimb (H)	101	Marsada boli (M)	30
Mahogany Family (English)	68	Mashaganı (Tam)	105

	PAGE		PAGE
Masha-Parui (S.)	100	Mulberry Family (English)	89
Mashparui (H)	100	Mullanjakka (S.)	101
Masur (H & S.)	105	Mullu-chitta (Tam)	101
Masuri (B)	105	Mulluvellari (M)	75
Mat (H)	99	Mumugu (M)	25
Matazor (H)	85	Mundiri-kai (M)	69
Maurabikh (Vern)	22, 62	Munditika (S)	78, 100
Mazri (H)	106	Mungas kajur (H)	84
Meetha-tellia (Vern)	62	Mungphali (H)	25
Menphal (B)	46, 77	Munjal (M)	35
Meradu (H)	66	Munjariki (S)	106
Merasingi (H & B)	39	Munni-vayz (M)	45
Meshasingi (S)	39	Murba (B)	100
Methi (H , B & Bo)	120	Murmuria (B)	78
Mexican Poppy (English)	25, 60	Murungi (M)	70
Mezereum Family (English)	87	Musabbar (Vern)	24
Mandanak-kattai (S)	47	Mushabbar (Vern)	92
Milkarana (M)	49	Mushkiara (P)	77
Milk-weed Family (English)	81	Mushrooms (English)	56
Milkwort Family (English)	66	Musna (H)	66
Mimulus Family (English)	84	Musta (S & Bo)	103
Minguta (Bo)	88	Mustard cake (English)	62
Mint (English)	255	Mustard Family (English)	65
Mint Family (English)	85	Mutha (B & H)	103
Mirch (H & P)	83	Muttia-lata (B)	42, 110
Mistletoe Family (English)	87	Myetpye (Burma)	105
Mitha indarjou (H)	109	Mylabris (English)	248
Modi-murukkan (M.)	29	Myrobolan Family (English)	73
Mohra (Vern)	22, 62	Myrtle Family (English)	73
Mohri (Vern)	22		
Mohua (Vern)	41	N	
Mohwa meal (English)	80		
Mohwa Family (English)	80	Nachuta (M)	88
Mooda cotton (M.)	69	Nagadamani (S)	26, 77
Moonseed Family (English)	64	Naga-danti (M)	88
Moravela (H & Bo)	63	Nagamusadi (M.)	82
Morwa (Bo)	100	Nagar Motha (H & B)	103
Moulds (English)	54	Nagar mustaka (S)	103
Muchi-tanki (M)	80	Nagbaia (S)	108
Mugani (H. & B)	99	Nagbail (H)	51
Muhuri (B)	43	Nagchampa (Bo.)	105
Mukajali (H.)	73	Nagdona (B.)	26, 77
Mukuk-rattai (M.)	29	Nagadouna (H.)	26, 77

	PAGE		PAGE
Nagdowan (Bo.)	91	Nimrudı (Bo.)	109
Nagkeshar (S , H & B.)	105	Nırbısı (H & B.)	33
Nahanı khapat (Bo)	101	Nırbamı (M)	27
Najunda (M)	68	Nırgundhı (S & H)	121
Nokchıknı (H)	37	Nırumel-neruppu (M)	74
Nala (B. & H)	114	Nırmulı (Bo)	37, 82
Nala tige (M)	71	Nırunjı (M)	50, 67
Nalı (S)	105	Nıshında (B)	111
Nalıchı bajı (Bo)	82	Nısomalı (S)	86
Nandıbriksha (S & B)	31	Nıpa (S.)	116
Nandru (P.)	83	Noalata (B)	71
Nannarı (M)	39	Nomenclature of Drugs (English)	252
Napalı dhanıa (B)	68	Nona (B)	64
Naranjı (M)	82	Norvishee (H)	73
Naraseja (Bo)	88	Nostoc (English)	53
Narcha (H. & B)	103	Nurma (H)	98
Narel (Bo)	97	Nutmeg Family (English)	86
Nargıs (P)	91	Nux-vomıca Family (English)	82
Nari-vengayam (M.)	50, 92	O	
Narıyal (H.)	97	Oandak (P)	63
Naro (Bo)	74	Olaktambol (Bo)	21
Nata-karanja (B)	30	Olang-karaı (M)	104
Nattu-ıreval-chınnı (M)	46, 86	Oleander Family (English)	80
Navaladı (M)	51	Oman (M)	50
Nayeti (Bo)	38, 88	Opium (English)	43
Nay-palaı	50, 82	Osarı (Bo.)	101
Neem Family (English)	68	Ot (Sant.)	55
Neglı (Bo)	66	Ouplate (Bo)	47, 78
Ner (P)	48, 67	P	
Nettavıl (M)	25, 89	Paas lata (H & B)	29
Nettle Family (English)	89	Pachak (B)	47, 78
Newrang (Bo)	88	Pochch-aı-aları (M)	49, 81
Nıghtshade Family (English)	83	Pacholı (H.)	85
Nıl (H. & Bo)	98	Packur-mul (B)	86
Nıla kadalaı (M)	25	Paddam (H)	72
Nılkattei (H. & P)	82	Padmaka (S)	72
Nıla-nırgundı (S)	105	Padma-kasta (Bo)	72
Nıla vakaı (M)	31	Paharı gandana (H)	110
Nıla vembu (M)	24, 48	Palai (M)	81
Nilika (S)	98	Palandu (S)	96
Nılı-nargandı (H)	105	Palla (M)	99
Nım (Veru., H , B & Bo)	27, 74	Palm Family (English)	92

	PAGE		PAGE
Palarvela (Vern)	29	Penicillium (English)	54
Palas (Vern , H & B)	29, 70	Pepper Family (English)	86
Palupaghel kalung (M)	99	Perala (Bo.)	110
Pamukh (P)	85	Pera-rattai (M)	24
Pan (H , B & Bo)	44, 99	Periya-yelakkay (M)	24
Pandharen-kamal (Bo)	106	Petthri (P.)	40
Pandharphali (P)	88	Peyttumatti (M)	34, 75
Pandhrakura (Bo.)	39, 80	Phala kantan (S)	36
Pandorina (English)	53	Phalsa (H & B.)	98
Pambira (M)	75	Phanasa-alambe (Cutch & Bomb)	55
Panirak (H)	67	Phanasa-alambe (Bo)	101
Panlata (B.)	71	Phanerogams (English)	59
Pan-mauri (B)	38	Phemila (S)	100
Pannax Family (English)	76	Phenila (S)	69, 75, 120
Papai (Bo)	31, 74	Phytolacca Family (English)	85
Papas (P)	98	Pichli (Vern)	23
Papaw Family (English)	74	Pikumkai (M)	40, 75
Papaya (H)	31, 74	Pila-kaner (H & Bo)	49, 81
Papey (B)	31, 74	Pilu (H., S & B)	80
Papita (Vern)	31	Pilvu (Bo.)	80
Pappayi (M)	31, 74	Pindalu (H.)	77
Papra (H)	45	Pindaluka (S)	91
Papri (P)	88	Pine-apple Family (English)	91
Parasikaya (S.)	40	Pine Family (English)	90
Parasi-kaya (S)	83	Pinjari (H)	49
Parusha (S)	98	Pipal (H)	99
Parvar (H)	75	Pippali (S)	99
Pashanbheda (Bo.)	102	Piralu (B)	77
Passion-flower Family (English)	74	Pisa (Bo)	23
Patalagaruda (S)	103	Pitadaru (S)	106
Patha (S)	33	Pithvan (H)	108
Patharchuri (B)	102	Pitz (Kash)	108
Patola (S)	75	Piyaj (B)	96
Patsan (H)	105	Piyaz (H.)	96
Patta karie (S)	88	Plant pigments (English)	252
Pattudu (Tel)	105	Plesmonium (English)	93
Pavakkachedi (M)	75	Plumbago Family (English)	79
Pawal-chhatta (B)	56	Poal-chhatea (B.)	56
Pea Family (English)	70	Podina (H.)	42
Pectin (English)	252	Poisonous Plants (English)	253
Pekarakai (M)	110	Pomegranata Family (English)	74
Penarvalli (M.)	75	Poonam-penmaraiva (Mal.)	100
Pendar (Bo.)	77	Ponnan- kottai (M.)	69

	PAGE		PAGE
Poppy Family (English)	64	Rasna (S., H., B & Bo.)	50
Postakatol (M.)	42, 64	Rasun (B)	23, 91
Potaki (S)	102	Ratna-gandhi (S. & Tam.)	110
Potal (B)	75	Rattanjog (P.)	63
Potari (B)	101	Rattu papillay (M)	78
Prashni (Bo.)	99	Rauwolfia (English)	80
Prime-rose Family (English)	79	Rechanaka (S)	41
Priyangu (S.)	72	Revandchini (H & B)	46, 86
Pudel (M)	75	Rhododendron Family (English)	79
Pudina (Bo., H , S & M.)	42, 105	Rhubarb Family (English)	86
Pudinah (B. & Bo.)	42	Rice Husk (English)	253
Puff-balls (English)	56	Ritha (H., B. & Bo)	69, 100, 120
Pugai-ilai (M.)	83	Riti (Sing)	89
Pulan-ki-zhanga (M)	35	Rohan (H , B. & Bo.)	108
Pu-maram (M)	69	Rohuna (S)	108
Punarnaba (B.)	29	Rosaceae (English)	72
Pungammaram (M.)	72	Rose Family (English)	72
Punnag (B.)	66	Ruba-barik (P)	84
Punnaga (S.)	66	Rudrajata (S)	25, 86
Punnagam (M.)	66	Rudraksha (S)	109
Purpuray timur (Nep.)	121	Rudranti (H & B)	98
Pusha (Vern.)	48	Rue Family (English)	67
Pussar (B.)	109	Rukh-alu (Bo)	93
Pyrethrum (English)	168, 185	Rush Family (English)	92
		Rusts (English)	53
		Rutthraksham (M.)	109
		S	
R			
Racta-vinda-chada (S.)	88	Sabuni (B.)	66
Rati-jaman (H.)	108	Sadab (H)	67
Rajadani (S)	99	Sada dhutura (B)	36, 87
Rajkoshataki (S.)	75	Sadul kou (B.)	81
Rakhalu (Bo)	118	Safed bikh (Vern.)	22
Rakra-posta (S.)	64	Safedchandam (H.)	47
Rakta kanchan (Bo)	97	Safed damar (H.)	100
Ramban (Bo)	108	Safed mushi (H.)	102
Ram dataum (H.)	108	Safed siris (H.)	70
Rametha (Bo.)	114	Safran (Vern.)	35
Ramoha (S.)	110	Sag-angur (H. & Vern)	27, 87
Ramphal (Bo.)	64, 87	Sagaragoth (Bo)	30
Ramsita (M.)	64	Sage Family (English)	85
Ramtulshi (H & B)	119	Sahadevi (S. & H.)	100
Ranghevada (Bo.)	103	Sajnah (H.)	70
Ranparul (Bo.)	75		

	PAGE		PAGE
Saka (S.)	108	Shadhurakkallı (M)	88
Sal (S., H., B. & Bo.)	108	Shakar-al-lighal (P.)	81
Salad (B.)	98	Shakar pitan (H. & P.)	88
Salvadora Family (English)	80	Shallakı (S.)	102
Samudra (M)	73	Shal Parnı (S & Bo)	109
Samudrapad (S.)	73	Shankeshvara (Bo.)	51, 78
Samudraphal (B.)	73	Shatamulı (S & B.)	97
Samudraphala (Bo.)	73	Shatapuspi (S.)	43
Samp-kı-kumb (P)	93	Shatawar (H)	47
Samutrapphullam (M)	73	Shayrang (M)	47, 70
Sandan (H)	71	Shendrı (Bo)	41
Sankhaphulı (H)	97	Sheoia (B)	100
Sankhapuspi (S.)	97	Shera (Bo)	88
Sanıpat (S , H & Bo)	110	Shetapuspa (S.)	43
Sanjna (Vern)	42	Shiajira (H)	31
Sankha pushpi (H)	104	Shialkata (English)	60
Sant (H)	29	Shikha-mulam (S.)	76
Saonf (H)	43	Shimai-azho-vanaı-vıraı (M)	43, 67
Saonjna (Vern)	42	Shimai-shombu (M)	31
Sapodilla Family (English)	80	Shiragam (Tam)	98
Sapta parna (S.)	24	Shırısha (S.)	96
Sarkarei-vallı (M.)	82	Shiru kurunja (M.)	38
Sarapunkha (S.)	72	Shirunarı-vengayam (M.)	47
Sarivan (H.)	109	Shivappu-postaka chedı (M.)	64
Sarphankha (H & Bo.)	72	Shiyappunellı (M)	89
Sarpagandha (S.)	46, 81	Shombu (M.)	38, 43
Sarson (Vern)	65	Shorı (B.)	35
Sasa (B)	75	Shothaghni (S.)	29
Satakuppi (M)	43	Sialkanta (H. & B.)	25, 64
Satap (B & Bo.)	67, 114	Sıj (B.)	88
Satapatri (S)	47	Silajit (Vern.)	253
Satavarı (Bo.)	97	Simainayuruvi (M.)	108
Satı (S)	35	Simla-Kashmal (Vern.)	28
Saya (M)	99	Sındilkodı (M)	49
Sebestan Family (English)	82	Sıora (H.)	100
Sedge Family (English)	94	Sıris (H., B & Bo.)	96
Segun (H. & B.)	108	Sıtama-purgonalu (M.)	36, 82
Sehud (H.)	88	Sıtapalam (M.)	64
Sehund (H.)	88	Sıtaphal (H.)	64
Selupa (M.)	68	Sıttrapaladı (M)	88
Senna (English)	163	Small Caltrops (English)	50
Sesamum Family (English)	84	Smuts (English)	53
Shada-burı (B)	28	Snuhi (S.)	88

	PAGE		PAGE
Soap-nut Family (English)	96	T	
Sobhanjana (S)	70	Tagar (H. & B)	50
Sojna (B)	70	Tagara (S)	50
Soma (S. & Bo.)	82	Tagar-ganthoda (Bo)	50
Somalata (S.)	67, 114	Takadasingi (Bo.)	69
Somlata (H & B)	82	Talim khona (B.)	27
Somaraḡ (S , H & B)	32	Talipanaḡ (M)	92
Sona (B)	106	Talmakhana (H.)	27, 102
Sonf (Bo)	43	Tamak (B)	83
Sonogaravi (Bo)	114	Tamaku (H)	83
Soolpha (B)	43	Taman (B)	74
Sopari (Bo)	25, 92	Tamara valli (S.)	89
Sourabhi-nimba (S)	106	Tamarta (M)	102
Spinach Family (English)	85	Tambaku (Bo.)	83
Spindle-tree Family (English)	68	Tambula (S)	44
Srigala-kantaka (S)	25, 64	Tanba (M.)	97
Sriphal (Vern & S)	23	Tannins (English)	73
Steroid hormones (English)	253	Tarur (H & B)	103
Strophanthus seeds (English)	263	Tatara (H)	73
Subali (H)	88	Tattu dattura (P)	83, 36
Sugandha muricha (S)	44	Tatwen (P.)	26
Sugandha-vacha (B)	24	Tazak-tsun (Kash)	79
Sujna (Bo.)	70	Tea Family (English)	66
Sukasa (S)	75	Teak Family (English)	84
Sukh-darsan (H & B)	91	Tea plant (English)	30
Sukra puspita (S)	92	Tekata siḡ (B)	88
Sultana (H)	66	Tejmal (H)	68
Sumlu (Vern)	28	Telakucha (B)	34
Sundew Family (English)	73	Tel-kodukki (M)	82
Sun-flower Family (English)	77	Tellajonna (M)	95
Supari (H & B)	25, 92	Tella-motuku (M.)	71
Suphadie-khus (H & B.)	47	Temru (Bo)	98
Surati sonnamukaḡ (Vern)	71	Tendu (H & Bo)	80, 104
Surinjan (Vern)	34, 92	Tessul (Bazinali, Bo)	110
Sushavi (S)	75	The Anise (English)	43
Sutr-sowa (B)	106	The Betelnut palm (English)	25
Svetakanchan (S)	97	The Asoka tree (English)	47
Swarnakshira (S)	65	The Barberry (English)	28
Sweet-gale Family (English)	90	The Caraway seed (English)	31
Swet-berela (H & B.)	67	The common Jowar (English)	61
Swet-chandan (S)	47	The Creat (English)	24
Syonaka (S.)	106	The East Indian Screw Tree (English)	39

	PAGE		PAGE
The Flowering Plants (English)	59	Toria (Vern)	65
The greater Cardamon (English)	24	Trayamana (S.)	104
The greater Galangal (English)	24	Tridhara-sehund (H.)	88
The Indian Millet (English)	61	Triputi (S)	71
The Indigenous Drugs of		Truffles (English)	56
India (English)	19	Tsjmul (M)	118
The Lemon Tree (English)	34	Tubah (Malay)	71
The Lime Tree (English)	34	Tudovullay (M)	100
The Lodh tree (English)	49	Tukhmalanga (P)	106
Thelu-Kodi (M)	114	Tulkapyre (Tam)	99
The Mahu tree (English)	40	Tumala (S)	80, 117
The marking nut tree (English)	47	Tumburu (S.)	68
The marsh mint (English)	42	Tun (H)	31
The Morell (English)	56	Tuni (Bo)	31
The Mushroom (English)	55	Tunumaram (M.)	31
The Neem (English)	27		
The Opium (English)	42		U
The Papaya (English)	31		
The Potato (English)	62	Udala (Bo)	115
The Rose (English)	47	Udsalap (H)	63
The sweet flag (English)	22	Udumbara (S)	104
The tru barberry (English)	29	Ughaiputtai (M)	80
The Upas tree (English)	25	Ukshi (Bo)	30
The Wealth of India 1949-53		Ulatkambal (H & B.)	21
(English)	20	Umatai (M.)	36, 83
Thikri (Vernl)	29	Unankodi (Tam)	109
Thimbawmagyi (Burm)	110	Undi (Bo)	66
Thinduka (S)	80, 117	Uravu (Mal)	99
Tholkuri (B)	32	Utangan (Bo.)	29
Thorapimple (Bo)	107	Utaajan (H)	29
Tikhi (Bo)	103	Utarni (Bo. & M.)	36
Tikhur (H & B.)	98	Utran (H)	36
Til (S , H , B. & Bo)	84		
Tilia Kachnag (Vern)	22		V
Tinduka (S.)	104		
Tinis (B)	71	Vachnag (Bo)	22, 62
Tinisasegandun (S.)	71	Vada ganneru (M)	81
Tisi (H & B)	67	Vadam-Kottai (M.)	72
Tita (B)	35	Vahisa (S)	99
Titakunga (B)	37	Vaj (Bo)	22, 93
Tithwan (Vern.)	26	Vajrakantaka (S)	88
Tiwas (Bo.)	71	Vakuchi (S)	45
Torai (H. & Bo.)	40, 75	Vallaimurdu (M.)	73

	PAGE		PAGE
Vallaipundu (M.)	23, 91	Vishamandala (S.)	91
Vallarai (M)	27, 32	Visha-mushti (S.)	48, 82
Val milaku (M.)	44	Vishnugandhi (S.)	104
Valuluwal (M)	32	Vishomangil (M.)	91
Vana-palandam (S)	50, 92	Viralı (M)	-69
Vanatik-tika (S)	120	Vitaraka (B)	100
Vaduka parni (S)	32	Volvox (English)	53
Vansa (S.)	27, 94	Vrischikali (S.)	89
Vantulshi (S)	119	Vuir (Kash)	107
Vardara (Bo)	100	Vulgarly phansamba (English)	55
Varuna (S)	98	W	
Vasaka (S)	23	Wagata (M)	77
Vasanvel (Bo)	97	Walnut Family (English)	90
Vashambu (M.)	23, 93	Walsura (Bo. & M)	68
Vashanavi (M)	22, 62	Wee-chhata (Beng.)	55
Vata (S)	104	White mustard (English)	65
Vegetable Drugs (English)	253	Wormwood (English)	26
Vegetable drug resources (English)	253	St John's Wort Family (English)	66
Vellai-maruda-maram (M)	49	Y	
Vellerku (M)	81	Yam Family (English)	91
Vembu (M.)	74	Yamani (S)	50
Verbena Family (English)	84	Yashti-madhu (S)	39
Verenda (B)	46, 89	Yavanala (S)	95
Vetasa (S)	102	Yebruı (B)	83
Vettılai (M)	44	Yellucheddie (M)	84
Vhaneri (Bo)	85	Yetti (M)	40, 48, 82
Vilayati afsantin (H & Dec)	26, 77	Ysjudemaram (M)	98
Vilayati mehdı (H.)	106	Z	
Vilayeti tamaku (H. & B)	83	Zıra (H)	98
Vilayati-vakhandı (Bo)	81	Zıra (Vern)	31
Visha (S.)	22, 26	Zırışk (Vern)	29
Vishala (S)	75	Zufah (Vern.)	98

